

14.5 Index Notation and the Transformation Law

Outside mathematics, a tensor is usually not a vector in $T_q^p(V)$ at all: it is an array of numbers with some indices written upstairs and some downstairs, together with a rule saying how the array changes when the coordinates change. That older definition is still the working one in physics and in differential geometry, and a reader who meets it after this chapter deserves a dictionary. This section supplies one. We write the components of a tensor, derive — not assert — how they transform under a change of basis, and then prove that the classical definition and ours describe the same objects in finite dimensions. The section ends with the warning the classical definition exists to make: an array of numbers is not automatically a tensor.

14.5.1 Components, upstairs and downstairs

Fix a finite-dimensional V over F with $n = \dim V \geq 1$, an ordered basis $\mathcal{B} = (\mathbf{v}_1, \dots, \mathbf{v}_n)$, and its dual basis $\mathcal{B}^* = (\varphi^1, \dots, \varphi^n)$, characterized by $\varphi^i(\mathbf{v}_j) = \delta_j^i$ (Definition 4.1.3). By Theorem 14.2.5, the n^{p+q} tensors

$$\mathbf{v}_{i_1} \otimes \cdots \otimes \mathbf{v}_{i_p} \otimes \varphi^{j_1} \otimes \cdots \otimes \varphi^{j_q}$$

form a basis of $T_q^p(V)$, so every tensor has unique coordinates in it.

Definition 14.5.1 — Components of a tensor

Let $t \in T_q^p(V)$. The **components of t in the basis $\mathcal{B}$** are the unique scalars $T_{j_1 \dots j_q}^{i_1 \dots i_p} \in F$, for $1 \leq i_1, \dots, i_p, j_1, \dots, j_q \leq n$, with

$$t = \sum T_{j_1 \dots j_q}^{i_1 \dots i_p} \mathbf{v}_{i_1} \otimes \cdots \otimes \mathbf{v}_{i_p} \otimes \varphi^{j_1} \otimes \cdots \otimes \varphi^{j_q},$$

the sum running over all $p + q$ indices from 1 to n . An index written **upstairs** belongs to a V -slot and is called **contravariant**; an index written **downstairs** belongs to a V^* -slot and is called **covariant**.

In words: the position of an index records which kind of slot it labels, so that the shape of a symbol already tells you the type of the tensor. Nothing here is new mathematics; it is the coordinate vector of Theorem 14.2.5, with the indices arranged to be readable.

Two pieces of shorthand come with it. The first is the placement rule: a vector has its coordinates upstairs, $\mathbf{v} = \sum_i v^i \mathbf{v}_i$, while a functional has them downstairs, $\xi = \sum_j \xi_j \varphi^j$ — which is Theorem 4.1.3 (c) written in the new notation. The second is the **summation convention**: an index appearing once upstairs and once downstairs in a product is summed over, and the $\sum$ is not written. With it, the displayed formula above reads

$$t = T_{j_1 \dots j_q}^{i_1 \dots i_p} \mathbf{v}_{i_1} \otimes \cdots \otimes \varphi^{j_q},$$

and the evaluation of a functional on a vector is $\xi(\mathbf{v}) = \xi_i v^i$. The rest of this book writes its sums out; this section is the one place where the convention is used, and it is used because the transformation law is unreadable without it.

Examples.

1. **Type (1, 1): an operator.** If $t \in T_1^1(V)$ corresponds to $S \in \mathcal{L}(V)$ under Theorem 14.3.9, then T_j^i is the (i, j) entry of $[S]_{\mathcal{B}}^{\mathcal{B}}$: the tensor $\mathbf{v}_i \otimes \varphi^j$ is the operator $\mathbf{u} \mapsto \varphi^j(\mathbf{u})\mathbf{v}_i$, whose matrix is $\mathbf{E}_{ij}$. Upper index = row, lower index = column.
2. **Type (0, 2): a bilinear form.** If $t \in T_2^0(V)$ corresponds to the bilinear form β (Proposition 14.4.3 and Exercise (B3: Which statements hold?)), then $T_{ij} = \beta(\mathbf{v}_i, \mathbf{v}_j)$, which is the Gram matrix $[\beta]_{\mathcal{B}}$ of Definition 13.1.2. Both indices downstairs.
3. **Contraction.** In components, Definition 14.4.3 is “set an upper index equal to a lower one and sum”. For $t \in T_1^1(V)$,

$$C_1^1(t) = T_i^i \quad \left(= \sum_{i=1}^n T_i^i \right),$$

which is the trace, by Theorem 14.4.1. This is the whole reason the convention pairs an upper with a lower index: a repeated index is a contraction, and contraction is the only basis-free way to remove slots.

Remark. The names are historically backwards, and it is worth saying so once. A *contravariant* index sits upstairs, on a V -slot, and its components transform by the **inverse** of the matrix that transforms the basis vectors; a *covariant* index sits downstairs and transforms by the same matrix as the basis vectors. The names describe the components, not the slots. [Theorem 14.5.2](#) below is what they refer to.

14.5.2 How the components change

Let $\mathcal{B} = (\mathbf{v}_1, \dots, \mathbf{v}_n)$ and $\mathcal{B}' = (\mathbf{v}'_1, \dots, \mathbf{v}'_n)$ be two ordered bases, and let

$$\mathbf{P} = [\text{id}]_{\mathcal{B}'}^{\mathcal{B}} = (p_j^i)$$

be the change-of-coordinates matrix from $\mathcal{B}'$ to $\mathcal{B}$ (Definition 3.7.1), whose j -th column is $[\mathbf{v}'_j]_{\mathcal{B}}$. That is,

$$\mathbf{v}'_j = \sum_{i=1}^n p_j^i \mathbf{v}_i,$$

with the row index upstairs and the column index downstairs. Write $\mathbf{Q} = \mathbf{P}^{-1} = (q_j^i)$, which exists because a change-of-coordinates matrix is invertible (Proposition 3.7.2).

The dual bases move by the inverse matrix. This is the one computation the whole section rests on, so we do it first.

Lemma 14.5.1 — Dual Bases Move by the Inverse Matrix

With the notation above, $(\varphi')^k = \sum_{l=1}^n q_l^k \varphi^l$ for each k , where $(\varphi')^1, \dots, (\varphi')^n$ is the dual basis of $\mathcal{B}'$ supplied by Theorem 4.1.3.

Proof. Both sides are linear maps on V , so by the uniqueness clause of Theorem 3.1.5 it is enough to check that they agree at each vector of the basis $\mathcal{B}'$. By Definition 4.1.3 the left side gives $(\varphi')^k(\mathbf{v}'_j) = \delta_j^k$. The right side gives

$$\sum_l q_l^k \varphi^l \left(\sum_i p_j^i \mathbf{v}_i \right) = \sum_l \sum_i q_l^k p_j^i \delta_i^l = \sum_i q_i^k p_j^i,$$

which is the (k, j) entry of $\mathbf{QP} = \mathbf{I}_n$, that is δ_j^k . The two agree. ■

So the dual basis vectors carry the entries of $\mathbf{P}^{-1}$, while the basis vectors carry the entries of $\mathbf{P}$. Substituting both into the expansion of a tensor gives the transformation law.

Theorem 14.5.2 — Transformation Law for Components

Let $t \in T_q^p(V)$ have components $T_{j_1 \dots j_q}^{i_1 \dots i_p}$ in $\mathcal{B}$ and $(T')_{l_1 \dots l_q}^{k_1 \dots k_p}$ in $\mathcal{B}'$, and let $\mathbf{P} = (p_j^i)$ and $\mathbf{Q} = \mathbf{P}^{-1} = (q_j^i)$ be as above. Then, with the summation convention,

$$(T')_{l_1 \dots l_q}^{k_1 \dots k_p} = q_{i_1}^{k_1} \dots q_{i_p}^{k_p} \times p_{l_1}^{j_1} \dots p_{l_q}^{j_q} T_{j_1 \dots j_q}^{i_1 \dots i_p}.$$

Each **upper** index is contracted with a factor of $\mathbf{P}^{-1}$, each **lower** index with a factor of $\mathbf{P}$.

Idea. There is only one move: write t in the new basis, replace every new basis vector and every new dual basis vector by its expression in the old ones, multiply out using multilinearity of $\otimes$, and compare with the expansion of t in the old basis. Coordinates in a basis are unique, so the comparison is an equality of coefficients. Doing it once for type $(1, 0)$ shows the whole pattern; the general case is the same computation with more indices.

Proof. Start with the expansion of t in the new basis:

$$t = (T')_{l_1 \dots l_q}^{k_1 \dots k_p} \mathbf{v}'_{k_1} \otimes \dots \otimes \mathbf{v}'_{k_p} \otimes (\varphi')^{l_1} \otimes \dots \otimes (\varphi')^{l_q}.$$

Substitute $\mathbf{v}'_k = p_k^i \mathbf{v}_i$ and, by Lemma 14.5.1, $(\varphi')^l = q_j^l \varphi^j$. Since $\otimes$ is linear in each of its slots, the scalars may be pulled out of all $p + q$ factors at once, giving

$$t = (T')_{l_1 \dots l_q}^{k_1 \dots k_p} p_{k_1}^{i_1} \dots p_{k_p}^{i_p} q_{j_1}^{l_1} \dots q_{j_q}^{l_q} \times \mathbf{v}_{i_1} \otimes \dots \otimes \mathbf{v}_{i_p} \otimes \varphi^{j_1} \otimes \dots \otimes \varphi^{j_q}.$$

The tensors on the right form a basis of $T_q^p(V)$ (Theorem 14.2.5), so coordinates in them are unique (Theorem 1.5.1). Comparing with Definition 14.5.1,

$$T_{j_1 \dots j_q}^{i_1 \dots i_p} = p_{k_1}^{i_1} \dots p_{k_p}^{i_p} q_{j_1}^{l_1} \dots q_{j_q}^{l_q} (T')_{l_1 \dots l_q}^{k_1 \dots k_p}. \quad (*)$$

It remains to invert $(*)$. Contract both sides of $(*)$ with $q_{i_1}^{m_1} \dots q_{i_p}^{m_p} p_{r_1}^{j_1} \dots p_{r_q}^{j_q}$ and sum over $i_1, \dots, i_p, j_1, \dots, j_q$. On the right, each pair $q_i^m p_k^j$ sums to $(\mathbf{QP})_k^m = \delta_k^m$, and each pair $q_j^l p_r^j$ sums to δ_r^l ; every delta then collapses one summation. What survives on the right is $(T')_{r_1 \dots r_q}^{m_1 \dots m_p}$, and what stands on the left is the asserted expression. Renaming m to k and r to l gives the theorem. ■

The two smallest cases are results the book already has, and they are worth reading off as a

check.

- **Type (1, 1).** The law reads $(T')_l^k = q_i^k p_l^j T_j^i$, that is $\mathbf{T}' = \mathbf{P}^{-1} \mathbf{T} \mathbf{P}$. This is Theorem 3.7.3: matrices of an operator in two bases are **similar**.
- **Type (0, 2).** The law reads $(T')_{kl} = p_k^i p_l^j T_{ij}$, that is $\mathbf{T}' = \mathbf{P}^\top \mathbf{T} \mathbf{P}$. This is Theorem 13.2.1: Gram matrices of a bilinear form in two bases are **congruent**.

So similarity and congruence are not two unrelated relations that happen to look alike. They are the transformation laws of type (1, 1) and type (0, 2), and the difference between $\mathbf{P}^{-1}$ and $\mathbf{P}^\top$ is exactly the difference between an upper and a lower index.

14.5.3 The classical definition

The classical definition turns the theorem into a definition: a tensor *is* a rule producing arrays that obey the law.

Definition 14.5.2 — Classical tensor

Let V be a vector space over F with $\dim V = n \geq 1$, and let $p, q \geq 0$. A **classical tensor of type (p, q)** on V is a rule $\mathcal{B} \mapsto A_{\mathcal{B}}$ assigning to **every** ordered basis $\mathcal{B}$ of V an array

$$A_{\mathcal{B}} = (A_{j_1 \dots j_q}^{i_1 \dots i_p})$$

of n^{p+q} scalars, such that for **any two** ordered bases $\mathcal{B}, \mathcal{B}'$, with $\mathbf{P} = [\text{id}]_{\mathcal{B}'}^{\mathcal{B}}$ and $\mathbf{Q} = \mathbf{P}^{-1}$,

$$(A_{\mathcal{B}'})_{l_1 \dots l_q}^{k_1 \dots k_p} = q_{i_1}^{k_1} \cdots q_{i_p}^{k_p} \times p_{l_1}^{j_1} \cdots p_{l_q}^{j_q} (A_{\mathcal{B}})_{j_1 \dots j_q}^{i_1 \dots i_p}.$$

Theorem 14.5.3 — The Two Definitions Agree

Let V be finite-dimensional over F with $\dim V = n \geq 1$, and let $p, q \geq 0$. The map sending $t \in T_q^p(V)$ to the rule

$$\mathcal{B} \longmapsto (\text{components of } t \text{ in } \mathcal{B})$$

is a bijection from $T_q^p(V)$ onto the set of classical tensors of type (p, q) on V . It is linear for the entrywise operations on arrays.

Idea. Well defined is [Theorem 14.5.2](#). Injective because a tensor is determined by its coordinates in one basis. Surjective is the only step with content, and it is short: fix one basis $\mathcal{B}_0$, build the tensor whose components there are the given array, and then observe that its components in any other basis obey the same law as the rule's array does, starting from the same array — so the two agree.

Proof. Well defined. If $t \in T_q^p(V)$, its components in the various bases satisfy the required identity, by [Theorem 14.5.2](#). So the rule attached to t is a classical tensor.

Injective. Fix a basis $\mathcal{B}_0$. If two tensors give the same rule, they have the same components in $\mathcal{B}_0$, hence are equal, since coordinates in the basis of Theorem 14.2.5 are

unique (Theorem 1.5.1).

Surjective. Let $\mathcal{B} \mapsto A_{\mathcal{B}}$ be a classical tensor. Fix an ordered basis $\mathcal{B}_0 = (\mathbf{v}_1, \dots, \mathbf{v}_n)$ with dual basis $(\varphi^1, \dots, \varphi^n)$, and define

$$t := (A_{\mathcal{B}_0})^{i_1 \dots i_p}_{j_1 \dots j_q} \mathbf{v}_{i_1} \otimes \dots \otimes \varphi^{j_q} \in T_q^p(V),$$

so that the components of t in $\mathcal{B}_0$ are exactly $A_{\mathcal{B}_0}$. Let $\mathcal{B}'$ be any other ordered basis and put $\mathbf{P} = [\text{id}]_{\mathcal{B}'}^{\mathcal{B}_0}$. By [Theorem 14.5.2](#) the components of t in $\mathcal{B}'$ are obtained from $A_{\mathcal{B}_0}$ by the law with this $\mathbf{P}$; by [Definition 14.5.2](#), $A_{\mathcal{B}'}$ is obtained from $A_{\mathcal{B}_0}$ by the law with the same $\mathbf{P}$. Both are the value of the same expression, so they are equal. Hence the rule attached to t is the given one.

Linearity. The components of $t + ct'$ in each basis are $A + cA'$, because coordinate maps are linear (Corollary 3.5.5). ■

Why the older definition survives. It is tempting to file [Definition 14.5.2](#) as a historical curiosity, and that would be a mistake. Three honest reasons keep it in use.

- **Components are what gets measured and computed.** A stress tensor, a metric, an inertia tensor: the physicist has numbers in a chosen frame, and the question that matters is what those numbers become in another frame. The transformation law is not a consequence to be derived afterwards; it is the datum.
- **There is often no global basis to be abstract about.** On a curved space, the objects of interest are tensor *fields*, and a manifold generally admits no single coordinate system covering it. What it does admit is many local coordinate patches with prescribed rules for passing between them. In that setting “an array in each chart, transforming correctly on overlaps” is a definition that works, while “an element of $T_q^p(V)$ ” needs a V that varies from point to point, which is a later and heavier construction.
- **It is a usable test.** Given a recipe that produces an array from a basis, checking the transformation law is a finite computation, and it is the standard way to certify that a quantity is geometric rather than an artifact of the coordinates. The next warning is exactly such a test, failed.

14.5.4 An array is not a tensor

Warning. **A matrix is not automatically a tensor.** Consider the rule that assigns to **every** ordered basis of V the same array, written as a matrix,

$$\mathbf{A}_{\mathcal{B}} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad \dim V = 2,$$

read as a type $(1, 1)$ array. Take $\mathcal{B} = (\mathbf{v}_1, \mathbf{v}_2)$ and $\mathcal{B}' = (\mathbf{v}_1, \mathbf{v}_1 + \mathbf{v}_2)$, so that

$$\mathbf{P} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad \mathbf{P}^{-1} = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}.$$

The law demands $\mathbf{A}_{\mathcal{B}'} = \mathbf{P}^{-1} \mathbf{A}_{\mathcal{B}} \mathbf{P}$, which is $\begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}$, not $\mathbf{A}_{\mathcal{B}}$. So this rule is **not** a classical tensor of type $(1, 1)$: it is a matrix, and nothing more. The operator it describes

in $\mathcal{B}$ — projection onto $\text{span}(\mathbf{v}_1)$ along $\text{span}(\mathbf{v}_2)$ — is a perfectly good tensor, but then its array in $\mathcal{B}'$ is the second matrix, not the first. “Write down the same numbers in every basis” is a rule, and almost never a tensor.

Two refinements of the same point are worth having.

The same array can be two different tensors. Take the array $\mathbf{I}_2$. As a type $(1,1)$ rule it *is* a tensor: $\mathbf{P}^{-1}\mathbf{I}_2\mathbf{P} = \mathbf{I}_2$ for every $\mathbf{P}$, and the tensor is the identity operator, which is the ι of Exercise (C2: The contraction of the identity, over any field). As a type $(0,2)$ rule, the same constant array is **not** a tensor: the law demands $\mathbf{P}^\top\mathbf{I}_2\mathbf{P}$, and for the $\mathbf{P}$ above that is $\begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \neq \mathbf{I}_2$. Geometrically: “the identity operator” is basis-free, while “the bilinear form whose Gram matrix is the identity”, that is, the choice of an inner product declaring the basis orthonormal, is not — it depends on the basis, and orthonormality, as Chapter 10 defines it, is relative to a chosen inner product.

Warning. Which type an array has is not visible in the array. The array $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ transforms as $\mathbf{P}^{-1}\mathbf{A}\mathbf{P}$ if it is the matrix of an operator and as $\mathbf{P}^\top\mathbf{A}\mathbf{P}$ if it is the Gram matrix of a form, and for the $\mathbf{P}$ above those give $\begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}$ and $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$. Same numbers, same basis change, different answers. This is why index position is not decoration: writing A_j^i rather than A_{ij} is a statement about what the object is.

Quick check. Let $\dim V = 2$. Is the rule assigning to every ordered basis the array $A_{ij} = \delta_{ij}$ of type $(0,2)$ ever a classical tensor — say, over $\mathbb{F}_2$, or if we only allow bases obtained from one another by permutation? Decide, with a reason.

Solution. Over $\mathbb{F}_2$ it still fails: with $\mathbf{P} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$, the law demands $\mathbf{P}^\top\mathbf{I}_2\mathbf{P} = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$ (since $2 = 0$ in $\mathbb{F}_2$), which is not $\mathbf{I}_2$. If the bases are restricted to permutations of one fixed basis, then $\mathbf{P}$ is a permutation matrix, so $\mathbf{P}^\top = \mathbf{P}^{-1}$ and $\mathbf{P}^\top\mathbf{I}_2\mathbf{P} = \mathbf{I}_2$: the rule survives. But Definition 14.5.2 quantifies over **every** ordered basis, and a rule that works only for a restricted family is exactly the kind of coordinate-dependent object the definition is designed to exclude. The lesson is that the law is a demand about all changes of basis, not about some.

14.5.5 Exercises

14.5.5.1 A. Check your understanding

Exercise A1.

- Define the components of a tensor of type (p, q) in an ordered basis, and say which indices go upstairs.
- State the summation convention, and write $\xi(\mathbf{v})$ with it.
- For a type $(1,1)$ tensor, write out the transformation law and name the relation it expresses between the two matrices.
- Give the transformation law for type $(0,2)$ and name the relation.

- (e) True or false: any $n \times n$ array of scalars, attached to every basis, is a tensor of type $(1, 1)$. Justify your answer.

Solution.

- (a) They are the unique coordinates of t in the basis of Theorem 14.2.5 built from $\mathcal{B}$ and $\mathcal{B}^*$ (Definition 14.5.1). An index belonging to a V -slot goes upstairs; one belonging to a V^* -slot goes downstairs.
- (b) An index occurring once upstairs and once downstairs in a product is summed from 1 to n , with the summation sign suppressed. So $\xi(\mathbf{v}) = \xi_i v^i$.
- (c) $(T')_l^k = q_i^k p_l^j T_j^i$, that is $\mathbf{T}' = \mathbf{P}^{-1} \mathbf{T} \mathbf{P}$: the two matrices are **similar** (Theorem 3.7.3).
- (d) $(T')_{kl} = p_k^i p_l^j T_{ij}$, that is $\mathbf{T}' = \mathbf{P}^\top \mathbf{T} \mathbf{P}$: the two matrices are **congruent** (Theorem 13.2.1, Definition 13.2.1).
- (e) False. The rule must satisfy the transformation law; the constant rule $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ does not, as the warning above shows.

14.5.5.2 B. Practice

Exercise B1: Transforming components.

Let V be a 2-dimensional space over $\mathbb{Q}$ with basis $\mathcal{B} = (\mathbf{v}_1, \mathbf{v}_2)$, and let $\mathcal{B}' = (\mathbf{v}_1 + \mathbf{v}_2, \mathbf{v}_1 + 2\mathbf{v}_2)$.

- (a) Write down $\mathbf{P} = [\text{id}]_{\mathcal{B}'}^{\mathcal{B}}$ and $\mathbf{Q} = \mathbf{P}^{-1}$.
- (b) Let $t \in T_1^1(V)$ have components $T_1^1 = 1$, $T_2^1 = 2$, $T_1^2 = 3$, $T_2^2 = 4$ in $\mathcal{B}$. Find its components in $\mathcal{B}'$.
- (c) Let $s \in T_2^0(V)$ have the same array S_{ij} of components in $\mathcal{B}$. Find its components in $\mathcal{B}'$, and compare with (b).
- (d) Compute $C_1^1(t)$ in both bases and check that the two answers agree.

Solution.

- (a) The columns of $\mathbf{P}$ are the $\mathcal{B}$ -coordinates of the vectors of $\mathcal{B}'$, so

$$\mathbf{P} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}, \quad \mathbf{Q} = \mathbf{P}^{-1} = \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix},$$

using Theorem 0.9.8 with $\det \mathbf{P} = 1$.

- (b) By Theorem 14.5.2, $\mathbf{T}' = \mathbf{P}^{-1} \mathbf{T} \mathbf{P}$. With $\mathbf{T} = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$,

$$\mathbf{P}^{-1} \mathbf{T} = \begin{pmatrix} -1 & 0 \\ 2 & 2 \end{pmatrix}, \quad \mathbf{T}' = \begin{pmatrix} -1 & -1 \\ 4 & 6 \end{pmatrix}.$$

- (c) Now the law gives $\mathbf{S}' = \mathbf{P}^\top \mathbf{S} \mathbf{P}$. With $\mathbf{P}^\top = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$ (this $\mathbf{P}$ is symmetric),

$$\mathbf{P}^\top \mathbf{S} = \begin{pmatrix} 4 & 6 \\ 7 & 10 \end{pmatrix}, \quad \mathbf{S}' = \begin{pmatrix} 10 & 16 \\ 17 & 27 \end{pmatrix}.$$

The same array in the same two bases produces completely different answers, because the two objects are of different types.

- (d) In $\mathcal{B}$: $T_1^1 + T_2^2 = 1 + 4 = 5$. In $\mathcal{B}'$: $-1 + 6 = 5$. They agree, as they must, since Definition 14.4.3 is defined without reference to a basis. (By contrast, the “trace” of the type $(0, 2)$ arrays is $1 + 4 = 5$ against $10 + 27 = 37$: there is no contraction of two lower indices, and summing S_{ii} is not a basis-free operation.)

Exercise B2: Tensor or not?.

Let $\dim V = 2$. For each rule below, determine whether it is a classical tensor of the stated type. Justify your answer, using $\mathcal{B} = (\mathbf{v}_1, \mathbf{v}_2)$ and $\mathcal{B}' = (\mathbf{v}_2, \mathbf{v}_1)$ or $\mathcal{B}' = (\mathbf{v}_1, \mathbf{v}_1 + \mathbf{v}_2)$ where a counterexample is needed.

- (a) Type $(1, 1)$: $A_{\mathcal{B}} = \mathbf{I}_2$ for every $\mathcal{B}$.
- (b) Type $(1, 1)$: $A_{\mathcal{B}} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ for every $\mathcal{B}$.
- (c) Type $(0, 2)$: $A_{\mathcal{B}} = \mathbf{0}$ for every $\mathcal{B}$.
- (d) Type $(1, 1)$: $A_{\mathcal{B}} = (\text{tr } \mathbf{M}_{\mathcal{B}}) \mathbf{I}_2$, where $\mathbf{M}_{\mathcal{B}}$ is the matrix in $\mathcal{B}$ of one fixed operator $M \in \mathcal{L}(V)$.

Solution.

- (a) Yes. The law demands $\mathbf{P}^{-1} \mathbf{I}_2 \mathbf{P} = \mathbf{I}_2$, which holds for every invertible $\mathbf{P}$. The tensor is id_V .
- (b) No. With $\mathcal{B}' = (\mathbf{v}_2, \mathbf{v}_1)$ we get $\mathbf{P} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \mathbf{P}^{-1}$, and $\mathbf{P}^{-1} A \mathbf{P} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \neq A$. (This is the standard non-diagonalizable matrix of Chapter 8; here it is failing a different test.)
- (c) Yes. The law reads $\mathbf{0} = \mathbf{P}^\top \mathbf{0} \mathbf{P}$, which holds. The tensor is the zero tensor, and the zero array is the one array that is a tensor of every type.
- (d) Yes. By Theorem 3.7.5 the scalar $c = \text{tr } \mathbf{M}_{\mathcal{B}}$ is the same for every $\mathcal{B}$, so the rule is the constant rule $c \mathbf{I}_2$, and $\mathbf{P}^{-1} (c \mathbf{I}_2) \mathbf{P} = c \mathbf{I}_2$. The tensor is $c \text{id}_V$. The point of contrast with (b) is that c was itself built basis-freely.

14.5.5.3 C. Going deeper

Exercise C1: Contraction in indices.

Let $\dim V = n$ and let $t \in T_1^1(V)$ have components T_j^i in $\mathcal{B}$ and $(T')_l^k$ in $\mathcal{B}'$.

- (a) Prove directly from Theorem 14.5.2 that $(T')_k^k = T_i^i$, without using Theorem 14.4.1.
- (b) Let $t \in T_1^2(V)$ with components T_k^{ij} . Show that $U^j := T_i^{ij}$ are the components of

a type $(1, 0)$ tensor, and identify it in terms of Definition 14.4.3.

- (c) Let $s \in T_0^2(V)$ have components S^{ij} . Show that the scalar $\sum_i S^{ii}$ is **not** basis-independent, and say which step of (b) fails for it.

Solution.

- (a) By the law, $(T')_l^k = q_i^k p_l^j T_j^i$. Setting $l = k$ and summing,

$$(T')_k^k = \sum_k \sum_{i,j} q_i^k p_k^j T_j^i = \sum_{i,j} \left(\sum_k p_k^j q_i^k \right) T_j^i = \sum_{i,j} \delta_i^j T_j^i = \sum_i T_i^i,$$

where the inner sum is the (j, i) entry of $\mathbf{P}\mathbf{Q} = \mathbf{I}_n$.

- (b) By Theorem 14.5.2 for type $(2, 1)$, $(T')_l^{km} = q_i^k q_j^m p_l^r T_r^{ij}$. Setting $l = k$ and summing over k ,

$$(U')^m = \sum_k (T')_k^{km} = q_j^m \left(\sum_{k,i,r} q_i^k p_k^r T_r^{ij} \right) = q_j^m \delta_i^r T_r^{ij} = q_j^m U^j,$$

which is the transformation law for a single upper index. So (U^j) is a classical tensor of type $(1, 0)$, and by Theorem 14.5.3 it is a vector; it is $C_1^1(t)$, the contraction of the first upper slot against the lower one (Definition 14.4.3).

- (c) Take $n = 2$ and $s = \mathbf{v}_1 \otimes \mathbf{v}_1$, so $S^{11} = 1$ and the other components vanish; then $\sum_i S^{ii} = 1$. Let $\mathcal{B}' = (\mathbf{v}_1 + \mathbf{v}_2, \mathbf{v}_2)$, so that $\mathbf{P} = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$ and $\mathbf{v}_1 = \mathbf{v}'_1 - \mathbf{v}'_2$. Expanding,

$$s = (\mathbf{v}'_1 - \mathbf{v}'_2) \otimes (\mathbf{v}'_1 - \mathbf{v}'_2),$$

so $(S')^{11} = (S')^{22} = 1$ and $(S')^{12} = (S')^{21} = -1$. Hence $\sum_k (S')^{kk} = 2 \neq 1$. The step that fails is the collapse in (b): setting the two indices equal pairs a factor q_i^k with another factor q_j^k , and $\sum_k q_i^k q_j^k$ is an entry of $\mathbf{Q}^\top \mathbf{Q}$, not of $\mathbf{P}\mathbf{Q} = \mathbf{I}_n$, so nothing cancels. Only an upper index repeated against a **lower** one is a contraction.

Exercise C2: Raising an index needs a form.

Let V be finite-dimensional over F and let β be a non-degenerate bilinear form on V , with components $g_{ij} = \beta(\mathbf{v}_i, \mathbf{v}_j)$ in $\mathcal{B}$.

- (a) Prove that the matrix $\mathbf{G} = (g_{ij})$ is invertible, and let (g^{ij}) denote the entries of $\mathbf{G}^{-1}$.
- (b) Let $t \in T_1^0(V)$ have components T_j . Show that $U^i := g^{ij} T_j$ satisfies the type $(1, 0)$ transformation law, so that “raising the index with β ” produces a genuine tensor.
- (c) Explain in one or two sentences why this does **not** contradict the warning that there is no canonical map $V^* \rightarrow V$.

Solution.

- (a) By Proposition 13.1.2, β is non-degenerate exactly when $\mathbf{G} = [\beta]_{\mathcal{B}}$ is invertible.
- (b) By Theorem 13.2.1, $\mathbf{G}' = \mathbf{P}^\top \mathbf{G} \mathbf{P}$, so $(\mathbf{G}')^{-1} = \mathbf{P}^{-1} \mathbf{G}^{-1} (\mathbf{P}^\top)^{-1} = \mathbf{Q} \mathbf{G}^{-1} \mathbf{Q}^\top$, that is $(g')^{kl} = q_i^k q_j^l g^{ij}$: the inverse array carries two upper indices. Also $(T')_l = p_l^r T_r$

by Theorem 14.5.2. Hence

$$\begin{aligned}(U')^k &= (g')^{kl}(T')_l = q_i^k g^{ij} \left(\sum_l q_j^l p_l^r \right) T_r \\ &= q_i^k g^{ij} \delta_j^r T_r = q_i^k g^{ij} T_j = q_i^k U^i,\end{aligned}$$

using $\sum_l p_l^r q_j^l = (\mathbf{PQ})_j^r = \delta_j^r$. That is the type $(1, 0)$ law.

- (c) The map is canonical *given* β , not given V alone: a different non-degenerate form raises indices differently, and there is no distinguished choice. This is the same point as the warning after Definition 14.4.1: the isomorphism $V \cong V^*$ is extra structure, which Chapter 10 supplies with an inner product and Chapter 13 with a form.