1.5 The Unit Circle and Polar Form

Addition in $\mathbb{C}$ is addition of arrows in the plane: $(a, b) + (c, d) = (a + c, b + d)$ puts the second arrow at the tip of the first. Multiplication has a picture too, and Section 1.2 already holds half of it: the modulus of a product is the product of the moduli (Theorem 1.2.2 (d)). The other half is about angles, and angles need the functions cosine and sine, which this volume does not construct. This section first proves everything about directions that needs no angle at all, then states exactly five facts about cosine and sine that it borrows, and from those five builds the polar form of a complex number and the rule "multiply the lengths, add the angles".

1.5.1 Length and direction

Every non-zero complex number should be a length times a direction. The length is the modulus. The directions are the complex numbers of length 1, and they deserve a name.

The unit circle is the set of complex numbers at distance 1 *from* 0.

We write

$$\mathbb{T} := \{u \in \mathbb{C} : |u| = 1\}$$

for the **unit circle**. For $u = x + yi$ with $x, y \in \mathbb{R}$, the condition $|u| = 1$ says $\sqrt{x^2 + y^2} = 1$, that is, $x^2 + y^2 = 1$: both sides of the first equation are non-negative, so it holds exactly when the squares agree (Number Systems: From \(\nN\) to \(\nR\), Proposition 2.4.3 (g)). So $\mathbb{T}$ is the circle of radius 1 about 0 in the plane. It contains 1, -1, i and $-i$, and also $\frac{3}{5} + \frac{4}{5}i$, since $\frac{9}{25} + \frac{16}{25} = 1$. It does not contain $1 + i$, whose modulus is $\sqrt{2} \neq 1$.

The next proposition collects what the multiplication of $\mathbb{C}$ does on $\mathbb{T}$. None of it uses an angle.

Proposition 1.5.1. The Unit Circle

(a) $1 \in \mathbb{T}$. If $u, v \in \mathbb{T}$, then $uv \in \mathbb{T}$. If $u \in \mathbb{T}$, then $u \neq 0$ and $u^{-1} = \overline{u} \in \mathbb{T}$.

(b) For every $z \in \mathbb{C}$ with $z \neq 0$ there is **exactly one** pair (r, u) with $r \in \mathbb{R}$, $r > 0$, $u \in \mathbb{T}$ and $z = ru$, namely $r = |z|$ and $u = z/|z|$.

(c) For $z \in \mathbb{C}$ with $z \neq 0$: $1/z = \overline{z}$ if and only if $|z| = 1$.

Idea. Every part is a statement about moduli, so each follows from the rules of Section 1.2: moduli multiply, $u\overline{u} = |u|^2$, and only 0 has modulus 0. For (b), divide z by its length; for the uniqueness, take moduli of $z = ru$.

Proof.

(a) By Proposition 1.2.1, the modulus of the real number 1 is its absolute value, which is 1; so $1 \in \mathbb{T}$. If $|u| = |v| = 1$, then $|uv| = |u||v| = 1$ by Theorem 1.2.2 (d). Let $u \in \mathbb{T}$. Then $u \neq 0$, since $|0| = 0 \neq 1$ (Theorem 1.2.2 (e)). By Theorem 1.2.2 (c), $u\overline{u} = |u|^2 = 1$, so $\overline{u}$ satisfies the defining equation of u^{-1}. Since $\mathbb{C}$ is a field (Theorem 1.2.4), inverses are unique (Number Systems: From \(\nN\) to \(\nR\), Lemma 4.1.1 (a)), and $u^{-1} = \overline{u}$. Finally $|\overline{u}| = |u| = 1$ by Theorem 1.2.2 (d).

(b) *Existence.* Let $r := |z|$. Then $r \geq 0$, and $r \neq 0$ because $z \neq 0$ (Theorem 1.2.2 (e)); so $r > 0$, and its inverse r^{-1} is a positive real number (Number Systems: From \(\nN\)

to \(\nR\), Proposition 3.2.2 (a)). By Theorem 1.2.4 the inverse of r in $\mathbb{C}$ is this real number, so $u := z/|z| = r^{-1}z$ is defined, and $z = ru$. By Theorem 1.2.2 (d) and Proposition 1.2.1, $|u| = |r^{-1}||z| = r^{-1}r = 1$, where $|r^{-1}| = r^{-1}$ because the absolute value of a positive real number is the number itself (Number Systems: From \(\nN\) to \(\nR\), Definition 4.1.1). So $u \in \mathbb{T}$.
Uniqueness. Let $z = ru$ with $r > 0$ and $u \in \mathbb{T}$. Taking moduli, $|z| = |r||u| = r \cdot 1 = r$, again by Theorem 1.2.2 (d) and Proposition 1.2.1. Then $u = r^{-1}z = z/|z|$. So the pair is the one named in the statement.

(c) Since inverses are unique, $1/z = \overline{z}$ holds if and only if $z\overline{z} = 1$, that is, by Theorem 1.2.2 (c), if and only if $|z|^2 = 1 = 1^2$. Both $|z|$ and 1 are non-negative, so by Number Systems: From \(\nN\) to \(\nR\), Proposition 2.4.3 (g) neither can be strictly smaller than the other, and $|z|^2 = 1$ holds if and only if $|z| = 1$.

■

Part (b) is the picture we wanted: $z = |z| \cdot u$, a length times a direction, and both factors are forced. Part (a) says that the directions are closed under multiplication and inversion and contain 1; *Algebraic Preliminaries*, Chapter 3 calls a set with these properties, under an associative product, a group, and $\mathbb{T}$ is one of its first examples. Part (c) is a practical test: on the unit circle, and only there, dividing by z is the same as conjugating.

Quick check. Write $z = 3 + 4i$ as ru with $r > 0$ and $u \in \mathbb{T}$. What is $1/u$?

Solution. By Proposition 1.5.1 (b), $r = |3+4i| = 5$ and $u = \frac{3}{5} + \frac{4}{5}i$. By (a), $1/u = \overline{u} = \frac{3}{5} - \frac{4}{5}i$. Check: $(\frac{3}{5} + \frac{4}{5}i)(\frac{3}{5} - \frac{4}{5}i) = \frac{9}{25} + \frac{16}{25} = 1$.

So far a direction is a point u of $\mathbb{T}$, and multiplying two directions gives a third. What we cannot yet say is **how much** a direction turns the plane. For that we need to measure positions on the circle by a real number, an angle, so that multiplying directions adds angles. That is a job for cosine and sine.

1.5.2 What this volume borrows about cosine and sine

An angle should be a real number θ that walks around the circle: a point $(\cos\theta, \sin\theta)$ on $\mathbb{T}$, moving as θ moves, returning to its start after a full turn of length 2π. Defining such functions needs limits of functions, by power series or by integrals. This volume develops no limits of functions, so it does not define cosine and sine. *Analysis on the Real Line*, Chapter 10 does: there cos and sin are defined by power series, π is defined as twice the least positive zero of cos, and the facts below are proved. Its $\mathbb{R}$ is a complete ordered field, so by Number Systems: From \(\nN\) to \(\nR\), Theorem 6.3.1 it is our $\mathbb{R}$, up to the one order-preserving field isomorphism; the functions and the number carry over along it.

We write $2\pi\mathbb{Z} := \{2\pi k : k \in \mathbb{Z}\}$, the integer multiples of 2π in $\mathbb{R}$.

Axiom 1.5.2. What This Volume Borrows about Cosine and Sine

There are functions $\cos, \sin\colon \mathbb{R} \to \mathbb{R}$ and a real number $\pi > 0$ with the following properties. They are **assumed here and proved in *Analysis on the Real Line*, Chapter 10**.

(T1) $\cos^2\theta + \sin^2\theta = 1$ for every $\theta \in \mathbb{R}$.

(T2) For all $\theta, \varphi \in \mathbb{R}$,

$$\cos(\theta+\varphi) = \cos\theta\cos\varphi - \sin\theta\sin\varphi,$$
$$\sin(\theta+\varphi) = \sin\theta\cos\varphi + \cos\theta\sin\varphi.$$

(T3) For $\theta \in \mathbb{R}$: $\cos\theta = 1$ if and only if $\theta \in 2\pi\mathbb{Z}$.

(T4) For every $(x, y) \in \mathbb{R}^2$ with $x^2 + y^2 = 1$ there is $\theta \in \mathbb{R}$ with $\cos\theta = x$ and $\sin\theta = y$.

(T5) $\sin\theta > 0$ whenever $0 < \theta < \pi$.

In words: (T1) puts the point $(\cos\theta, \sin\theta)$ on the unit circle; (T2) is the pair of addition formulas; (T3) says that the cosine reaches 1 exactly at the multiples of a full turn; (T4) says that every point of the circle is reached; (T5) says that during the first half turn the point stays in the upper half plane.

This box is where the chapter borrows. Like Logic and Proof, Axiom 4.2.1, which *Logic and Proof* assumed and *Number Systems: From* $\mathbb{N}$ *to* $\mathbb{R}$, Chapter 1 proved, it is a statement assumed in one volume and proved in another. Nothing in Sections 1.1–1.4, and nothing in this section above the box, uses it. From here on we use these five facts about cos, sin and π and **no others**: no value such as $\cos(\pi/3) = \frac{1}{2}$, no periodicity, no picture of a graph. Whatever else we need, we derive, and each proof that uses one of the facts names it by its number. Two consequences come at once.

- $\cos 0 = 1$, by (T3), since $0 = 2\pi \cdot 0 \in 2\pi\mathbb{Z}$. Then (T1) gives $\sin^2 0 = 1 - 1 = 0$, so $\sin 0 = 0$, because a non-zero real number has a positive square (Number Systems: From \(\nN\) to \(\nR\), Proposition 2.4.3 (e)).
- $\cos(2\pi k) = 1$ and $\sin(2\pi k) = 0$ for every $k \in \mathbb{Z}$, by (T3) and then (T1) in the same way.

Warning. The five facts are about cos, sin and π only. A school fact such as $\sin(\pi/6) = \frac{1}{2}$ is not among them, and a proof in this volume may not quote it: Example (Special Angles) below derives the few values we use. Nor do the facts say that cos and sin are "periodic with period 2π" as functions; that is true, and derivable from (T2) and (T3), but no proof here needs it.

1.5.3 Polar form

With angles available, we can name the point of $\mathbb{T}$ at angle θ.

$e^{i\theta}$ is the point of the unit circle at angle θ.

Definition 1.5.1. Polar Form

For $\theta \in \mathbb{R}$ we write

$$e^{i\theta} := \cos\theta + i\sin\theta.$$

Let $z \in \mathbb{C}$ with $z \neq 0$. A **polar form** of z is an expression $z = re^{i\theta}$ with $r \in \mathbb{R}$, $r > 0$, and

$\theta \in \mathbb{R}$. Then θ is **an** argument of z.

In words: the first line is an abbreviation, not an exponential. This volume has no exponential function, the letter e on its own means nothing here, and the only rule of exponents we will use for $e^{i\theta}$ is the one proved below (Theorem 1.5.3 (a)). *Analysis on the Real Line*, Chapter 10 shows that $e^{i\theta}$ is indeed the value of the complex exponential function at $i\theta$; the notation anticipates that, and the linear algebra volumes write their roots of unity with it. The exponent may be written in any form equal to $i\theta$ with θ real, and the symbol then means $e^{i\theta}$: for example $e^{-i\theta} = e^{i(-\theta)}$, $e^{2\pi i} = e^{i\cdot 2\pi}$, $e^{in\theta} = e^{i(n\theta)}$ and $e^{-2\pi i} = e^{i(-2\pi)}$. This is unambiguous, because the real number θ is determined by $i\theta$ (it is its imaginary part). The second part asks for a **positive** real r and allows **any** real θ.

The point $e^{i\theta}$ lies on $\mathbb{T}$. Indeed $|e^{i\theta}| = \sqrt{\cos^2\theta + \sin^2\theta} = \sqrt{1} = 1$, by Definition 1.2.1 and (T1). So $re^{i\theta}$ is a length r times a direction, as in Proposition 1.5.1 (b), and that proposition gives $r = |z|$ for every polar form of z.

Examples.

- $e^{i0} = \cos 0 + i \sin 0 = 1$, by the consequences of the box. So $1 = 1 \cdot e^{i0}$ and $5 = 5e^{i0}$ are polar forms.
- $e^{2\pi i} = \cos 2\pi + i \sin 2\pi = 1$, by the second consequence with $k = 1$. So $1 = 1 \cdot e^{2\pi i}$ as well: 0 and 2π are both arguments of 1, and they differ, since $\pi > 0$.
- The number 0 has no polar form, since $|re^{i\theta}| = r > 0$.

A non-example by minimal change. Allow $r < 0$: then $-1 = (-1) \cdot e^{i0}$ looks like a polar form of -1 with argument 0. The clause that fails is $r > 0$, and it matters: with negative r allowed, the length could no longer be read off as $|z|$, and $1 = 1 \cdot e^{i0}$ and $-1 = (-1)e^{i0}$ would share the argument 0. We will see in Example (Special Angles) that $-1 = 1 \cdot e^{i\pi}$ is a genuine polar form.

The examples already show why the definition says **an** argument: $1 = e^{i0} = e^{2\pi i}$. How much freedom there is, exactly, is part (b) of Proposition 1.5.4 below. First, the reason for the notation: these numbers multiply the way exponentials do.

Theorem 1.5.3. Multiplication in Polar Form

(a) $e^{i\theta}e^{i\varphi} = e^{i(\theta+\varphi)}$ for all $\theta, \varphi \in \mathbb{R}$.

(b) $e^{i0} = 1$, and $(e^{i\theta})^{-1} = e^{-i\theta} = \overline{e^{i\theta}}$ for every $\theta \in \mathbb{R}$.

(c) If $r, s \in \mathbb{R}$ with $r, s > 0$ and $\theta, \varphi \in \mathbb{R}$, then $(re^{i\theta})(se^{i\varphi}) = rs\, e^{i(\theta+\varphi)}$, and this is a polar form of the product.

Idea. Multiply out $(\cos\theta + i\sin\theta)(\cos\varphi + i\sin\varphi)$ with the rule of Definition 1.1.1: the real and imaginary parts that appear are exactly the right-hand sides of the addition formulas (T2). The rest is bookkeeping with inverses and moduli.

Proof.

(a) By the multiplication rule of Definition 1.1.1,

$$\begin{aligned} e^{i\theta}e^{i\varphi} &= (\cos\theta\cos\varphi - \sin\theta\sin\varphi) + (\cos\theta\sin\varphi + \sin\theta\cos\varphi)i \\ &= \cos(\theta+\varphi) + i\sin(\theta+\varphi), \end{aligned}$$


where the second equality is (T2), with the two products in the imaginary part reordered by commutativity in $\mathbb{R}$. The right-hand side is $e^{i(\theta+\varphi)}$.

(b) $e^{i0} = 1$ was shown in the first example above. By (a) with $\varphi = -\theta$, $e^{i\theta}e^{-i\theta} = e^{i0} = 1$, so $e^{-i\theta}$ is the inverse of $e^{i\theta}$, by uniqueness of inverses in the field $\mathbb{C}$ (Theorem 1.2.4, Number Systems: From \(\nN\) to \(\nR\), Lemma 4.1.1 (a)). Since $e^{i\theta} \in \mathbb{T}$, its inverse is also its conjugate (Proposition 1.5.1 (a)).

(c) By Proposition 1.1.2 (commutativity and associativity of $\cdot$), $(re^{i\theta})(se^{i\varphi}) = (rs)(e^{i\theta}e^{i\varphi})$, and (a) turns this into $rs\,e^{i(\theta+\varphi)}$. Here $rs > 0$ by Number Systems: From \(\nN\) to \(\nR\), Proposition 2.4.3 (c) (multiply $0 < r$ by $s > 0$), so this is a polar form.

■

In words: **to multiply two complex numbers, multiply their moduli and add their arguments**. This is the chapter's third named move, **multiply moduli, add angles**. Multiplying by a fixed $w = se^{i\varphi}$ therefore turns the plane about 0 by the angle φ and stretches it by the factor s. The picture shows z, w and zw for $|z| = 1.5$, $|w| = 1.2$, and arguments of 20° and 50° (a full turn being 360°, that is, 2π).



The arrow for zw has length $1.5 \times 1.2 = 1.8$ and makes the angle $20° + 50° = 70°$ with the positive real axis: it is the arrow for z, turned by the argument of w and stretched by the modulus of w. The small arc marks the argument of z, the large arc the extra turn by the argument of w, and the dashed arc is part of the unit circle, where the numbers $e^{i\theta}$ live.

Now the freedom in the argument. Every non-zero number has a polar form, and its arguments are determined up to a whole number of full turns.

Proposition 1.5.4. Existence and Uniqueness of Polar Form

(a) Every $z \in \mathbb{C}$ with $z \neq 0$ has a polar form, and every polar form $z = re^{i\theta}$ has $r = |z|$.

(b) For $r, s \in \mathbb{R}$ with $r, s > 0$ and $\theta, \varphi \in \mathbb{R}$,

$$re^{i\theta} = se^{i\varphi} \iff r = s \text{ and } \theta - \varphi \in 2\pi\mathbb{Z}.$$

Proof.

(a) By Proposition 1.5.1 (b), $z = |z| \cdot u$ with $u = x + yi \in \mathbb{T}$, so $x^2 + y^2 = 1$. By (T4) there is $\theta \in \mathbb{R}$ with $\cos\theta = x$ and $\sin\theta = y$, that is, $u = e^{i\theta}$. Hence $z = |z|e^{i\theta}$, with $|z| > 0$. Conversely, if $z = re^{i\theta}$ is a polar form, then $e^{i\theta} \in \mathbb{T}$, and the uniqueness in Proposition 1.5.1 (b) gives $r = |z|$.

(b) Suppose first that $re^{i\theta} = se^{i\varphi}$. The number $re^{i\theta}$ is non-zero: $r \neq 0$, $e^{i\theta} \in \mathbb{T}$ is non-zero (Proposition 1.5.1 (a)), and a product of non-zero elements of the field $\mathbb{C}$ is non-zero (Theorem 1.2.4, Number Systems: From \(\nN\) to \(\nR\), Lemma 4.1.1 (d)). So both sides are polar forms of the same non-zero number, and $r = s$ by (a). Multiplying by r^{-1} gives $e^{i\theta} = e^{i\varphi}$. In the other direction, if $r = s$, the two sides are equal exactly when $e^{i\theta} = e^{i\varphi}$. So it remains to show that $e^{i\theta} = e^{i\varphi}$ if and only if $\theta - \varphi \in 2\pi\mathbb{Z}$.

Multiplying by $e^{-i\varphi}$, and back by $e^{i\varphi}$, shows that $e^{i\theta} = e^{i\varphi}$ holds if and only if $e^{i\theta}e^{-i\varphi} = e^{i\varphi}e^{-i\varphi}$, that is, by Theorem 1.5.3 (a) and (b), if and only if $e^{i(\theta-\varphi)} = 1$. Comparing real and imaginary parts, this says $\cos(\theta - \varphi) = 1$ and $\sin(\theta - \varphi) = 0$. The second condition follows from the first: by (T1), $\cos(\theta - \varphi) = 1$ gives $\sin^2(\theta - \varphi) = 0$, hence $\sin(\theta - \varphi) = 0$ (Number Systems: From \(\nN\) to \(\nR\), Proposition 2.4.3 (e)). So $e^{i\theta} = e^{i\varphi}$ if and only if $\cos(\theta - \varphi) = 1$, and by (T3) that holds if and only if $\theta - \varphi \in 2\pi\mathbb{Z}$.

■

Existence came from (T4), and uniqueness from (T3), with (T1) helping and (T2) entering through Theorem 1.5.3. "Comparing arguments" therefore always means comparing them up to a multiple of 2π, and the next warning is the trap this sets.

Warning. From $e^{i\theta} = e^{i\varphi}$ one may **not** conclude $\theta = \varphi$: $e^{i0} = e^{2\pi i} = 1$, and $0 \neq 2\pi$. Only $\theta - \varphi \in 2\pi\mathbb{Z}$ follows (Proposition 1.5.4 (b)). Forgetting this loses solutions as soon as an equation is solved for an angle; Section 1.6 meets an equation where it would lose all but one of them.

1.5.4 Powers

Repeating Theorem 1.5.3 gives powers. The powers z^n are those of the recursion $z^0 = 1$, $z^{n+1} = z^n z$ (Proposition 1.1.3).

Theorem 1.5.5. De Moivre's Formula

For every $\theta \in \mathbb{R}$ and every $n \in \mathbb{N}$,

$$(e^{i\theta})^n = e^{in\theta}, \quad \text{that is,} \quad (\cos\theta + i\sin\theta)^n = \cos n\theta + i\sin n\theta.$$

Consequently $(re^{i\theta})^n = r^n e^{in\theta}$ for every real $r > 0$, and this is a polar form.

Idea. A power is a repeated product, and each new factor $e^{i\theta}$ adds θ to the argument, by Theorem 1.5.3 (a). So induction on n, along the recursion $z^{n+1} = z^n z$.

Proof. We use Logic and Proof, Axiom 4.2.1 on n, read on $\mathbb{N} = \omega$ through Number Systems: From \(\nN\) to \(\nR\), Corollary 1.4.3 (b); here $n\theta$ is the product in $\mathbb{R}$ of $n \cdot 1_{\mathbb{R}}$ and θ (Number Systems: From \(\nN\) to \(\nR\), Definition 4.2.1 (a)). For $n = 0$, the left side is $(e^{i\theta})^0 = 1$ by the recursion, and the right side is $e^{i0} = 1$ by Theorem 1.5.3 (b). Suppose $(e^{i\theta})^n = e^{in\theta}$ for some $n \in \mathbb{N}$. Then

$$(e^{i\theta})^{n+1} = (e^{i\theta})^n e^{i\theta} = e^{in\theta} e^{i\theta} = e^{i(n\theta+\theta)} = e^{i(n+1)\theta},$$

by the recursion, the induction hypothesis, Theorem 1.5.3 (a), and $(n+1) \cdot 1_{\mathbb{R}} = n \cdot 1_{\mathbb{R}} + 1_{\mathbb{R}}$ (Number Systems: From \(\nN\) to \(\nR\), Definition 4.2.1 (a)) with distributivity in $\mathbb{R}$. This proves the formula for every $n \in \mathbb{N}$.

For the consequence, Proposition 1.1.3 (c) gives $(re^{i\theta})^n = r^n (e^{i\theta})^n = r^n e^{in\theta}$. The power r^n is the same whether it is computed in $\mathbb{R}$ or in $\mathbb{C}$, by the remark after Proposition 1.1.3 in Section 1.1, and $r^n > 0$ because a product of positive reals is positive (Number Systems: From \(\nN\) to \(\nR\), Proposition 2.4.3 (c); induction on n, with $r^0 = 1 > 0$).

■

For a negative exponent, Exercise C1 below defines z^{-n} locally and proves the matching formula; nothing in the text uses it.

The five borrowed facts give no particular value of cos or sin except at multiples of 2π. Every other value this chapter uses is derived next, from (T1)–(T5) and the algebra of Sections 1.1–1.4.

Example. Special Angles

Show, using only (T1)–(T5) and the results proved so far, that

$$e^{i\pi} = -1, \qquad e^{i\pi/2} = i, \qquad e^{i\pi/3} = \tfrac{1}{2} + \tfrac{\sqrt{3}}{2} i,$$
$$e^{i\pi/6} = \tfrac{\sqrt{3}}{2} + \tfrac{1}{2} i, \qquad e^{i\pi/4} = \tfrac{1}{\sqrt{2}} + \tfrac{1}{\sqrt{2}} i.$$

Solution. Each value is found the same way: a power of the unknown point is a point we already know, and (T5) chooses between the candidates. Since $\pi > 0$, each of $\pi/2$, $\pi/3$, $\pi/4$, $\pi/6$ lies strictly between 0 and π, so (T5) gives $\operatorname{Im} e^{i\theta} = \sin\theta > 0$ for each of them.

(i) $e^{i\pi} = -1$. Let $u := e^{i\pi}$. By Theorem 1.5.5, $u^2 = e^{2\pi i} = 1$, so $(u-1)(u+1) = u^2 - 1 = 0$, and since $\mathbb{C}$ has no zero divisors (Theorem 1.2.4, Number Systems: From \(\nN\) to \(\nR\), Lemma 4.1.1 (d)), $u = 1$ or $u = -1$. Suppose $u = 1$. Then $\cos\pi = 1$, so $\pi = 2\pi k$ for some $k \in \mathbb{Z}$ by (T3); dividing by $\pi \neq 0$ gives $1 = 2k$ in $\mathbb{R}$, an equation between images of integers. The embedding of $\mathbb{Z}$ into $\mathbb{R}$, through $\mathbb{Q}$ (Number Systems: From \(\nN\) to \(\nR\), Proposition 3.1.4) and κ (Number Systems: From \(\nN\) to \(\nR\), Theorem 5.5.2), is injective and preserves $+$ and $\cdot$, so $1 = 2k$ holds in $\mathbb{Z}$. But $1 = 2 \cdot 0 + 1$ is odd, and an odd integer is not of the form $2k$ (Number Systems: From \(\nN\) to \(\nR\), Corollary 2.5.3 (a)). So $u \neq 1$, and $e^{i\pi} = -1$.

(ii) $e^{i\pi/2} = i$. Let $u := e^{i\pi/2}$. By Theorem 1.5.3 (a), $u^2 = e^{i\pi} = -1 = i^2$, so $(u-i)(u+i) = 0$ and $u = \pm i$. Since $\operatorname{Im} u > 0$ and $\operatorname{Im}(-i) = -1 < 0$, $u = i$.

(iii) $e^{i\pi/3} = \frac{1}{2} + \frac{\sqrt{3}}{2} i$. Let $u := e^{i\pi/3}$. By Theorem 1.5.5, $u^3 = e^{i\pi} = -1$. Multiplying out shows $(u+1)(u^2-u+1) = u^3+1 = 0$. Now $u \neq -1$, since $\operatorname{Im} u > 0$; so $u^2-u+1 = 0$. By

Corollary 1.4.2 with $a = 1$, $b = -1$, $c = 1$, and $d := \sqrt{3}\,i$, a square root of $b^2 - 4ac = -3$ (indeed $(\sqrt{3}\,i)^2 = 3i^2 = -3$),

$$u = \frac{1 + \sqrt{3}\,i}{2} \quad \text{or} \quad u = \frac{1 - \sqrt{3}\,i}{2}.$$

Since $\operatorname{Im} u > 0$, $u = \frac{1}{2} + \frac{\sqrt{3}}{2}i$.

(iv) $e^{i\pi/6} = \frac{\sqrt{3}}{2} + \frac{1}{2}i$. Since $\pi/6 = \pi/2 + (-\pi/3)$, Theorem 1.5.3 (a) and (b) give

$$e^{i\pi/6} = e^{i\pi/2}e^{-i\pi/3} = i \cdot \overline{e^{i\pi/3}} = i\Big(\frac{1}{2} - \frac{\sqrt{3}}{2}i\Big) = \frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

by (ii) and (iii).

(v) $e^{i\pi/4} = \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i$. Let $u := e^{i\pi/4}$. By Theorem 1.5.3 (a) and (ii), $u^2 = e^{i\pi/2} = i$. The formula of Proposition 1.4.1 for $z = i$ (so $a = 0$, $b = 1$, $r = |i| = 1$) gives the square root $w = \sqrt{1/2} + \sqrt{1/2}\,i$, and $\sqrt{1/2} = 1/\sqrt{2}$, since both are non-negative and square to $\frac{1}{2}$. By Proposition 1.4.1 (b), $u = w$ or $u = -w$, and $\operatorname{Im} u > 0$ rules out $-w$. So $u = w$.

Comparing real and imaginary parts turns these into values of cos and sin:

θ	0	$\pi/6$	$\pi/4$	$\pi/3$	$\pi/2$	π
$\cos\theta$	1	$\sqrt{3}/2$	$1/\sqrt{2}$	$1/2$	0	-1
$\sin\theta$	0	$1/2$	$1/\sqrt{2}$	$\sqrt{3}/2$	1	0

These are the familiar values, now proved from the five facts.

Two special cases of the rotation picture are worth isolating. Multiplying by $i = e^{i\pi/2}$ turns the plane by a quarter turn, counterclockwise; two quarter turns make a half turn, and this is the picture behind $i^2 = -1 = e^{i\pi}$. And multiplying by $e^{-i\theta} = \overline{e^{i\theta}}$ undoes the turn by θ (Theorem 1.5.3 (b)).

Powers that are tedious to expand become one line in polar form.

Example. A Power via Polar Form

Compute $(\sqrt{3} + i)^6$.

Solution. We have $\sqrt{3} + i = 2(\frac{\sqrt{3}}{2} + \frac{1}{2}i) = 2e^{i\pi/6}$, by Example (Special Angles) (iv). By Theorem 1.5.5 and Example (Special Angles) (i),

$$(\sqrt{3} + i)^6 = 2^6 e^{i\cdot 6\pi/6} = 64\,e^{i\pi} = -64.$$

By hand: $(\sqrt{3}+i)^2 = 3+2\sqrt{3}\,i-1 = 2+2\sqrt{3}\,i = 2(1+\sqrt{3}\,i)$, and $(1+\sqrt{3}\,i)^3 = (1+\sqrt{3}\,i)(-2+2\sqrt{3}\,i) = -2-6 = -8$, so $(\sqrt{3}+i)^6 = \big(2(1+\sqrt{3}\,i)\big)^3 = 8\cdot(-8) = -64$ (Proposition 1.1.3 (b), (c)), the same answer.

Quick check. Use polar form to compute $(1+i)^8$, and check the answer with $(1+i)^2 = 2i$.

Solution. By Example (Special Angles) (v), $1+i = \sqrt{2}(\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i) = \sqrt{2}\,e^{i\pi/4}$. By Theorem 1.5.5, $(1+i)^8 = (\sqrt{2})^8 e^{i\cdot 8\pi/4} = 16\,e^{2\pi i} = 16$, since $(\sqrt{2})^8 = ((\sqrt{2})^2)^4 = 2^4 = 16$ and $e^{2\pi i} = 1$. Check: $(1+i)^8 = ((1+i)^2)^4 = (2i)^4 = 16i^4 = 16$, by Proposition 1.1.3 (b), (c).

1.5.5 Exercises

1.5.5.1 A. Check your understanding

Exercise. A1

(a) Give a polar form of -2 and of $3i$.

(b) True or false: if $e^{i\theta} = e^{i\varphi}$, then $\theta = \varphi$. Justify your answer.

(c) In Proposition 1.5.4, which of the borrowed facts (T1)–(T5) gives the existence of a polar form, and which gives the uniqueness of the argument up to $2\pi\mathbb{Z}$?

(d) Which results of Sections 1.1–1.5 use none of the borrowed facts? Is Proposition 1.5.1 among them?

Solution.

(a) $-2 = 2e^{i\pi}$ and $3i = 3e^{i\pi/2}$, by Example (Special Angles) (i) and (ii); the moduli 2 and 3 are positive, as a polar form requires. (Any $\pi + 2\pi k$, respectively $\pi/2 + 2\pi k$, with $k \in \mathbb{Z}$, is also an argument, by Proposition 1.5.4 (b).)

(b) False. $e^{i0} = 1 = e^{2\pi i}$, but $0 \neq 2\pi$ since $\pi > 0$. What is true is $\theta - \varphi \in 2\pi\mathbb{Z}$ (Proposition 1.5.4 (b)).

(c) Existence: (T4), which provides an angle for the point $z/|z|$ of $\mathbb{T}$. Uniqueness: (T3), which turns $\cos(\theta-\varphi) = 1$ into $\theta - \varphi \in 2\pi\mathbb{Z}$; (T1) is used alongside it, to see that $\sin(\theta-\varphi) = 0$ follows from $\cos(\theta-\varphi) = 1$, and (T2) enters through Theorem 1.5.3, which turns $e^{i\theta} = e^{i\varphi}$ into $e^{i(\theta-\varphi)} = 1$.

(d) Everything in Sections 1.1–1.4, and the part of this section before Axiom 1.5.2; after the box, only Exercise C2, which is set up to avoid them. In particular Proposition 1.5.1 uses none of them: its proof uses only the modulus rules of Section 1.2, the field properties of $\mathbb{C}$, and order facts about $\mathbb{R}$ from *Number Systems: From $\mathbb{N}$ to $\mathbb{R}$*.

1.5.5.2 B. Practice

Exercise. B1

Compute $(1 - \sqrt{3}\,i)^6$.

Solution. We have $1 - \sqrt{3}\,i = 2(\frac{1}{2} - \frac{\sqrt{3}}{2}i) = 2\,\overline{e^{i\pi/3}} = 2e^{-i\pi/3}$, by Example (Special Angles)

(iii) and Theorem 1.5.3 (b). By Theorem 1.5.5,

$$(1-\sqrt{3}\,i)^6 = 2^6 e^{-6i\pi/3} = 64\,e^{-2\pi i} = 64,$$

since $-2\pi = 2\pi\cdot(-1) \in 2\pi\mathbb{Z}$, so $\cos(-2\pi) = 1$ and $\sin(-2\pi) = 0$ by (T3) and (T1), and $e^{-2\pi i} = 1$.

Exercise. B2: Symmetries of Cosine and Sine

Prove that for every $\theta \in \mathbb{R}$:

$$\cos(-\theta) = \cos\theta, \quad \sin(-\theta) = -\sin\theta, \quad \cos(\theta+\pi) = -\cos\theta,$$
$$\sin(\theta+\pi) = -\sin\theta, \quad \cos\theta = \sin(\theta+\pi/2).$$

Hint: write each left side as a part of some $e^{i\psi}$, and compute that $e^{i\psi}$ with Theorem 1.5.3.

Solution. By Theorem 1.5.3 (b), $e^{-i\theta} = \overline{e^{i\theta}}$, that is, $\cos(-\theta) + i\sin(-\theta) = \cos\theta - i\sin\theta$. Comparing real and imaginary parts gives $\cos(-\theta) = \cos\theta$ and $\sin(-\theta) = -\sin\theta$.

By Theorem 1.5.3 (a) and Example (Special Angles) (i), $e^{i(\theta+\pi)} = e^{i\theta}e^{i\pi} = -e^{i\theta}$, that is, $\cos(\theta+\pi) + i\sin(\theta+\pi) = -\cos\theta - i\sin\theta$. Comparing parts gives the next two identities.

By Theorem 1.5.3 (a) and Example (Special Angles) (ii), $e^{i(\theta+\pi/2)} = ie^{i\theta} = i\cos\theta - \sin\theta$. Comparing imaginary parts gives $\sin(\theta+\pi/2) = \cos\theta$. (Comparing real parts gives, in addition, $\cos(\theta+\pi/2) = -\sin\theta$.)

Exercise. B3

Let $\theta \in \mathbb{R}$. Show that $|e^{i\theta} - 1|^2 = 2 - 2\cos\theta$. Hence show again that $e^{i\theta} = 1$ if and only if $\cos\theta = 1$.

Solution. Let $u := e^{i\theta}$, so $u\overline{u} = |u|^2 = 1$. Squaring the modulus, by Theorem 1.2.2 (a)–(c),

$$|u-1|^2 = (u-1)\overline{(u-1)} = (u-1)(\overline{u}-1) = u\overline{u} - (u+\overline{u}) + 1 = 2 - 2\operatorname{Re} u,$$

and $\operatorname{Re} u = \cos\theta$. Hence $|u-1|^2 = 2 - 2\cos\theta$.

Now $u = 1$ if and only if $|u-1| = 0$ (Theorem 1.2.2 (e)), if and only if $|u-1|^2 = 0$ (a real number is 0 exactly when its square is: $0^2 = 0$ by Number Systems: From \(\nN\) to \(\nR\), Proposition 2.3.2 (b), and a non-zero square is positive by Number Systems: From \(\nN\) to \(\nR\), Proposition 2.4.3 (e)), if and only if $2 - 2\cos\theta = 0$, that is, $\cos\theta = 1$.

1.5.5.3 C. Going deeper

Exercise. C1: Triple Angles and Negative Powers

(a) Prove that $\cos 3\theta = 4\cos^3\theta - 3\cos\theta$ and $\sin 3\theta = 3\sin\theta - 4\sin^3\theta$ for every $\theta \in \mathbb{R}$.

(b) For $z \in \mathbb{C}$ with $z \neq 0$ and $n \in \mathbb{N}$, put $z^{-n} := (z^{-1})^n$ (a definition local to this exercise). Show that $(e^{i\theta})^{-n} = e^{-in\theta}$.

Solution.

(a) Write $c := \cos\theta$ and $s := \sin\theta$. By Theorem 1.5.5, $\cos 3\theta + i \sin 3\theta = (c + is)^3$. Now $(c + is)^2 = (c^2 - s^2) + 2cs\,i$, and

$$\begin{aligned}(c + is)^3 &= ((c^2 - s^2) + 2cs\,i)(c + is) \\ &= (c^3 - 3cs^2) + (3c^2 s - s^3)i.\end{aligned}$$

Comparing real parts and using $s^2 = 1 - c^2$ (T1), $\cos 3\theta = c^3 - 3c(1 - c^2) = 4c^3 - 3c$. Comparing imaginary parts and using $c^2 = 1 - s^2$, $\sin 3\theta = 3(1 - s^2)s - s^3 = 3s - 4s^3$.

(b) By Theorem 1.5.3 (b), $(e^{i\theta})^{-1} = e^{-i\theta} = e^{i(-\theta)}$. By Theorem 1.5.5 applied to $-\theta$, $(e^{i\theta})^{-n} = (e^{i(-\theta)})^n = e^{in(-\theta)} = e^{-in\theta}$.

Nothing in this chapter uses negative powers of complex numbers; the definition in (b) is for practice only.

Exercise. C2: Points on the Circle without Angles

For $t \in \mathbb{R}$ put

$$u_t := \frac{1 + it}{1 - it}.$$

(a) Show that u_t is defined, that $u_t \in \mathbb{T}$, and that $u_t \neq -1$.

(b) Show that every $u \in \mathbb{T}$ with $u \neq -1$ equals u_t for **exactly one** $t \in \mathbb{R}$, namely $t = \operatorname{Im} u/(1 + \operatorname{Re} u)$.

(c) Compute $u_{1/2}$ in the form $a + bi$.

Use none of the borrowed facts (T1)–(T5).

Solution (Hint 1). For (b), on the unit circle the conjugate is the inverse: $\overline{u} = 1/u$.

Solution (Hint 2). Clearing the denominator, $u = u_t$ is equivalent to $u - 1 = it(u + 1)$, and $u + 1 \neq 0$, so t is determined; it remains to show that $(u - 1)/(i(u + 1))$ is real.

Solution.

(a) The real part of $1 - it$ is 1, so $1 - it \neq 0$ and the quotient is defined. The numbers $1 + it$ and $1 - it$ are conjugate, so they have the same modulus (Theorem 1.2.2 (d)), which is non-zero. From $u_t(1 - it) = 1 + it$ and Theorem 1.2.2 (d), $|u_t||1 - it| = |1 + it| = |1 - it|$, and canceling the non-zero real number $|1 - it|$ gives $|u_t| = 1$. If $u_t = -1$, then $1 + it = -(1 - it) = -1 + it$, so $1 = -1$, which is false. So $u_t \neq -1$.

(b) Let $u \in \mathbb{T}$ with $u \neq -1$, and let $t \in \mathbb{R}$. Since $1 - it \neq 0$, the equation $u = u_t$ is

equivalent to $u(1-it) = 1+it$, that is, to

$$u - 1 = it(u+1).$$

Since $i(u+1) \neq 0$, this holds if and only if $t = t_0$, where $t_0 \coloneqq (u-1)/(i(u+1)) \in \mathbb{C}$. So at most one real t has $u_t = u$, and one does exactly when t_0 is real. Using $u\overline{u} = 1$,

$$(u-1)(\overline{u}+1) = u\overline{u} + u - \overline{u} - 1 = u - \overline{u} = 2i \operatorname{Im} u,$$

where $u - \overline{u} = 2i \operatorname{Im} u$ by Definition 1.2.1 (with $u = a+bi$, $(a+bi)-(a-bi) = 2bi$); and $(u+1)(\overline{u}+1) = |u+1|^2 = 2 + 2\operatorname{Re} u$, by Theorem 1.2.2 (a), (c). This is a positive real number, since $u + 1 \neq 0$. Multiplying numerator and denominator of $(u-1)/(u+1)$ by $\overline{u}+1 = \overline{u+1} \neq 0$,

$$t_0 = \frac{1}{i} \cdot \frac{2i \operatorname{Im} u}{2 + 2\operatorname{Re} u} = \frac{\operatorname{Im} u}{1 + \operatorname{Re} u},$$

a real number. So exactly one $t \in \mathbb{R}$ has $u_t = u$, namely t_0.

(c) Multiplying numerator and denominator by the conjugate $1 + \frac{1}{2}i$ of the denominator,

$$u_{1/2} = \frac{(1+\frac{1}{2}i)^2}{1+\frac{1}{4}} = \frac{\frac{3}{4}+i}{\frac{5}{4}} = \frac{3}{5} + \frac{4}{5}i.$$

Check against (b): $\operatorname{Im} u/(1+\operatorname{Re} u) = \frac{4}{5}/\frac{8}{5} = \frac{1}{2}$.

So the map $t \mapsto u_t$ is a bijection from $\mathbb{R}$ onto $\mathbb{T} \setminus \{-1\}$, and it was found with no angle. For rational t, the point u_t has rational coordinates: the computation in (c) with t in place of $\frac{1}{2}$ gives $u_t = ((1-t^2) + 2t\,i)/(1+t^2)$, and $t = 1/2$ gives the point $(3/5, 4/5)$ behind the right triangle with sides 3, 4, 5.

Marking scheme. (10 points) - 2: $1 - it \neq 0$ with a reason, and $|u_t| = 1$ from the equal moduli of conjugates. - 1: $u_t \neq -1$. - 2: $u = u_t$ rewritten as a linear equation in t, with $u + 1 \neq 0$ used to solve it uniquely. - 3: shows that the unique solution is real and equals $\operatorname{Im} u/(1+\operatorname{Re} u)$, with $1 + \operatorname{Re} u \neq 0$ justified (for example by multiplying by $\overline{u+1}$ and using $u\overline{u} = 1$, with $2 + 2\operatorname{Re} u > 0$). - Second route (coordinates $u = x + yi$, $x^2 + y^2 = 1$): the two lines above (5 points) become 2 for splitting $u - 1 = it(u+1)$ into $x - 1 = -ty$ and $y = t(1+x)$, showing $x \neq -1$ (otherwise $u = -1$) and so $t = y/(1+x)$ is forced; and 3 for checking that this t satisfies $x - 1 = -ty$, using $x^2 + y^2 = 1$. - 2: (c) computed and checked. Fatal errors (0 for the step): writing $u = e^{i\theta}$ and using a half-angle formula (borrowed facts, and values not derived here); dividing by $1 + \operatorname{Re} u$ without showing it is non-zero.