4.1 Working in an Ordered Field

Section 2.4 proved the rules for inequalities from the order axioms, and Section 3.2 added the one rule that needs division: reciprocals reverse order. A limit argument asks for a few more. It halves a tolerance, so it needs $0 < \varepsilon/2 < \varepsilon$; it subtracts fractions, so it needs $1/b - 1/d = (d-b)/(bd)$; and above all it measures how far apart two numbers are, so it needs the absolute value and the triangle inequality. Each of these is a few lines from the rules we have. This section proves them once, in an arbitrary ordered field, and the rest of the chapter uses them on nearly every page.

4.1.1 The standing setup

Throughout this chapter, F is an **ordered field** (Definition 2.4.1): a field with a relation $<$ satisfying (O1)–(O4). The rationals are one, by Theorem 3.2.1, and for most of the chapter they are the one that matters: the next chapter applies these results first with $F = \mathbb{Q}$, to build the real numbers from Cauchy sequences of rationals, and then again with $F = \mathbb{R}$, once $\mathbb{R}$ is shown to be an ordered field. We state results for a general F because the proofs never use anything about $\mathbb{Q}$ beyond (F1)–(F9) and (O1)–(O4), and saying so on the page makes it checkable.

From here on we use the field rules and the rules for inequalities of Sections 2.3, 2.4 and 3.2 without naming them, as in school algebra. A step is cited only where it needs care: multiplying or dividing an inequality, which needs the sign of the factor; taking reciprocals, which reverses order (Proposition 3.2.2); and a weak inequality, which hides an equality case. Such a step cites its rule once in each proof. In a field $1 \neq 0$, so $1 > 0$ (Proposition 2.4.3 (e)). We write

$$F_{>0} := \{x \in F : x > 0\}, \qquad F_{\geq 0} := \{x \in F : x \geq 0\},$$

so that $\mathbb{Q}_{>0}$ of Section 3.2 is the case $F = \mathbb{Q}$. Recall that $a/b := ab^{-1}$ for $b \neq 0$, so $1/b = b^{-1}$, and that $2 := 1 + 1$ and $x^2 := xx$ (Proposition 2.3.2 (d)).

4.1.2 Fractions

School algebra adds fractions by cross-multiplying and subtracts reciprocals by a common denominator. Both are identities in any field, and the second is how we will compare $1/b_n$ with $1/b$ in Section 4.4. The only point needing care is that a product of non-zero elements is non-zero, so that bd may stand in a denominator.

Lemma 4.1.1. Fraction Rules in a Field

Let F be a field (Definition 2.3.1), and let $a, b, c, d \in F$ with $b \neq 0$ and $d \neq 0$.

(a) b^{-1} is the **only** element y with $by = 1$, and $(b^{-1})^{-1} = b$. Moreover $bd \neq 0$ and $(bd)^{-1} = b^{-1}d^{-1}$.

(b) $\dfrac{a}{b} + \dfrac{c}{d} = \dfrac{ad + bc}{bd}$ and $\dfrac{a}{b} \cdot \dfrac{c}{d} = \dfrac{ac}{bd}$.

(c) $\dfrac{1}{b} - \dfrac{1}{d} = \dfrac{d - b}{bd}$.

(d) For **all** $x, y \in F$: if $xy = 0$, then $x = 0$ or $y = 0$.

Idea. Everything rests on one move from Section 2.3: to identify an element, show it satisfies the **defining equation** of the element you want and let uniqueness finish. The product $b^{-1}d^{-1}$ satisfies the defining equation of $(bd)^{-1}$; once bd is known to be a legal denominator, (b) and (c) are bookkeeping.

Proof.

(a) Suppose $by = 1$. Then

$$y = 1 \cdot y = (b^{-1}b)y = b^{-1}(by) = b^{-1} \cdot 1 = b^{-1}.$$

This is the uniqueness argument of Proposition 2.3.2 (a), run with multiplication. Next, $b^{-1} \neq 0$: otherwise $1 = bb^{-1} = b \cdot 0 = 0$ by Proposition 2.3.2 (b), against $1 \neq 0$. So $(b^{-1})^{-1}$ exists, and since $b^{-1}b = bb^{-1} = 1$, the element b satisfies its defining equation; uniqueness gives $(b^{-1})^{-1} = b$. Now

$$(bd)(b^{-1}d^{-1}) = (bb^{-1})(dd^{-1}) = 1 \cdot 1 = 1.$$

If $bd = 0$, the left side would be $0 \cdot (b^{-1}d^{-1}) = 0$ by Proposition 2.3.2 (b), so $1 = 0$, which is false in a field. Hence $bd \neq 0$, and the display says that $b^{-1}d^{-1}$ satisfies the defining equation of $(bd)^{-1}$; by uniqueness, $(bd)^{-1} = b^{-1}d^{-1}$.

(b) By (a), $bd \neq 0$, so both right-hand sides are defined. For the product, by (a),

$$\begin{aligned} \frac{a}{b} \cdot \frac{c}{d} &= (ab^{-1})(cd^{-1}) = (ac)(b^{-1}d^{-1}) \\ &= (ac)(bd)^{-1} = \frac{ac}{bd}. \end{aligned}$$

For the sum, first $\dfrac{ad}{bd} = (ad)(b^{-1}d^{-1}) = (ab^{-1})(dd^{-1}) = \dfrac{a}{b}$, and likewise $\dfrac{bc}{bd} = \dfrac{c}{d}$. Then distributivity gives

$$\begin{aligned} \frac{a}{b} + \frac{c}{d} &= (ad)(bd)^{-1} + (bc)(bd)^{-1} \\ &= (ad + bc)(bd)^{-1} = \frac{ad + bc}{bd}. \end{aligned}$$

(c) Apply the sum rule of (b) with $a = 1$ and $c = -1$. Since $(-1)d^{-1} = -(d^{-1})$ by Proposition 2.3.2 (c), the left side is $1/b + (-1)/d = 1/b - 1/d$, and the numerator on the right is $1 \cdot d + b(-1) = d - b$.

(d) Let $xy = 0$, and suppose $x \neq 0$. Then $y = 1 \cdot y = (x^{-1}x)y = x^{-1}(xy) = x^{-1} \cdot 0 = 0$. So $x = 0$ or $y = 0$. This proves the lemma.

■

In an ordered field, (d) is also Proposition 2.4.3 (f). The point of (d) here is that it needs **no** order: a field has no zero divisors. The commutative ring of integers modulo 6, in the warning after Definition 2.3.1, has them ($[2][3] = [0]$ with $[2] \neq [0] \neq [3]$), so by (d) it is not a field.

4.1.3 Halving and comparing

The commonest move in a limit proof is to give each of two terms half the tolerance. That needs $\varepsilon/2$ to be positive, smaller than ε, and to add up to ε with itself. The next proposition collects

these facts with two comparisons we will need as often.

Proposition 4.1.2. Halves and Comparisons in an Ordered Field

Let F be an ordered field.

(a) $\leq$ is a total order on F (Sets and Functions, Definition 5.3.1), and its strict part is $<$.

(b) $2 := 1 + 1 > 0$. For every $\varepsilon \in F_{>0}$,

$$0 < \frac{\varepsilon}{2} < \varepsilon \qquad \text{and} \qquad \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.$$

For all $a, b \in F$ with $a < b$, $a < \dfrac{a+b}{2} < b$.

(c) For $a, b \in F_{>0}$: if $a \leq b$, then $\dfrac{1}{b} \leq \dfrac{1}{a}$.

(d) For $a, b, c, d \in F$: if $0 \leq a \leq b$ and $0 \leq c \leq d$, then $ac \leq bd$.

Idea. Each part is one or two of the rules of Sections 2.4 and 3.2 in sequence. The only care needed is with $\leq$, which hides an "or": each non-strict hypothesis splits into an equality case and a strict case, and the strict rules apply only to the second.

Proof.

(a) This is Proposition 2.4.3 (i).

(b) Since $1 > 0$, (O3) gives $2 = 1 + 1 > 0 + 1 = 1$, and $2 > 0$ by (O2). So $2 \neq 0$, and $2^{-1} > 0$ by Proposition 3.2.2 (a). For $\varepsilon > 0$, $\varepsilon/2 = \varepsilon \cdot 2^{-1} > 0$ by (O4), and

$$\begin{aligned} \frac{\varepsilon}{2} + \frac{\varepsilon}{2} &= \varepsilon \cdot 2^{-1} \cdot 1 + \varepsilon \cdot 2^{-1} \cdot 1 \\ &= (\varepsilon \cdot 2^{-1})(1 + 1) = \varepsilon(2^{-1} \cdot 2) = \varepsilon. \end{aligned}$$

Adding $\varepsilon/2$ to $0 < \varepsilon/2$ by (O3) gives $\varepsilon/2 < \varepsilon/2 + \varepsilon/2 = \varepsilon$. For the midpoint, let $a < b$. By (O3), $a + a < a + b$ and $a + b < b + b$, and $a + a = a(1 + 1) = 2a$, $b + b = 2b$. Multiplying by $2^{-1} > 0$ (Proposition 2.4.3 (c)) gives $a < (a + b)/2 < b$, since $(2a)2^{-1} = a(2 \cdot 2^{-1}) = a$.

(c) If $a = b$, then $1/b = 1/a$. Otherwise $0 < a < b$, and $1/b < 1/a$ by Proposition 3.2.2 (b). In both cases $1/b \leq 1/a$.

(d) First, $ac \leq bc$: if $a = b$ or $c = 0$, the two sides are equal (for $c = 0$ both are 0); otherwise $a < b$ and $c > 0$, and $ac < bc$ by Proposition 2.4.3 (c). Next, $bc \leq bd$, by the same argument with the roles of the factors exchanged: $c \leq d$, and $b \geq 0$ since $0 \leq a \leq b$ (transitivity, part (a)). By transitivity of $\leq$ (part (a)), $ac \leq bd$.

■

Part (b) is used whenever a proof splits a tolerance: the sum of two quantities, each below $\varepsilon/2$, is below ε. The midpoint is the same fact read for two points: between any two elements of F lies a third. Part (d) is how a product of two small, or two bounded, quantities is estimated.

Warning. The non-negativity in (d) cannot be dropped. In $\mathbb{Q}$, take $a = c = -2$ and $b = d = 0$: then $a \leq b$ and $c \leq d$, but $ac = 4$ and $bd = 0$, and $4 \leq 0$ is false. Before multiplying two inequalities, check that both sides are non-negative.

4.1.4 Absolute value

To say that x is close to a, we want one element of F measuring the gap between them, whichever of the two is larger. In school this is $|x - a|$, and $|x|$ is sometimes written $\sqrt{x^2}$. That formula needs a square-root function, and an ordered field has none in general: no rational squares to 2 (Theorem 3.3.1). The square x^2 has the two square roots x and $-x$, and picking the non-negative one is already a definition by cases. So we define the absolute value by cases, using only the order.

The absolute value of x is x with its sign made non-negative.

Definition 4.1.1. Absolute Value in an Ordered Field

Let F be an ordered field. For $x \in F$, the **absolute value** of x is

$$|x| \coloneqq \begin{cases} x & \text{if } x \geq 0, \\ -x & \text{if } x < 0. \end{cases}$$

For $x, y \in F$, the element $|x - y|$ is the **distance** from x to y.

The two cases do not overlap and cover F, by trichotomy (O1), so $|x|$ is a well-defined element of F, and $x \mapsto |x|$ is a function $F \to F$.

Examples. In $\mathbb{Q}$, $|3/4| = 3/4$ and $|-3/4| = 3/4$, the second because $-3/4 < 0$ and $-(-3/4) = 3/4$. The distance from $1/3$ to $1/2$ is $|1/3 - 1/2| = |-1/6| = 1/6$. The degenerate case is $|0| = 0$. Part (a) below shows that 0 is the only element with absolute value 0, so the distance from x to y is 0 exactly when $x = y$.

↻ **Second reading** · *Why not x^2?*

A non-example by minimal change. Replace the definition by $x \mapsto x^2$. This map is also non-negative, zero only at 0, and multiplicative, so it passes (a) and the product rule of (b) below. It fails the triangle inequality (e): in $\mathbb{Q}$, with $x = y = 1$, the "size" of $x + y$ would be $2^2 = 4$, while the sizes of x and y add up to $1 + 1 = 2$. A measure of size must not grow faster than its argument, and that is the clause the square breaks.

Proposition 4.1.3. Rules for the Absolute Value

Let F be an ordered field, and let $x, y, z, r \in F$.

(a) $|x| \geq 0$, and $|x| = 0$ if and only if $x = 0$.
(b) $|-x| = |x|$, $|xy| = |x||y|$ and $|x|^2 = x^2$; if $x \neq 0$, then $|1/x| = 1/|x|$.

(c) $-|x| \le x \le |x|$.

(d) $|x| \le r$ if and only if $-r \le x \le r$; and $|x| < r$ if and only if $-r < x < r$.

(e) *(Triangle inequality.)* $|x+y| \le |x| + |y|$, and $|x-z| \le |x-y| + |y-z|$.

(f) *(Reverse triangle inequality.)* $\big||x| - |y|\big| \le |x-y|$.

Idea. Parts (a)–(d) are case checks on the sign of x (and of y). The triangle inequality then needs **no** cases: by (c), x and y are each trapped between $\pm$ their absolute values; adding the two traps traps $x+y$ between $\pm(|x|+|y|)$, and (d) reads that as the inequality. The reverse triangle inequality is the triangle inequality applied to $x = (x-y)+y$ and to $y = (y-x)+x$.

Proof.

(a) If $x \ge 0$, then $|x| = x \ge 0$. If $x < 0$, then $|x| = -x > 0$ by Proposition 2.4.3 (a), since $-0 = 0$. So $|x| \ge 0$ always, and $|x| = 0$ only in the first case with $x = 0$; conversely $|0| = 0$.

(b) *Negation.* If $x > 0$, then $-x < 0$ (Proposition 2.4.3 (a)), so $|-x| = -(-x) = x = |x|$. If $x = 0$, both sides are 0. If $x < 0$, then $-x > 0$, so $|-x| = -x = |x|$.
Product. If $x \ge 0$ and $y \ge 0$, then $xy \ge 0$ (by (O4), or Proposition 2.3.2 (b) if a factor is 0), so $|xy| = xy = |x||y|$. If $x \ge 0$ and $y < 0$, then $-y > 0$ and $|x||y| = x(-y) = -(xy)$; and $-(xy) \ge 0$ by the first case, so $xy \le 0$ and $|xy| = -(xy)$ (if $xy = 0$, both readings give 0). The case $x < 0 \le y$ is the same with the roles exchanged. If $x < 0$ and $y < 0$, then $xy = (-x)(-y) > 0$ by (O4), so $|xy| = xy = (-x)(-y) = |x||y|$.
Square. $|x|^2$ is xx or $(-x)(-x)$, and both are x^2.
Reciprocal. Let $x \neq 0$. By the product rule, $|x|\,|1/x| = |x \cdot x^{-1}| = |1| = 1$, and $|x| \neq 0$ by (a). So $|1/x|$ satisfies the defining equation of $|x|^{-1}$, and Lemma 4.1.1 (a) gives $|1/x| = 1/|x|$.

(c) If $x \ge 0$, then $-|x| = -x \le 0 \le x = |x|$, using Proposition 2.4.3 (a) for $-x \le 0$. If $x < 0$, then $-|x| = -(-x) = x$, and $x < 0 < -x = |x|$. In both cases $-|x| \le x \le |x|$.

(d) ($\Rightarrow$) Let $|x| \le r$. By (c), $x \le |x| \le r$. By (c) and (b) applied to $-x$, $-x \le |-x| = |x| \le r$, and negating reverses the inequality (Proposition 2.4.3 (a), or equality), so $-r \le x$. ($\Leftarrow$) Let $-r \le x \le r$. If $x \ge 0$, then $|x| = x \le r$. If $x < 0$, then $|x| = -x \le r$, by negating $-r \le x$. The strict version is the same argument with $<$ in place of $\le$ at the steps that involve r: for instance $x \le |x| < r$ gives $x < r$ by transitivity.

(e) By (c), $-|x| \le x \le |x|$ and $-|y| \le y \le |y|$. Adding (Proposition 2.4.3 (b)),

$$-\big(|x| + |y|\big) \le x + y \le |x| + |y|,$$

the left end because $-(u+v) = (-u)+(-v)$. By (d) with $r := |x|+|y|$, $|x+y| \le |x|+|y|$. For the second form, apply the first to $x - z = (x-y) + (y-z)$.

(f) By (e), $|x| = |(x-y)+y| \le |x-y| + |y|$, so $|x| - |y| \le |x-y|$ by (O3) (or equality). Exchanging x and y, and using $|y-x| = |-(x-y)| = |x-y|$ from (b), gives $|y| - |x| \le |x-y|$. So $-|x-y| \le -(|y| - |x|)$ by Proposition 2.4.3 (a) (or equality), and $-(|y| - |x|) = |x| - |y|$, since $-(u+v) = (-u) + (-v)$; that is, $-|x-y| \le |x| - |y|$. By (d), $\big||x| - |y|\big| \le |x-y|$. This proves the proposition.

■

The form of (d) that the rest of the chapter uses is about a distance. Apply the strict half of (d) to $x - a$ with $r = \varepsilon$, and add a throughout; adding $-a$ goes back, so the step is reversible:

$$|x - a| < \varepsilon \iff a - \varepsilon < x < a + \varepsilon.$$

In words: x lies within ε of a exactly when it lies strictly between $a - \varepsilon$ and $a + \varepsilon$. In Section 4.3 the role of x is played by the terms a_n of a sequence and a is a candidate limit.

Example. In $\mathbb{Q}$, which x satisfy $|x - 1/2| < 1/4$? By the display above, exactly those with $1/2 - 1/4 < x < 1/2 + 1/4$, that is, $1/4 < x < 3/4$. So $2/3$ qualifies and $3/4$ does not: the inequality is strict, and $|3/4 - 1/2| = 1/4$.

Quick check. Is $|a| - |b| \le |a - b|$ for all $a, b \in F$? Is $|a - b| \le |a| - |b|$ for all $a, b \in F$?

Solution. The first: yes. By (f) and (c), $|a| - |b| \le ||a| - |b|| \le |a - b|$. The second: no. In $\mathbb{Q}$, take $a = 0$ and $b = 1$: then $|a - b| = |-1| = 1$, while $|a| - |b| = -1$, and $1 \le -1$ is false.

Quick check. For which $x \in \mathbb{Q}$ do both $|x - 1| < 2$ and $|x + 1| < 2$ hold?

Solution. By the display above, $|x - 1| < 2$ says $-1 < x < 3$, and $|x + 1| = |x - (-1)| < 2$ says $-3 < x < 1$. Both hold exactly when $-1 < x < 1$.

The triangle inequality is the tool of the whole chapter. In its second form it says that a detour through a third point y cannot shorten the way from x to z. Run backwards, it turns two small distances, from x to y and from y to z, into one small distance from x to z. That is the move of almost every proof in Sections 4.3–4.5.

4.1.5 Exercises

4.1.5.1 A. Check your understanding

Exercise. A1

Let F be an ordered field.

(a) Here is a proof that $0 < 1/2 < 1$ in F: (1) $1 > 0$; (2) $2 = 1 + 1 > 1 > 0$; (3) $0 < 1/2 < 1/1 = 1$. Name the rule behind each of the three steps.

(b) True or false in every ordered field: "if $a < b$, then $a^2 < b^2$". Give a reason, or a counterexample in $\mathbb{Q}$.

(c) True or false in every ordered field: "if $|a| < |b|$, then $a < b$". Give a reason, or a counterexample in $\mathbb{Q}$.

(d) True or false in every ordered field: "$x^2 + 1 > 0$ for every x". Give a reason, or a counterexample in $\mathbb{Q}$.

Solution.

(a) (a) Proposition 2.4.3 (e), since $1 \neq 0$ in a field. (2) $1+1 > 0+1 = 1$ by (O3) (or Proposition 2.4.3 (b), strict), then $2 > 0$ by (O2). (3) Proposition 3.2.2 (b), applied to $0 < 1 < 2$; and $1/1 = 1 \cdot 1^{-1} = 1$, since 1 is its own inverse ($1 \cdot 1 = 1$, and Lemma 4.1.1 (a)).

(b) False. In $\mathbb{Q}$, $-2 < 1$, but $(-2)^2 = 4 > 1 = 1^2$. The true statement needs $a, b \geq 0$: Proposition 2.4.3 (g).

(c) False. In $\mathbb{Q}$, $a = 1$ and $b = -2$ give $|a| = 1 < 2 = |b|$, but $a = 1 > -2 = b$.

(d) True. By Proposition 2.4.3 (e), $x^2 > 0$ if $x \neq 0$, and $x^2 = 0$ if $x = 0$ (Proposition 2.3.2 (b)); so $x^2 \geq 0$. Also $1 > 0$. Hence $x^2 + 1 > 0 + 0 = 0$ by Proposition 2.4.3 (b), strict in the second summand.

4.1.5.2 B. Practice

Exercise. B1: Two companions of (c) and (d)

Let F be an ordered field.

(a) Prove that if $0 \leq a < b$ and $0 < c \leq d$, then $ac < bd$.

(b) Prove that for $a, b \in F_{>0}$, $a \leq b$ **if and only if** $1/b \leq 1/a$.

Name the rule behind each step.

Solution.

(a) Since $0 \leq a < b$, $b > 0$ by Proposition 2.4.3 (i). First, $ac < bc$: $a < b$ and $c > 0$, so Proposition 2.4.3 (c) applies. Next, $bc \leq bd$: if $c = d$ the two sides are equal; otherwise $c < d$ and $b > 0$, so $cb < db$ by Proposition 2.4.3 (c), that is, $bc < bd$ by (F6). Combining $ac < bc \leq bd$ by Proposition 2.4.3 (i) gives $ac < bd$.

(b) ($\Rightarrow$) is Proposition 4.1.2 (c). ($\Leftarrow$) Let $1/b \leq 1/a$. Both are positive, by Proposition 3.2.2 (a). Applying Proposition 4.1.2 (c) to the positive elements $1/b \leq 1/a$ gives $1/(1/a) \leq 1/(1/b)$, and $1/(1/a) = (a^{-1})^{-1} = a$, $1/(1/b) = b$ by Lemma 4.1.1 (a). So $a \leq b$.

Exercise. B3: Two more triangle inequalities

Prove that for all a, b, c in an ordered field,

(a) $|a + b + c| \leq |a| + |b| + |c|$;

(b) $||a| - |b|| \leq |a + b|$.

Solution.

(a) By Proposition 4.1.3 (e) twice, $|(a+b)+c| \leq |a+b| + |c| \leq (|a| + |b|) + |c|$, the second step by adding $|c|$ to $|a+b| \leq |a|+|b|$ (Proposition 2.4.3 (b)). Conclude by transitivity (Proposition 2.4.3 (i)).

(b) Apply Proposition 4.1.3 (f) to a and $-b$: $||a| - |-b|| \leq |a - (-b)|$. By Proposition 4.1.3

(b), $|-b| = |b|$, and $a - (-b) = a + b$ by Proposition 2.3.2 (a). This is the claim.

4.1.5.3 C. Going deeper

Exercise. C1: A field that cannot be ordered

(a) Prove that in an ordered field no x satisfies $x^2 = -1$.

(b) Deduce that if a field contains an element i with $i^2 = -1$, then no relation $<$ makes it an ordered field.

Solution (Hint 1). Which sign does a square have, and which sign does -1 have?

Solution (Hint 2). Show first that $x^2 \geq 0$ for every x, splitting on whether $x = 0$.

Solution.

(a) Let F be an ordered field and $x \in F$. If $x = 0$, then $x^2 = 0$ (Proposition 2.3.2 (b)); if $x \neq 0$, then $x^2 > 0$ (Proposition 2.4.3 (e)). So $x^2 \geq 0$. On the other hand $1 > 0$ (Proposition 2.4.3 (e), as $1 \neq 0$ in a field), so $-1 < 0$ by Proposition 2.4.3 (a). If $x^2 = -1$, then $0 \leq x^2 = -1 < 0$, so $0 < 0$ by Proposition 2.4.3 (i), against (O1).

(b) If some relation $<$ made the field an ordered field, then (a) would forbid $i^2 = -1$. So no such relation exists.

The complex numbers of *Analytic Preliminaries*, Chapter 1 contain such an i, so they form a field that cannot be ordered. Compare Example (A field with two elements that cannot be ordered), where the obstacle was $1 + 1 = 0$; here it is that -1 is a square.

Marking scheme. (10 points) - 4: shows $x^2 \geq 0$ for every x (cases $x = 0$, $x \neq 0$), or rules out $x = 0$ for the x with $x^2 = -1$ and then shows $x^2 > 0$, giving the rule for each step. - 2: shows $-1 < 0$ from $1 > 0$ (from $1 \neq 0$ and Proposition 2.4.3 (e), or the standing setup of Section 4.1) and from negation reversing order. - 2: derives the contradiction with trichotomy (O1) or irreflexivity. - 2: (b) states that any order would make (a) apply, so none exists. Fatal errors: using strictness on $x^2 > 0$ without ruling out $x = 0$; asserting $1 > 0$ with no reason; using $\mathbb{R}$ or $\sqrt{}$.

Exercise. C2: Maximum and minimum through the absolute value

For a, b in an ordered field F, let $\max\{a, b\}$ be b if $a \leq b$ and a otherwise, and $\min\{a, b\}$ be a if $a \leq b$ and b otherwise.

(a) Prove that

$$\max\{a, b\} = \frac{a + b + |a - b|}{2}, \qquad \min\{a, b\} = \frac{a + b - |a - b|}{2}.$$

(b) Deduce that $|x| = \max\{x, -x\}$ for every $x \in F$.

Solution.

(a) *Case* $a \leq b$. Then $a - b \leq 0$ (Proposition 2.4.3 (a), or equality), so $|a - b| = b - a$: by the second clause of Definition 4.1.1 if $a - b < 0$, and because both sides are 0 if $a = b$. Hence

$$\frac{a + b + |a - b|}{2} = \frac{a + b + b - a}{2} = \frac{2b}{2} = b = \max\{a, b\},$$

and likewise $\dfrac{a + b - (b - a)}{2} = \dfrac{2a}{2} = a = \min\{a, b\}$, using $b + b = 2b$ and $(2b)2^{-1} = b$ as in the proof of Proposition 4.1.2 (b).
Case $b < a$. Then $a - b > 0$, so $|a - b| = a - b$, and the two formulas give $(2a)/2 = a = \max\{a, b\}$ and $(2b)/2 = b = \min\{a, b\}$.

(b) By (a) with $a = x$ and $b = -x$: $\max\{x, -x\} = (x + (-x) + |2x|)/2 = |2x|/2$. By Proposition 4.1.3 (b), $|2x| = |2||x| = 2|x|$, since $2 > 0$. So $\max\{x, -x\} = |x|$.