

24.4 Instrumental variables

The corrections so far used a *number* from outside the regression. Instrumental variables use a *variable*, related to the mismeasured regressor and unrelated to everything that makes least squares fail. The idea goes back to Philip Wright (1928), who used it to estimate supply and demand curves.

24.4.1 Endogeneity

By Eq. 24.1.2, $Y = \alpha + \beta W + e$ with $e = \varepsilon - \beta U$ correlated with W . An omitted common cause (Theorem 19.1.2) or a regressor partly determined by the response has the same effect. Such regressors are called **endogenous**. In every case the equation

$$y_i = \mathbf{x}_{(i)}^\top \boldsymbol{\beta} + e_i, \quad \mathbb{E}(\mathbf{x}_{(i)} e_i) \neq \mathbf{0}, \quad (24.4.1)$$

describes a relation of scientific interest, not the best linear predictor that least squares estimates. Here $\mathbf{x}_{(i)}$ holds all the regressors, including the constant. In this section $\mathbf{x}$ collects all regressors and $\mathbf{z}$ all instruments; the exactly measured covariates of earlier sections appear in both.

Definition 24.4.1. Instruments

A random q -vector $\mathbf{z}$ is a vector of **instruments** for Eq. 24.4.1, with random regressor p -vector $\mathbf{x}$, if

- (i) *exogeneity*: $\mathbb{E}(\mathbf{z}e) = \mathbf{0}$;
- (ii) *relevance*: $\boldsymbol{\Sigma}_{zx} = \mathbb{E}(\mathbf{z}\mathbf{x}^\top)$ has rank p , which requires $q \geq p$;

and $\boldsymbol{\Sigma}_{zz} = \mathbb{E}(\mathbf{z}\mathbf{z}^\top)$ is positive definite.

Regressors uncorrelated with e , such as the constant, are their own instruments; the rank condition asks the *excluded* instruments to carry extra information about the endogenous regressors. With $q = p$ the equation is **just identified**, and with $q > p$ **overidentified**.

In the blood-pressure study the second reading instruments the first: $\mathbf{x} = (1, W_1, \text{age})$, $\mathbf{z} = (1, W_2, \text{age})$ and $e = \varepsilon - \beta U_1$, which is uncorrelated with $W_2 = X + U_2$. Nothing about σ_u^2 needs to be known, but the two errors must be *independent*, which rules out two readings that share a transient cause (Exercise (A2)).

24.4.2 The estimators

With the $n \times p$ matrix $\mathbf{X}$ of regressors and the $n \times q$ matrix $\mathbf{Z}$ of instruments, the sample version of $\mathbb{E}\{\mathbf{z}(y - \mathbf{x}^\top \boldsymbol{\beta})\} = \mathbf{0}$ in the just-identified case is $\mathbf{Z}^\top(\mathbf{y} - \mathbf{X}\boldsymbol{\beta}) = \mathbf{0}$, with solution

$$\hat{\boldsymbol{\beta}}_{\text{IV}} = (\mathbf{Z}^\top \mathbf{X})^{-1} \mathbf{Z}^\top \mathbf{y}.$$

When $q > p$ these are more equations than unknowns. **Two-stage least squares** (2SLS) combines them through the projection $\mathbf{M}_Z = \mathbf{Z}(\mathbf{Z}^\top \mathbf{Z})^{-1} \mathbf{Z}^\top$ onto $\mathcal{C}(\mathbf{Z})$:

$$\hat{\boldsymbol{\beta}}_{2\text{SLS}} = (\mathbf{X}^\top \mathbf{M}_Z \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{M}_Z \mathbf{y}. \quad (24.4.2)$$

The name comes from a two-step computation.

Lemma 24.4.1. Two-stage least squares as projection

Let $\mathbf{Z}$ have full column rank and $\mathbf{X}^\top \mathbf{M}_\mathbf{Z} \mathbf{X}$ be nonsingular, and let $\hat{\mathbf{X}} = \mathbf{M}_\mathbf{Z} \mathbf{X}$ be the matrix of first-stage fitted values, obtained by regressing each column of $\mathbf{X}$ on $\mathbf{Z}$.

- (a) $\hat{\beta}_{2SLS}$ is the least squares coefficient of $\mathbf{y}$ on $\hat{\mathbf{X}}$, and also the IV estimator with instruments $\hat{\mathbf{X}}$: $\hat{\beta}_{2SLS} = (\hat{\mathbf{X}}^\top \hat{\mathbf{X}})^{-1} \hat{\mathbf{X}}^\top \mathbf{y} = (\hat{\mathbf{X}}^\top \mathbf{X})^{-1} \hat{\mathbf{X}}^\top \mathbf{y}$.
- (b) Columns of $\mathbf{X}$ that are also columns of $\mathbf{Z}$ are unchanged by the first stage.
- (c) If $q = p$ and $\mathbf{Z}^\top \mathbf{X}$ is nonsingular, $\hat{\beta}_{2SLS} = \hat{\beta}_{IV}$.

Proof. (a) $\mathbf{M}_\mathbf{Z}$ is the orthogonal projection onto $\mathcal{C}(\mathbf{Z})$ (Theorem 6.3.6), so it is symmetric and idempotent (Theorem 6.3.3). Hence $\hat{\mathbf{X}}^\top \hat{\mathbf{X}} = \mathbf{X}^\top \mathbf{M}_\mathbf{Z} \mathbf{M}_\mathbf{Z} \mathbf{X} = \mathbf{X}^\top \mathbf{M}_\mathbf{Z} \mathbf{X}$, and $\hat{\mathbf{X}}^\top \mathbf{X} = \mathbf{X}^\top \mathbf{M}_\mathbf{Z} \mathbf{X}$ and $\hat{\mathbf{X}}^\top \mathbf{y} = \mathbf{X}^\top \mathbf{M}_\mathbf{Z} \mathbf{y}$ as well. Substitute these in Eq. 24.4.2. (b) A column in $\mathcal{C}(\mathbf{Z})$ is fixed by $\mathbf{M}_\mathbf{Z}$. (c) With $\mathbf{Z}^\top \mathbf{X}$ square and invertible, $(\mathbf{X}^\top \mathbf{Z}(\mathbf{Z}^\top \mathbf{Z})^{-1} \mathbf{Z}^\top \mathbf{X})^{-1} = (\mathbf{Z}^\top \mathbf{X})^{-1}(\mathbf{Z}^\top \mathbf{Z})(\mathbf{X}^\top \mathbf{Z})^{-1}$, and multiplying by $\mathbf{X}^\top \mathbf{Z}(\mathbf{Z}^\top \mathbf{Z})^{-1} \mathbf{Z}^\top \mathbf{y}$ leaves $(\mathbf{Z}^\top \mathbf{X})^{-1} \mathbf{Z}^\top \mathbf{y}$. ■

The first stage replaces each endogenous regressor by the part of it that the instruments explain, which is uncorrelated with e . As with the residualized regressions of Section 6.6, the second-stage *coefficients* are right but its *residuals* $\mathbf{y} - \hat{\mathbf{X}}\hat{\beta}$ are not; the equation's residuals are $\mathbf{y} - \mathbf{X}\hat{\beta}$.

Theorem 24.4.2. Consistency and asymptotic normality of IV

Let $(\mathbf{x}_{(i)}, \mathbf{z}_{(i)}, e_i)$ be independent copies of $(\mathbf{x}, \mathbf{z}, e)$ with finite second moments, satisfying Eq. 24.4.1 and Definition 24.4.1. Let $\mathbf{H} = \Sigma_{xz} \Sigma_{zz}^{-1} \Sigma_{zx}$, where $\Sigma_{xz} = \Sigma_{zx}^\top$.

- (a) $\hat{\beta}_{2SLS} \rightarrow \beta$ with probability one.
- (b) If $E(e^2 \|\mathbf{z}\|^2) < \infty$, then $\sqrt{n}(\hat{\beta}_{2SLS} - \beta) \rightarrow N_p(\mathbf{0}, \mathbf{V})$ in distribution, where

$$\mathbf{V} = \mathbf{H}^{-1} \Sigma_{xz} \Sigma_{zz}^{-1} \Omega \Sigma_{zz}^{-1} \Sigma_{zx} \mathbf{H}^{-1}, \quad \Omega = E(e^2 \mathbf{z} \mathbf{z}^\top).$$

If $E(e^2 \mid \mathbf{z}) = \sigma_e^2$ is constant, then $\mathbf{V} = \sigma_e^2 \mathbf{H}^{-1}$.

- (c) $\hat{\sigma}_e^2 = n^{-1} \|\mathbf{y} - \mathbf{X}\hat{\beta}_{2SLS}\|^2 \rightarrow \sigma_e^2 = E(e^2)$ with probability one.

Proof. (a) Write $\hat{\mathbf{C}} = n^{-1} \mathbf{Z}^\top \mathbf{X}$ and $\hat{\mathbf{S}} = n^{-1} \mathbf{Z}^\top \mathbf{Z}$. Substituting $\mathbf{y} = \mathbf{X}\beta + \boldsymbol{\varepsilon}$ in Eq. 24.4.2,

$$\hat{\beta}_{2SLS} - \beta = (\hat{\mathbf{C}}^\top \hat{\mathbf{S}}^{-1} \hat{\mathbf{C}})^{-1} \hat{\mathbf{C}}^\top \hat{\mathbf{S}}^{-1} \frac{1}{n} \mathbf{Z}^\top \boldsymbol{\varepsilon}.$$

By the strong law, $\hat{\mathbf{C}} \rightarrow \Sigma_{zx}$, $\hat{\mathbf{S}} \rightarrow \Sigma_{zz}$ and $n^{-1} \mathbf{Z}^\top \boldsymbol{\varepsilon} \rightarrow E(\mathbf{z}e) = \mathbf{0}$. The matrix $\mathbf{H}$ is positive definite: for $\mathbf{a} \neq \mathbf{0}$, $\Sigma_{zx} \mathbf{a} \neq \mathbf{0}$ by the rank condition, so $\mathbf{a}^\top \mathbf{H} \mathbf{a} = (\Sigma_{zx} \mathbf{a})^\top \Sigma_{zz}^{-1} (\Sigma_{zx} \mathbf{a}) > 0$. By continuity the right side tends to $\mathbf{H}^{-1} \Sigma_{xz} \Sigma_{zz}^{-1} \mathbf{0} = \mathbf{0}$.

(b) Multiply the display by $\sqrt{n}$. The vectors $\mathbf{z}_{(i)} e_i$ are independent and identically distributed with mean $\mathbf{0}$ and covariance Ω , finite by assumption, so $n^{-1/2} \mathbf{Z}^\top \boldsymbol{\varepsilon} \rightarrow N_q(\mathbf{0}, \Omega)$ by the central limit theorem. The matrix in front converges to $\mathbf{L} = \mathbf{H}^{-1} \Sigma_{xz} \Sigma_{zz}^{-1}$, and the splitting used for Theorem 24.3.1(b) gives the limit $\mathbf{L} N_q(\mathbf{0}, \Omega) = N_p(\mathbf{0}, \mathbf{V})$. If $E(e^2 \mid \mathbf{z}) = \sigma_e^2$, then $\Omega = E\{E(e^2 \mid \mathbf{z}) \mathbf{z} \mathbf{z}^\top\} = \sigma_e^2 \Sigma_{zz}$, and $\mathbf{V}$ collapses to $\sigma_e^2 \mathbf{H}^{-1}$.

(c) With $\delta = \hat{\beta}_{2\text{SLS}} - \beta$, $y - X\hat{\beta}_{2\text{SLS}} = \varepsilon - X\delta$, so

$$\hat{\sigma}_e^2 = \frac{1}{n} \varepsilon^\top \varepsilon - 2\delta^\top \frac{1}{n} X^\top \varepsilon + \delta^\top \left(\frac{1}{n} X^\top X \right) \delta.$$

The averages converge to $E(e^2)$, $E(xe)$ and $E(xx^\top)$, all finite, and $\delta \rightarrow 0$ by (a).

■

In practice V is estimated with sample moments and $\hat{\Omega} = n^{-1} \sum_i \hat{e}_i^2 \mathbf{z}_{(i)} \mathbf{z}_{(i)}^\top$, as in Theorem 21.4.2, or under homoscedasticity by $\hat{\sigma}_e^2 (X^\top M_Z X)^{-1}$. Among estimators using linear combinations of the instruments, 2SLS has the smallest asymptotic covariance under homoscedasticity ([Exercise \(C2\)](#)).

Example. The second reading as an instrument

Instrument the first reading with the second, and let age and the constant instrument themselves. The IV estimates are 0.0708 for blood pressure, with standard error 0.0085 (0.0081 from the sandwich), and 0.0054 for age (standard error 0.0081). The spurious age effect has gone. The first-stage F statistic for the second reading is 160.1, a strong instrument. The correct residuals give $\hat{\sigma}_e^2 = 1.438$, estimating $\sigma^2 + \beta^2 \sigma_u^2 = 1.5184$; the second-stage residuals give 1.376. The moment correction of Example (Correcting with two readings) gave 0.0757 with standard error 0.0076. It uses both readings symmetrically, while IV uses one only as an instrument and pays in precision. But IV needs no estimate of the error variance, and the instrument need only be correlated with X , not a second unbiased reading of it.

```
import numpy as np

rng = np.random.default_rng(2401)
n = 300
age = rng.normal(55, 10, n)
x = 130 + 0.6 * (age - 55) + rng.normal(0, np.sqrt(108), n)  # long-run blood
                        pressure, sd 12
w = x[:, None] + rng.normal(0, 9, (n, 2))                  # two clinic readings
y = -4 + 0.08 * x + 0.0 * age + rng.normal(0, 1.0, n)      # age has no effect
                        given x

X = np.column_stack([np.ones(n), w[:, 0], age])            # regressors: first reading, age
Z = np.column_stack([np.ones(n), w[:, 1], age])            # instruments: second reading,
                        age
b_iv = np.linalg.solve(Z.T @ X, Z.T @ y)
e = y - X @ b_iv                                          # residuals use X, not the first
                        -stage fit
ZXinv = np.linalg.inv(Z.T @ X)
cov_h = e @ e / n * ZXinv @ (Z.T @ Z) @ ZXinv.T          # homoscedastic
cov_s = ZXinv @ ((Z * e[:, None] ** 2).T @ Z) @ ZXinv.T   # heteroscedasticity-
                        robust
print("IV (1, x, age):", b_iv.round(4))
print("se, homoscedastic:", np.sqrt(np.diag(cov_h)).round(4))
print("se, sandwich:      ", np.sqrt(np.diag(cov_s)).round(4))

# first stage: regress the first reading on the instruments; F for the excluded
                        instrument
g, rss1 = np.linalg.lstsq(Z, w[:, 0], rcond=None)[:2]
rss0 = np.sum((w[:, 0] - np.column_stack([np.ones(n), age]) @
```

```

np.linalg.lstsq(np.column_stack([np.ones(n), age]), w[:, 0], rcond=
None)[0]) ** 2)
F1 = (rss0 - rss1[0]) / (rss1[0] / (n - 3))
print(f"first-stage F = {F1:.1f}")

```

[Run this code online](#)

Wald's (1940) grouping estimator ([Section 7.1](#)) is an IV estimator whose instrument is a group indicator ([Exercise \(B1\)](#)).

24.4.3 Weak instruments

If the instruments explain little of the endogenous regressor, the limit theory of [Theorem 24.4.2](#) is a poor guide at realistic n . The simplest case can be solved exactly.

Proposition 24.4.3. The exact distribution with one instrument

Let $y_i = \beta x_i + e_i$ and $x_i = \pi z_i + v_i$, $i = 1, \dots, n$, with fixed z_i , $S_{zz} = \sum_i z_i^2 > 0$, and (e_i, v_i) independent bivariate normal with means zero, standard deviations σ_e, σ_v and correlation ρ , $|\rho| < 1$. Let $\hat{\beta} = \sum_i z_i y_i / \sum_i z_i x_i$ and $\mu = \pi \sqrt{S_{zz}} / \sigma_v$.

(a) $\hat{\beta} - \beta$ has the same distribution as

$$\frac{\sigma_e}{\sigma_v} \cdot \frac{\rho\eta + \sqrt{1 - \rho^2} \xi}{\mu + \eta}, \quad \eta, \xi \stackrel{\text{iid}}{\sim} N(0, 1). \quad (24.4.3)$$

(b) $E|\hat{\beta}| = \infty$ for every μ .

(c) As $\mu \rightarrow \infty$, $(\mu\sigma_v/\sigma_e)(\hat{\beta} - \beta) \rightarrow N(0, 1)$ in distribution. If $\mu = 0$, $\hat{\beta} - \beta$ has median $\rho\sigma_e/\sigma_v$, which is the large-sample bias of least squares when $\pi = 0$.

(d) $(\sum_i z_i x_i)^2 / (\sigma_v^2 S_{zz}) \sim \chi^2(1, \mu^2)$, with mean $1 + \mu^2$.

Proof. (a) Put $\eta = \sum_i z_i v_i / (\sigma_v \sqrt{S_{zz}})$ and $\zeta = \sum_i z_i e_i / (\sigma_e \sqrt{S_{zz}})$. They are jointly normal (Theorem 3.3.1) with unit variances and covariance $\sum_i z_i^2 \rho \sigma_e \sigma_v / (\sigma_e \sigma_v S_{zz}) = \rho$. By Theorem 3.5.2, $\xi = (\zeta - \rho\eta) / \sqrt{1 - \rho^2}$ is standard normal and independent of η . Now $\sum_i z_i x_i = \pi S_{zz} + \sigma_v \sqrt{S_{zz}} \eta = \sigma_v \sqrt{S_{zz}} (\mu + \eta)$, and $\hat{\beta} - \beta = \sum_i z_i e_i / \sum_i z_i x_i = \sigma_e \zeta / \{\sigma_v (\mu + \eta)\}$.

(b) Given η , the numerator $\rho\eta + \sqrt{1 - \rho^2} \xi$ is normal with standard deviation $\sqrt{1 - \rho^2} > 0$, so its conditional absolute mean is at least $c = \sqrt{2(1 - \rho^2)}/\pi$. Hence $E|\hat{\beta} - \beta| \geq (\sigma_e/\sigma_v) c E|\mu + \eta|^{-1} = \infty$, because the density of $\mu + \eta$ is continuous and positive at zero, as in the proof of Proposition 12.5.2. Finally $|\hat{\beta}| \geq |\hat{\beta} - \beta| - |\beta|$.

(c) $(\mu\sigma_v/\sigma_e)(\hat{\beta} - \beta)$ has the law of $(\rho\eta + \sqrt{1 - \rho^2} \xi) / (1 + \eta/\mu)$. The numerator is standard normal and $\eta/\mu \rightarrow 0$ in probability, so Slutsky's lemma applies. For $\mu = 0$ the representation is $(\sigma_e/\sigma_v)(\rho + \sqrt{1 - \rho^2} \xi/\eta)$, and ξ/η is symmetric about zero, so the median is $\rho\sigma_e/\sigma_v$. When $\pi = 0$, $x = v$ and least squares converges to $\beta + \text{Cov}(v, e)/\sigma_v^2 = \beta + \rho\sigma_e/\sigma_v$.

(d) $\sum_i z_i x_i / (\sigma_v \sqrt{S_{zz}}) = \mu + \eta \sim N(\mu, 1)$, and its square is $\chi^2(1, \mu^2)$, whose mean is $1 + \mu^2$ (the convention of [Chapter 4](#), without a factor $\frac{1}{2}$). ■

The design and π enter only through the **concentration parameter** μ^2 , which acts as a sample




Figure 24.4.1: Sampling distributions of the IV estimate of $\beta = 1$ for $\mu^2 = 1, 10, 100$ on $[-1, 3]$; dotted, the least squares limit.

size, and by (d) the first-stage statistic estimates $1 + \mu^2$. Hence the rule of thumb of Staiger and Stock (1997): an instrument is weak if the first-stage F is below about 10. A useless instrument ($\mu = 0$) gives an estimator centred on the least squares bias, with Cauchy tails.

Example. How weak is too weak?

With $n = 200$, $\beta = 1$, $\sigma_e = \sigma_v = 1$ and $\rho = 0.9$, least squares tends to about 1.9 for a weak instrument. In 20000 samples for each concentration:

μ^2	median	quartiles	coverage of nominal 95% Wald interval
1	1.298	0.711 to 1.646	0.843
10	1.008	0.740 to 1.181	0.921
100	1.002	0.929 to 1.064	0.947

With $\mu^2 = 1$ the median moves a third of the way to the least squares limit, a proportion 0.200 of estimates fall outside $[-1, 3]$, and the Wald interval undercovers, as it still does at $\mu^2 = 10$ (Figure 24.4.1).

```
def iv_draws(mu2, reps, seed, n=200, rho=0.9):
    """IV estimates of beta = 1 and Wald-interval coverage when the concentration is
    mu2."""
    r = np.random.default_rng(seed)
    z = r.normal(size=n)
    z = z * np.sqrt(n / (z @ z))          # fixed instrument with sum z^2 =
    n
    pi = np.sqrt(mu2 / n)
    ev = r.multivariate_normal([0, 0], [[1, rho], [rho, 1]], size=(reps, n))
    xx = pi * z + ev[:, :, 1]
    yy = 1.0 * xx + ev[:, :, 0]
    b = (yy @ z) / (xx @ z)
    res = yy - b[:, None] * xx
    se = np.sqrt((res ** 2).mean(axis=1) * n / (xx @ z) ** 2)
```

```

return b, np.abs(b - 1) <= 1.96 * se

for mu2 in [1, 10, 100]:
    b, hit = iv_draws(mu2, 2000, 2440)
    print(f"mu^2 = {mu2:3d}: median {np.median(b):.3f}, "
          f"quartiles {np.percentile(b, 25):.3f} to {np.percentile(b, 75):.3f}, "
          f"coverage {hit.mean():.3f}")

```

[Run this code online](#)

With many weak instruments, 2SLS is biased towards least squares, and Bound, Jaeger and Baker (1995) showed that it can nearly reproduce least squares; we do not prove this. For inference, invert a test whose null distribution does not depend on instrument strength, such as the Anderson–Rubin test ([Exercise C1](#)). Its confidence sets are of Fieller type (Proposition 12.5.1) and can be unbounded, as they must sometimes be (Proposition 12.5.4).

Warning. Exogeneity cannot be checked in a just-identified model: all the equations $E(\mathbf{z}e) = \mathbf{0}$ are used to estimate β . With more instruments, tests of overidentifying restrictions (Sargan 1958) check only that the instruments *agree*. An instrument is justified by knowledge of how the data arose; for replicates, by the independence of their errors. Instruments for causal effects are taken up in [Chapter 25](#).

24.4.4 Exercises

24.4.4.1 A. Check your understanding

Exercise. A1

For $\mathbf{x} = (1, W)$ and $\mathbf{z} = (1, Z)$, show that the IV slope is s_{zy}/s_{zw} , the ratio of two sample covariances. Show that the population ratio equals β when Z is correlated with X and uncorrelated with U and ε .

Exercise. A2

If the errors of two readings are correlated, $\text{Cov}(U_1, U_2) = \kappa$, find the limit of the IV slope in simple regression that instruments W_1 with W_2 . In which direction is it biased when $\kappa > 0$?

Solution. $\text{Cov}(W_2, Y) = \beta\sigma_x^2$ and $\text{Cov}(W_2, W_1) = \sigma_x^2 + \kappa$, so the limit is $\beta\sigma_x^2/(\sigma_x^2 + \kappa)$. With $\kappa > 0$ it is attenuated, though less than least squares unless $\kappa = \sigma_u^2$. The instrument is not exogenous: $\text{Cov}(W_2, \varepsilon - \beta U_1) = -\beta\kappa$.

24.4.4.2 B. Practice

Exercise. B1

Let $g_i \in \{0, 1\}$ be a group indicator, and let $\mathbf{z}_{(i)} = (1, g_i)$ instrument $\mathbf{x}_{(i)} = (1, w_i)$. Show that the IV slope is $(\bar{y}_1 - \bar{y}_0)/(\bar{w}_1 - \bar{w}_0)$, the difference of group means of y divided by that

of w . This is Wald's (1940) estimator. Show that it is consistent if g is independent of (U, ε) and $E(X | g)$ differs between the groups. Explain why forming the groups by splitting at the median of w itself, as Wald proposed, does not satisfy these conditions under classical error.

Solution. By [Exercise \(A1\)](#) the slope is s_{gy}/s_{gw} . For a binary g with n_1 ones, $\sum_i (g_i - \bar{g})y_i = n_1(\bar{y}_1 - \bar{y})$ and $\bar{y}_1 - \bar{y} = (n_0/n)(\bar{y}_1 - \bar{y}_0)$. The same holds for w , and the factors cancel. Exogeneity needs $\text{Cov}(g, \varepsilon - \beta U) = 0$, which holds when g is independent of (U, ε) . Relevance needs $\text{Cov}(g, W) = \text{Cov}(g, X) \neq 0$, that is, different group means of X . If the groups are formed from w , membership depends on U : readings with large positive errors tend to land in the upper group, so $\text{Cov}(g, U) > 0$ and exogeneity fails. The grouping must come from information independent of the measurement error, such as a second reading or a design variable.

24.4.4.3 C. Going deeper

Exercise. C1

In [Eq. 24.4.1](#) with $\mathbf{x} = (1, W)$, let $\mathbf{z} = (1, \mathbf{z}_1)$ with $\mathbf{z}_1$ a k -vector of excluded instruments, and suppose e is independent of $\mathbf{z}$ and $N(0, \sigma_e^2)$. To test $H_0 : \beta = \beta_0$, regress $y - \beta_0 w$ on $\mathbf{z}$ and compute the F statistic for $\mathbf{z}_1$. Show that under H_0 it has an exact $F(k, n - k - 1)$ distribution, whatever the strength of the instruments. This is the test of Anderson and Rubin (1949). Inverting it gives a confidence set for β . Show that for $k = 1$ the set is of the form in [Proposition 12.5.1](#).

Solution. Under H_0 , $y - \beta_0 w = \alpha + e$ with e normal and independent of $\mathbf{z}$: given $\mathbf{Z}$, a normal linear model in which the coefficients of $\mathbf{z}_1$ are zero. The F statistic of [Theorem 11.1.1](#) is exactly $F(k, n - k - 1)$ given $\mathbf{Z}$, hence unconditionally, and w plays no role. For $k = 1$, let $\hat{a}$ and $\hat{b}$ be the slopes of y and of w on $(1, z)$, let $v = 1 / \sum_i (z_i - \bar{z})^2$, and let s_{11} , s_{12} , s_{22} be the residual mean squares and mean cross-product of y and w from these two regressions. The regression of $y - \beta_0 w$ on $(1, z)$ has slope $\hat{a} - \beta_0 \hat{b}$ and residual mean square $s_{11} - 2\beta_0 s_{12} + \beta_0^2 s_{22}$, so $F = t^2$ and the acceptance region is

$$(\hat{a} - \beta_0 \hat{b})^2 \leq t_{n-2, \alpha/2}^2 v (s_{11} - 2\beta_0 s_{12} + \beta_0^2 s_{22}).$$

This is [Eq. 12.5.1](#) with $s^2 v_{ij}$ replaced by $v s_{ij}$: a quadratic inequality in β_0 . Like Fieller's set, it can be an interval, the complement of an interval, or the whole line.

Exercise. C2

Let $\mathbf{F}$ be a fixed $q \times p$ matrix with $\Sigma_{zx}^\top \mathbf{F}$ nonsingular, and consider the IV estimator with instruments $\mathbf{ZF}$. Under homoscedasticity its asymptotic covariance is $\sigma_e^2 (\mathbf{F}^\top \Sigma_{zx})^{-1} \mathbf{F}^\top \Sigma_{zz} \mathbf{F} (\Sigma_{zx}^\top \mathbf{F})^{-1}$. Show that this is never smaller, in the nonnegative definite ordering, than $\sigma_e^2 \mathbf{H}^{-1}$, with equality when $\mathbf{F} = \Sigma_{zz}^{-1} \Sigma_{zx}$, which is the population version of 2SLS.