

12.5 Inverse prediction and calibration

Prediction runs from x to y . Calibration runs the other way: an instrument is calibrated on specimens of known value, a new specimen is measured, and its value is wanted. The estimate reads the fitted line backwards; the interval reads the prediction band backwards, and it is not always an interval.

12.5.1 Ratios of estimable functions

Inverse prediction belongs to a wider problem: a confidence set for a *ratio* of linear functions of the coefficients, such as the zero crossing $-\beta_0/\beta_1$ of a line or the turning point $-\beta_1/(2\beta_2)$ of a quadratic. A delta-method interval ignores that the denominator is uncertain and may be near zero. Fieller's approach tests, for each candidate value, a hypothesis that is linear in the coefficients.

Proposition 12.5.1. Fieller's confidence set for a ratio

Let $\lambda_1, \lambda_2 \in \mathcal{C}(\mathbf{X}^\top)$ be linearly independent, put $a = \lambda_1^\top \beta$, $b = \lambda_2^\top \beta$, their estimates $\hat{a} = \lambda_1^\top \hat{\beta}$, $\hat{b} = \lambda_2^\top \hat{\beta}$, and $v_{ij} = \lambda_i^\top \mathbf{G} \lambda_j$, so that $\mathbf{V} = (v_{ij})$ is positive definite. Suppose $b \neq 0$ and let $\theta = a/b$. With $t = t_{n-r, \alpha/2}$, let

$$\mathcal{F}(\mathbf{Y}) = \{\theta_0 \in \mathbb{R} : (\hat{a} - \theta_0 \hat{b})^2 \leq t^2 s^2 (v_{11} - 2\theta_0 v_{12} + \theta_0^2 v_{22})\}. \quad (12.5.1)$$

- (a) $\mathcal{F}(\mathbf{Y})$ is an exact $1 - \alpha$ confidence set for θ , at every parameter value with $b \neq 0$.
- (b) Let $Q = (\hat{a}, \hat{b}) \mathbf{V}^{-1} (\hat{a}, \hat{b})^\top$. Outside an event of probability zero, $\mathcal{F}(\mathbf{Y})$ is
 - a bounded closed interval containing $\hat{\theta} = \hat{a}/\hat{b}$, if $\hat{b}^2 > t^2 s^2 v_{22}$;
 - the union of two disjoint closed rays, if $\hat{b}^2 < t^2 s^2 v_{22}$ and $Q > t^2 s^2$;
 - the whole real line, if $\hat{b}^2 < t^2 s^2 v_{22}$ and $Q < t^2 s^2$.

Proof.

- (a) Fix θ_0 . The vector $\lambda_1 - \theta_0 \lambda_2$ lies in $\mathcal{C}(\mathbf{X}^\top)$ and is not zero, so $(\lambda_1 - \theta_0 \lambda_2)^\top \beta = a - \theta_0 b$ is estimable, with estimate $\hat{a} - \theta_0 \hat{b}$ and $(\lambda_1 - \theta_0 \lambda_2)^\top \mathbf{G} (\lambda_1 - \theta_0 \lambda_2) = v_{11} - 2\theta_0 v_{12} + \theta_0^2 v_{22}$. The inequality in Eq. 12.5.1 says that the t interval of Theorem 12.1.1 for $a - \theta_0 b$ contains 0. At $\theta_0 = \theta$, $a - \theta b = 0$, so the probability of this event is exactly $1 - \alpha$.
- (b) The defining inequality is $q(\theta_0) = A\theta_0^2 - 2B\theta_0 + K \leq 0$, with

$$A = \hat{b}^2 - t^2 s^2 v_{22}, \quad B = \hat{a}\hat{b} - t^2 s^2 v_{12}, \quad K = \hat{a}^2 - t^2 s^2 v_{11}.$$

Expanding and using $(\hat{a}, \hat{b}) \mathbf{V}^{-1} (\hat{a}, \hat{b})^\top = (v_{22}\hat{a}^2 - 2v_{12}\hat{a}\hat{b} + v_{11}\hat{b}^2)/\det \mathbf{V}$,

$$\begin{aligned} B^2 - AK &= t^2 s^2 (v_{22}\hat{a}^2 - 2v_{12}\hat{a}\hat{b} + v_{11}\hat{b}^2 - t^2 s^2 \det \mathbf{V}) \\ &= t^2 s^2 \det(\mathbf{V}) (Q - t^2 s^2). \end{aligned}$$

If $A > 0$, then $q(\hat{\theta}) = -t^2 s^2 (v_{11} - 2\hat{\theta}v_{12} + \hat{\theta}^2 v_{22}) < 0$, so the upward parabola q has two real roots, and $\{q \leq 0\}$ is the closed interval between them, which contains $\hat{\theta}$. If $A < 0$, the parabola opens downward. When $B^2 - AK < 0$ it is negative everywhere and $\mathcal{F}(\mathbf{Y}) = \mathbb{R}$. When $B^2 - AK > 0$ it is nonpositive exactly outside the open interval

between its roots. The events $A = 0$ and $Q = t^2 s^2$ have probability zero, because $\hat{b}$ and Q/s^2 have continuous distributions. ■

The first case says that the t test rejects $b = 0$: the denominator is away from zero and the ratio is confined. Otherwise very large $|\theta|$ cannot be excluded, and $Q/(2s^2)$, the F statistic for $a = b = 0$, decides between two rays and the whole line. [Exercise \(C1\)](#) gives a picture: $\mathcal{F}(\mathbf{Y})$ is the set of slopes of lines through the origin that meet a certain confidence ellipse for (a, b) .

12.5.2 The calibration problem

A calibration experiment measures n standards with known values $x_1, \dots, x_n$, not all equal, and records readings

$$Y_i = \beta_0 + \beta_1 x_i + \varepsilon_i, \quad i = 1, \dots, n.$$

A new specimen with unknown value x_0 is then measured $m \geq 1$ times, $Y_{0j} = \beta_0 + \beta_1 x_0 + \varepsilon_{0j}$, and all $n + m$ errors are independent $N(0, \sigma^2)$. Let $\bar{Y}_0$ be the mean of the new readings, and let $\hat{\beta}_0$, $\hat{\beta}_1$ and s^2 (with $n - 2$ degrees of freedom) come from the standards alone. The **classical estimator** reads the fitted line backwards:

$$\hat{x}_0 = \frac{\bar{Y}_0 - \hat{\beta}_0}{\hat{\beta}_1} = \bar{x} + \frac{\bar{Y}_0 - \bar{Y}}{\hat{\beta}_1}.$$

It is the maximum likelihood estimator ([Exercise \(A2\)](#)) and, being a ratio, inherits a pathology of ratios.

Proposition 12.5.2. The classical estimator has no mean

If $\Pr(\bar{Y}_0 \neq \bar{Y}) > 0$, which always holds here, then $E|\hat{x}_0| = \infty$.

Proof. $\bar{Y}$ and $\hat{\beta}_1$ are jointly normal and uncorrelated (Corollary 5.5.4), hence independent (Theorem 3.4.1), and $\bar{Y}_0$ is independent of both. So $g = \bar{Y}_0 - \bar{Y}$ is independent of $\hat{\beta}_1$ and $E|\hat{x}_0 - \bar{x}| = E|g| E|1/\hat{\beta}_1|$, with $E|g| > 0$. The density f of $\hat{\beta}_1$ is continuous and positive at 0, so $f \geq c > 0$ on some $[-\epsilon, \epsilon]$ and $E|1/\hat{\beta}_1| \geq c \int_{-\epsilon}^{\epsilon} |u|^{-1} du = \infty$. Finally $|\hat{x}_0| \geq |\hat{x}_0 - \bar{x}| - |\bar{x}|$. ■

So a delta-method “standard error” of $\hat{x}_0$ approximates a quantity that does not exist. That is harmless when $\hat{\beta}_1$ is many standard errors from zero, and harmful exactly when the slope is poorly determined.

Theorem 12.5.3. Calibration set

In the calibration model, let $t = t_{n-2, \alpha/2}$ and

$$\begin{aligned} \mathcal{C} &= \left\{ x \in \mathbb{R} : (\bar{Y}_0 - \hat{\beta}_0 - \hat{\beta}_1 x)^2 \right. \\ &\quad \left. \leq t^2 s^2 \left(\frac{1}{m} + \frac{1}{n} + \frac{(x - \bar{x})^2}{S_{xx}} \right) \right\}. \end{aligned} \tag{12.5.2}$$

- (a) $\mathcal{C}$ is an exact $1 - \alpha$ confidence set for x_0 , at every $(\beta_0, \beta_1, \sigma^2, x_0)$, including $\beta_1 = 0$.
- (b) $\mathcal{C}$ is the set of x at which $\bar{Y}_0$ lies inside the $1 - \alpha$ prediction interval of Theorem 12.3.1(c) for the mean of m new readings at x .
- (c) Let $T_1 = \hat{\beta}_1 \sqrt{S_{xx}}/s$ be the t statistic for the slope and $g = \bar{Y}_0 - \bar{Y}$. Outside an event of probability zero, $\mathcal{C}$ is
- a bounded interval containing $\hat{x}_0$, if $T_1^2 > t^2$;
 - the union of two disjoint closed rays, if $T_1^2 < t^2$ and $T_1^2 + g^2/\{s^2(1/m + 1/n)\} > t^2$;
 - the whole real line, if $T_1^2 + g^2/\{s^2(1/m + 1/n)\} < t^2$.

Proof.

- (a) and (b). For fixed x , $\bar{Y}_0 - \hat{\beta}_0 - \hat{\beta}_1 x$ is the difference between the mean of the new readings and the estimated mean response at x . At $x = x_0$, Theorem 12.3.1(c), with $\mathbf{x}_0 = (1, x_0)^\top$ and $h_0 = 1/n + (x_0 - \bar{x})^2/S_{xx}$ (Eq. 5.5.2), shows that

$$\frac{\bar{Y}_0 - \hat{\beta}_0 - \hat{\beta}_1 x_0}{s\sqrt{1/m + 1/n + (x_0 - \bar{x})^2/S_{xx}}} \sim t(n-2),$$

whatever the parameters are. This is a pivot, and $x_0 \in \mathcal{C}$ is the event that its absolute value is at most t . Nothing in the argument requires $\beta_1 \neq 0$. Part (b) restates the definition.

- (b) Put $d = x - \bar{x}$ and $c = 1/m + 1/n$. Since $\bar{Y}_0 - \hat{\beta}_0 - \hat{\beta}_1 x = g - \hat{\beta}_1 d$, the defining inequality is $Ad^2 - 2\hat{\beta}_1 g d + (g^2 - t^2 s^2 c) \leq 0$ with $A = \hat{\beta}_1^2 - t^2 s^2/S_{xx}$. Its discriminant is

$$\begin{aligned} \hat{\beta}_1^2 g^2 - A(g^2 - t^2 s^2 c) &= t^2 s^2 \left(\frac{g^2}{S_{xx}} + cA \right) \\ &= \frac{t^2 s^4 c}{S_{xx}} \left(T_1^2 + \frac{g^2}{s^2 c} - t^2 \right). \end{aligned}$$

The sign of A is that of $T_1^2 - t^2$. If $A > 0$, the quadratic is negative at $d = g/\hat{\beta}_1$, that is at $x = \hat{x}_0$, where it equals $-t^2 s^2(c + d^2/S_{xx})$; so it has two real roots and $\mathcal{C}$ is the interval between them. If $A < 0$, the argument of Proposition 12.5.1(b) applies: two rays if the discriminant is positive and the whole line if it is negative. Equalities have probability zero. ■

Part (b) is the practical recipe (Figure 12.5.1): draw the prediction band for the mean of m readings and the horizontal line at $\bar{Y}_0$, and read off where the line is inside the band. A steep calibration line crosses the band in a short interval. A flat one lets both hyperbolic edges pass the horizontal line, giving two rays or everything. The set is bounded iff the slope test rejects, as it does for any useful instrument.

Proposition 12.5.4. Unbounded calibration sets cannot be avoided

Let $\mathcal{D}$ be any confidence set for x_0 with coverage at least $1 - \alpha > 0$ at every parameter value, including those with $\beta_1 = 0$. Then at every parameter value with $\beta_1 = 0$, $\Pr\{v \in \mathcal{D}\} \geq 1 - \alpha$ for every real v , and if $\mathcal{D}$ is measurable in the sense of Proposition 12.1.3, its expected length is infinite.

Proof. When $\beta_1 = 0$, the joint distribution of the standards and the new readings does not involve x_0 . So for every v , the probability that $v \in \mathcal{D}$ is the same as it would be if x_0 were v , and that probability is at least $1 - \alpha$. The rest is the Fubini argument in the proof of Proposition 12.1.3. ■

By continuity the same happens approximately for small β_1 . A procedure that always reports a bounded interval, like the delta method, pays with coverage below $1 - \alpha$ somewhere. The Fieller set's unbounded answers are the honest answer to a question the data cannot settle.

Example. Calibrating an assay

In a simulated assay, nine standards of concentration $0, 1, \dots, 8$ are each measured three times ($n = 27$). The fitted line is $0.0338 + 0.1161x$, with $s = 0.0131$ and slope t statistic 118.7. A new specimen is read $m = 3$ times, with mean reading 0.6521. The classical estimate is $\hat{x}_0 = 5.327$, and the 95% calibration interval is $(5.184, 5.471)$ ($t_{25,0.025} = 2.060$). For an assay this precise the delta-method interval, $\hat{x}_0 \pm t(s/|\hat{\beta}_1|)\sqrt{1/m + 1/n + (\hat{x}_0 - \bar{x})^2/S_{xx}}$, agrees to three decimals: $(5.184, 5.471)$.

Now consider a weak assay with the same design, in which the fitted slope is 0.0021 and its t statistic is only 1.67. The mean reading of the standards is 0.0475. A specimen with mean reading 0.0525, close to that mean, gives the whole real line: nothing about its concentration can be concluded. A specimen with mean reading 0.0775 gives two rays, $(-\infty, -55.3] \cup [7.6, \infty)$: the middle of the scale is excluded, but not extreme concentrations of either sign. Figure 12.5.1 shows both assays.

In 20000 simulated experiments at $x_0 = 5.3$, both intervals cover in 0.9517 for the precise assay. With the slope three standard errors from zero ($\beta_1 = 0.00335$) the Fieller set again covers in 0.9517, since its event involves only the errors, and is bounded in 0.817 of the experiments, close to the power 0.822 of the slope t test; the delta-method interval covers in 0.9827, wastefully. With the slope two standard errors from zero and $x_0 = 12$, beyond the standards, the Fieller set covers in 0.9528 and the delta-method interval in only 0.9337.

```
import numpy as np
from scipy import stats

rng = np.random.default_rng(1206)
x = np.repeat(np.arange(9.0), 3)           # standards, in triplicate
n = len(x)
y = 0.04 + 0.115 * x + 0.015 * rng.normal(size=n) # absorbance readings
m = 3
y0 = 0.04 + 0.115 * 5.3 + 0.015 * rng.normal(size=m) # the unknown sample, true x0
    = 5.3

def calibrate(x, y, y0bar, m, level=0.95):
    """Fieller set {x0 : |y0bar - b0 - b1 x0| <= t s sqrt(1/m + 1/n + (x0 - xbar)^2
    / Sxx)}. """
    n = len(x)
    xbar, Sxx = x.mean(), np.sum((x - x.mean()) ** 2)
    b1 = np.sum((x - xbar) * (y - y.mean())) / Sxx
    s2 = np.sum((y - y.mean() - b1 * (x - xbar)) ** 2) / (n - 2)
```




Figure 12.5.1: Calibration inverts the 95% prediction band for the mean of $m = 3$ readings (shaded); bars mark the calibration sets for the observed readings (horizontal lines). (a) A precise assay, magnified in the inset. (b) A weak assay: whole line (lower bar) and two rays (upper bar).

```

t2 = stats.t.ppf(0.5 + level / 2, n - 2) ** 2
g = y0bar - y.mean()
# quadratic in d = x0 - xbar: A d^2 - 2 B d + K <= 0
A = b1**2 - t2 * s2 / Sxx
B = b1 * g
K = g**2 - t2 * s2 * (1 / m + 1 / n)
disc = B**2 - A * K
if disc < 0:
    return "whole line", (-np.inf, np.inf), xbar + g / b1
r1, r2 = sorted([(B - np.sqrt(disc)) / A, (B + np.sqrt(disc)) / A])
if A > 0:
    return "interval", (xbar + r1, xbar + r2), xbar + g / b1
return "two rays", (xbar + r1, xbar + r2), xbar + g / b1 # (-inf, first] and [
second, inf)

```

```

shape, (lo, hi), x0_hat = calibrate(x, y, y0.mean(), m)
print(f"estimate {x0_hat:.3f}; 95% calibration set: {shape} ({lo:.3f}, {hi:.3f})")

```

[Run this code online](#)

Remark (Classical or inverse estimation?). The *inverse* estimator regresses the x_i on the readings and predicts from $\bar{Y}_0$: $\tilde{x}_0 = \bar{x} + (S_{xy}/S_{yy})(\bar{Y}_0 - \bar{Y})$, with $S_{xy} = \sum(x_i - \bar{x})(Y_i - \bar{Y})$ and $S_{yy} = \sum(Y_i - \bar{Y})^2$. Since $\bar{Y}_0 - \bar{Y} = \hat{\beta}_1(\hat{x}_0 - \bar{x})$ and $\hat{\beta}_1 = S_{xy}/S_{xx}$,

$$\tilde{x}_0 - \bar{x} = \frac{S_{xy}^2}{S_{xx}S_{yy}}(\hat{x}_0 - \bar{x}) = R^2(\hat{x}_0 - \bar{x}),$$

where R is the sample correlation of the calibration pairs. So $\tilde{x}_0$ shrinks $\hat{x}_0$ toward $\bar{x}$ by the factor $R^2 \in [0, 1]$. It has finite moments and can beat $\hat{x}_0$ in mean squared error near the centre of the standards (Krutchkoff 1967); away from the centre it does worse ([Exercise \(B3\)](#)). Hoadley (1970) showed that it is a Bayes estimator under a particular prior for x_0 .

12.5.3 Exercises

12.5.3.1 A. Check your understanding

Exercise. A1

For a straight line, write down Fieller's $1 - \alpha$ confidence set for the point $\xi = -\beta_0/\beta_1$ where the line crosses zero, and state the condition under which it is a bounded interval. Show that it is the calibration set of [Theorem 12.5.3](#) with $\bar{Y}_0$ replaced by 0 and $1/m$ replaced by 0.

Solution. With $\lambda_1 = (1, 0)^\top$ and $\lambda_2 = (0, 1)^\top$, $\xi = -a/b$. Apply [Proposition 12.5.1](#) to $\theta = a/b$ and change sign, or directly: ξ_0 is in the set iff the t interval for $\beta_0 + \beta_1\xi_0$ contains 0, that is iff $(\hat{\beta}_0 + \hat{\beta}_1\xi_0)^2 \leq t^2 s^2 \{1/n + (\xi_0 - \bar{x})^2/S_{xx}\}$. This is [Eq. 12.5.2](#) with $\bar{Y}_0 = 0$ and the $1/m$ term absent, because 0 is a known number rather than an average of new readings. It is bounded iff $T_1^2 > t^2$.

Exercise. A2

Show that $\hat{x}_0$, together with the least squares estimates from the standards, maximizes the joint likelihood of the $n+m$ readings over $(\beta_0, \beta_1, \sigma^2, x_0)$ when $\hat{\beta}_1 \neq 0$. What is the maximum likelihood estimate of σ^2 ?

Solution. For fixed σ^2 the likelihood is maximized by minimizing $\sum_i (Y_i - \beta_0 - \beta_1 x_i)^2 + \sum_j (Y_{0j} - \beta_0 - \beta_1 x_0)^2$. For any $\beta_1 \neq 0$ the second sum is minimized over x_0 by choosing $\beta_0 + \beta_1 x_0 = \bar{Y}_0$, leaving $\sum_j (Y_{0j} - \bar{Y}_0)^2$, which is free of the parameters. The first sum is then minimized by least squares on the standards, and $x_0 = (\bar{Y}_0 - \hat{\beta}_0)/\hat{\beta}_1 = \hat{x}_0$. The maximum likelihood estimate of σ^2 is $\{\text{SSE} + \sum_j (Y_{0j} - \bar{Y}_0)^2\}/(n+m)$.

12.5.3.2 B. Practice

Exercise. B1

When $m \geq 2$, the new readings carry their own information about σ^2 . Let $s_p^2 = \{\text{SSE} + \sum_j (Y_{0j} - \bar{Y}_0)^2\}/(n+m-3)$. Show that replacing s by s_p and $t_{n-2, \alpha/2}$ by $t_{n+m-3, \alpha/2}$ in [Eq. 12.5.2](#) gives another exact $1 - \alpha$ set. Why is it shorter on average, and why is it more often bounded?

Solution. The within-specimen sum of squares is $\sigma^2 \chi^2(m-1)$, independent of $\bar{Y}_0$ (the sample mean and sample variance of a normal sample are independent by Corollary 4.4.5) and of the standards. So $\{\text{SSE} + \sum_j (Y_{0j} - \bar{Y}_0)^2\}/\sigma^2 \sim \chi^2(n+m-3)$, independent of the numerator of the pivot, and the pivot with s_p is $t(n+m-3)$. More degrees of freedom give a smaller multiplier and a less variable estimate of σ , so the set is shorter on average. The bounded case requires $\hat{\beta}_1^2 S_{xx} > t^2 s_p^2$, and a smaller t makes this easier.

Exercise. B2

In the quadratic regression $EY_i = \beta_0 + \beta_1 x_i + \beta_2 x_i^2$, use [Proposition 12.5.1](#) to give a confidence set for the location $x^* = -\beta_1/(2\beta_2)$ of the turning point. When is it a bounded interval, and what does that condition mean for the curvature?

Solution. Take $\lambda_1 = (0, 1, 0)^\top$, $\lambda_2 = (0, 0, 2)^\top$ and $\theta = \beta_1/(2\beta_2)$, so $x^* = -\theta$. The set is $\{x_0 : (\hat{\beta}_1 + 2\hat{\beta}_2 x_0)^2 \leq t^2 s^2 (c_{11} + 4c_{12}x_0 + 4c_{22}x_0^2)\}$, where c_{ij} are the entries of $(\mathbf{X}^\top \mathbf{X})^{-1}$ belonging to β_1, β_2 . It is the set of x_0 at which the estimated derivative $\beta_1 + 2\beta_2 x_0$ is not significantly different from zero. It is bounded iff $\hat{\beta}_2^2 > t^2 s^2 c_{22}$, that is iff the curvature is significant: without significant curvature, the fitted curve may be nearly straight and its turning point anywhere.

Exercise. B3

Study the two estimators of the Remark “Classical or inverse estimation?” in an idealized large calibration experiment: the standards are so numerous that $\hat{\beta}_0$, $\hat{\beta}_1$ and R^2 may be replaced by constants β_0 , $\beta_1 \neq 0$ and $\rho^2 \in (0, 1)$, while $\bar{Y}_0$ is still the mean of m new readings. Let $v = \sigma^2/(m\beta_1^2)$.

1. Show that $\hat{x}_0$ is unbiased with mean squared error v , and that $\tilde{x}_0 = \bar{x} + \rho^2(\hat{x}_0 - \bar{x})$ has mean squared error $\rho^4 v + (1 - \rho^2)^2(x_0 - \bar{x})^2$.
2. Show that $\tilde{x}_0$ has the smaller mean squared error iff $(x_0 - \bar{x})^2 < v(1 + \rho^2)/(1 - \rho^2)$.
3. Let $\tau^2 = S_{xx}/n$, so that in this limit $\rho^2 = \beta_1^2 \tau^2 / (\beta_1^2 \tau^2 + \sigma^2)$. Show that the condition of part 2 reads $(x_0 - \bar{x})^2 < v + 2\tau^2/m$. With $m = 1$ and a precise instrument, roughly how far from $\bar{x}$, in units of the spread τ of the standards, does the inverse estimator stay ahead?

Solution.

1. Here $\hat{x}_0 = x_0 + \bar{\varepsilon}_0/\beta_1$ with $\bar{\varepsilon}_0 \sim N(0, \sigma^2/m)$, so it is unbiased with variance v . Then $\tilde{x}_0 - x_0 = \rho^2(\hat{x}_0 - x_0) - (1 - \rho^2)(x_0 - \bar{x})$: variance $\rho^4 v$ plus squared bias $(1 - \rho^2)^2(x_0 - \bar{x})^2$.
2. The inequality $\rho^4 v + (1 - \rho^2)^2(x_0 - \bar{x})^2 < v$ is $(1 - \rho^2)^2(x_0 - \bar{x})^2 < (1 - \rho^2)(1 + \rho^2)v$; divide by $(1 - \rho^2)^2 > 0$.
3. $(1 + \rho^2)/(1 - \rho^2) = (2\beta_1^2 \tau^2 + \sigma^2)/\sigma^2$, so $v(1 + \rho^2)/(1 - \rho^2) = v + 2\tau^2/m$. When σ is small, $v \approx 0$ and the inverse estimator wins only for $|x_0 - \bar{x}| < \sqrt{2}\tau$, a little more than one spread from the centre; beyond that its bias dominates.

12.5.3.3 C. Going deeper**Exercise. C1**

In the setting of [Proposition 12.5.1](#), let $E_t = \{\boldsymbol{\phi} \in \mathbb{R}^2 : ((\hat{a}, \hat{b})^\top - \boldsymbol{\phi})^\top \mathbf{V}^{-1}((\hat{a}, \hat{b})^\top - \boldsymbol{\phi}) \leq t^2 s^2\}$, an ellipse with the t constant in place of $2F_\alpha(2, n - r)$. Show that $\theta_0 \in \mathcal{F}(\mathbf{Y})$ iff the line $\{(\phi_1, \phi_2) : \phi_1 = \theta_0 \phi_2\}$ meets E_t . Use the picture to explain the three cases: the set is the whole line iff E_t contains the origin, and it is bounded iff E_t does not meet the axis $\phi_2 = 0$.

Solution. The line is $\{\boldsymbol{\phi} : \mathbf{c}^\top \boldsymbol{\phi} = 0\}$ with $\mathbf{c} = (1, -\theta_0)^\top$. By Corollary 12.2.4 (with the constant $t^2 s^2$), the values of $\mathbf{c}^\top \boldsymbol{\phi}$ over E_t form the interval $\hat{a} - \theta_0 \hat{b} \pm ts\sqrt{\mathbf{c}^\top \mathbf{V} \mathbf{c}}$, and $\mathbf{c}^\top \mathbf{V} \mathbf{c} = v_{11} - 2\theta_0 v_{12} + \theta_0^2 v_{22}$. So the line meets E_t iff this interval contains 0, which is Eq. 12.5.1. Every line through the origin meets E_t iff the origin is in E_t , which is $Q \leq t^2 s^2$. Lines of large slope approach the axis $\phi_2 = 0$ (the ϕ_1 -axis in these coordinates); the set is unbounded iff lines of arbitrarily large slope meet E_t , iff E_t meets that axis, which by Corollary 12.2.4 with $\mathbf{c} = (0, 1)^\top$ is $\hat{b}^2 \leq t^2 s^2 v_{22}$.

Exercise. C2

Two straight lines $EY = \alpha_k + \beta_k x$, $k = 1, 2$, are fitted to two independent samples in one model with a common σ^2 , as in a two-phase regression. Use Proposition 12.5.1 to construct a confidence set for the x -coordinate $(\alpha_1 - \alpha_2)/(\beta_2 - \beta_1)$ of their intersection. When is it bounded? Interpret the unbounded cases in terms of parallel lines.