

18.3 Heterogeneous slopes

The covariance model of [Section 18.1](#) assumes a common slope, so each treatment comparison is a vertical shift between parallel lines, one number. If the slopes differ, the treatment effect depends on the covariate, and “the” adjusted difference no longer exists. This section tests parallelism and says what to report when it fails.

18.3.1 The separate lines model and the test of parallelism

Keep one factor with g levels and one covariate, and allow each level its own line:

$$E(y_i) = \mu_k + \beta_k x_i \quad \text{for unit } i \text{ in group } k.$$

This is the separate lines model Eq. 8.6.4 of [Section 8.6](#). The parallel lines model is the special case $\beta_1 = \dots = \beta_g$. For group k write

$$E_{xx,k} = \sum_{i \in k} (x_i - \bar{x}_k)^2, \quad E_{xy,k} = \sum_{i \in k} (x_i - \bar{x}_k)(y_i - \bar{y}_k),$$

so that the pooled within-group quantities of [Section 18.1](#) are $E_{xx} = \sum_k E_{xx,k}$ and $E_{xy} = \sum_k E_{xy,k}$.

Theorem 18.3.1 — The test of parallel slopes

Assume x varies within every group, so that every $E_{xx,k} > 0$, and $n > 2g$.

- (a) In the separate lines model, $\hat{\beta}_k = E_{xy,k}/E_{xx,k}$, these estimates are uncorrelated with variances $\sigma^2/E_{xx,k}$, and the residual sum of squares is

$$\text{SSE}_{\text{sep}} = \sum_k \left(E_{yy,k} - \frac{E_{xy,k}^2}{E_{xx,k}} \right)$$

on $n - 2g$ degrees of freedom.

- (b) The pooled slope of the parallel lines model is the weighted mean $\hat{\beta} = \sum_k E_{xx,k} \hat{\beta}_k / \sum_k E_{xx,k}$, and the extra sum of squares for the separate slopes is

$$\text{SS}_{\text{slopes}} = \sum_k E_{xx,k} (\hat{\beta}_k - \hat{\beta})^2 = \sum_k \frac{E_{xy,k}^2}{E_{xx,k}} - \frac{E_{xy}^2}{E_{xx}} \quad (18.3.1)$$

on $g - 1$ degrees of freedom.

- (c) If $\mathbf{Y}$ is normal,

$$F = \frac{\text{SS}_{\text{slopes}}/(g-1)}{\text{SSE}_{\text{sep}}/(n-2g)} \sim F(g-1, n-2g, \gamma),$$

with

$$\gamma = \frac{1}{\sigma^2} \sum_k E_{xx,k} (\beta_k - \bar{\beta})^2,$$

where $\bar{\beta} = \sum_k E_{xx,k} \beta_k / \sum_k E_{xx,k}$. The test that rejects for large F is the F test of parallelism.

Proof. Let $\mathbf{M}$ project onto the span of the group indicators, and let $\mathbf{w}_k$ be the vector that equals $x_i - \bar{x}_k$ for units of group k and zero elsewhere. The k th slope column of the separate lines model is $\mathbf{x}$ restricted to group k , and its residual after projection onto $\mathcal{C}(\mathbf{M})$ is $\mathbf{w}_k$. By Lemma 9.2.1 the separate lines model space is $\mathcal{C}(\mathbf{M}) \oplus \text{span}\{\mathbf{w}_1, \dots, \mathbf{w}_g\}$, and the parallel lines space is $\mathcal{C}(\mathbf{M}) \oplus \text{span}\{\mathbf{w}\}$ with $\mathbf{w} = \sum_k \mathbf{w}_k = (\mathbf{I} - \mathbf{M})\mathbf{x}$. The vectors $\mathbf{w}_k$ have disjoint supports, so they are mutually orthogonal, with $\|\mathbf{w}_k\|^2 = E_{xx,k}$ and $\mathbf{w}_k^\top \mathbf{y} = E_{xy,k}$.

- (a) The projection of $\mathbf{y}$ onto $\text{span}\{\mathbf{w}_k\}$ is $\sum_k (E_{xy,k}/E_{xx,k}) \mathbf{w}_k$. Reading off the coefficient of the k th slope column gives $\hat{\beta}_k$, which is $\mathbf{w}_k^\top \mathbf{y} / \|\mathbf{w}_k\|^2$. Hence $\text{Cov}(\hat{\beta}_k, \hat{\beta}_l) = \sigma^2 \mathbf{w}_k^\top \mathbf{w}_l / (E_{xx,k} E_{xx,l})$, which is $\sigma^2 / E_{xx,k}$ for $k = l$ and zero otherwise. The residual sum of squares is $\|(\mathbf{I} - \mathbf{M})\mathbf{y}\|^2 - \sum_k E_{xy,k}^2 / E_{xx,k}$, and $\|(\mathbf{I} - \mathbf{M})\mathbf{y}\|^2 = \sum_k E_{yy,k}$. The model has rank $2g$.
- (b) The pooled slope is $\mathbf{w}^\top \mathbf{y} / \|\mathbf{w}\|^2 = E_{xy} / E_{xx}$, and $E_{xy} = \sum_k E_{xx,k} \hat{\beta}_k$. By Theorem 6.5.1 the extra sum of squares is the squared length of the projection onto the orthogonal complement of $\mathbf{w}$ in $\text{span}\{\mathbf{w}_k\}$, which equals $\sum_k E_{xy,k}^2 / E_{xx,k} - E_{xy}^2 / E_{xx}$. Writing $E_{xy,k} = E_{xx,k} \hat{\beta}_k$ and expanding $\sum_k E_{xx,k} (\hat{\beta}_k - \bar{\beta})^2 = \sum_k E_{xx,k} \hat{\beta}_k^2 - \bar{\beta}^2 E_{xx}$ gives the first form.
- (c) By Theorem 11.1.1, with $\sigma^2 \gamma$ equal to the extra sum of squares evaluated at the mean vector. The mean vector has $\mathbf{w}_k^\top \boldsymbol{\mu} = \beta_k E_{xx,k}$, because $\mathbf{w}_k$ is orthogonal to the group indicators. Substituting into (b) with $\hat{\beta}_k$ replaced by β_k gives γ .

■

The test is a weighted test of homogeneity of g independent slope estimates, with weights equal to their precisions $E_{xx,k}/\sigma^2$.

Example. (Parallelism in the tutoring experiment)

For the four formats of Example (Four study formats with a pretest), the separate slopes are 0.703, 0.823, 1.035 and 0.992, and their weighted mean is the pooled slope 0.8670. The separate lines model has residual sum of squares 1347.47 on 32 degrees of freedom, against 1407.47 on 35 for parallel lines. The test gives $F = 0.47$ on 3 and 32 degrees of freedom, $p = 0.70$, and the covariance analysis stands.

```
import numpy as np
from scipy import stats

rng = np.random.default_rng(84)
g, m = 4, 10                                # four formats, ten students
    each
n = g * m
group = rng.permutation(np.repeat(np.arange(g), m)) # the randomization
x = np.round(rng.normal(60, 10, n))             # pretest, measured before
    assignment
mu_true = np.array([70.0, 74.0, 71.0, 77.0])
y = np.round(mu_true[group] + 0.8 * (x - 60) + rng.normal(0, 6, n), 1)
Z = np.eye(g)[group]                          # indicator matrix of the
```

```

    formats
n_k = Z.sum(axis=0)

def within(v):
    """(I - M) v: deviations from the group means (M projects onto C(Z))."""
    return v - (Z.T @ v / n_k)[group]

E_xx, E_xy, E_yy = within(x) @ within(x), within(x) @ within(y), within(y) @ within(
    y)
sse = E_yy - E_xy ** 2 / E_xx # ANCOVA residual sum of
    squares
X_sep = np.column_stack([Z, Z * x[:, None]]) # a separate line for each
    format
E_xx_k = np.array([within(x)[group == k] @ within(x)[group == k] for k in range(g)])
E_xy_k = np.array([within(x)[group == k] @ within(y)[group == k] for k in range(g)])
slopes = E_xy_k / E_xx_k
sse_sep = sse - (np.sum(E_xy_k ** 2 / E_xx_k) - E_xy ** 2 / E_xx)
F_par = ((sse - sse_sep) / (g - 1)) / (sse_sep / (n - 2 * g))
print("separate slopes", slopes.round(3),
      f" F = {F_par:.2f}, p = {stats.f.sf(F_par, g - 1, n - 2 * g):.3f}")

```

[Run this code online](#)

With ten students per group the test has little power, so the case for parallel lines rests mostly on the science. When a treatment is meant to help some units more than others, such as a remedial programme aimed at weak students, the separate lines model should be the starting point rather than a check.

18.3.2 When the slopes differ

If the slopes differ, the difference between treatments k and l at covariate value x is the line

$$\delta(x) = (\mu_k - \mu_l) + (\beta_k - \beta_l)x,$$

estimated by $\hat{\delta}(x) = (\hat{\mu}_k - \hat{\mu}_l) + (\hat{\beta}_k - \hat{\beta}_l)x$. The two lines are independent simple regressions, so by Corollary 5.5.4

$$\text{Var}(\hat{\delta}(x)) = \sigma^2 v(x), \quad v(x) = \frac{1}{n_k} + \frac{(x - \bar{x}_k)^2}{E_{xx,k}} + \frac{1}{n_l} + \frac{(x - \bar{x}_l)^2}{E_{xx,l}}. \quad (18.3.2)$$

Honest reports include:

- (i) Report $\hat{\delta}(x)$ with its standard error at a few covariate values chosen for their meaning, for instance the quartiles of x , or a clinical threshold. Each is a t interval (Theorem 12.1.1) with $n - 2g$ degrees of freedom.
- (ii) Draw $\hat{\delta}(x)$ as a function with a confidence band. The pointwise band uses $t_{n-2g, \alpha/2}$. A band that covers the whole line $\delta(\cdot)$ with probability exactly $1 - \alpha$ replaces this multiplier by $\sqrt{2F_\alpha(2, n - 2g)}$, because $\{\delta(x) : x \in \mathbb{R}\}$ lies in a two-dimensional space of estimable functions. This is the argument of Theorem 13.2.2, as used for the Working–Hotelling band (Theorem 12.4.1).
- (iii) Give the set of covariate values at which the treatments differ significantly. This is the **Johnson–Neyman region** (Proposition 18.3.2 below).

Reporting $\hat{\delta}(\bar{x})$ alone as *the* treatment effect hides the slope, which is the whole point.

Remark (The average effect in a randomized experiment). There is one exception. In a randomized experiment, δ at the population mean of the covariate is the *average* treatment effect, because δ is linear. Its estimate $\hat{\delta}(\bar{x})$ is the treatment coefficient in the regression with a treatment-by-covariate interaction and the covariate centred at $\bar{x}$ (Exercise (C1)). This “interacted” adjustment is asymptotically never less precise than the unadjusted difference, even when the model is wrong. Chapter 25 returns to it (Theorem 25.6.2).

18.3.3 The Johnson–Neyman region

Proposition 18.3.2 — Johnson–Neyman region

In the separate lines model with normal errors, let $s^2 = \text{SSE}_{\text{sep}}/(n - 2g)$ and fix a multiplier $c > 0$. Write $\Delta_\mu = \hat{\mu}_k - \hat{\mu}_l$ and $\Delta_\beta = \hat{\beta}_k - \hat{\beta}_l$. The set of x with $|\hat{\delta}(x)| > cs\sqrt{v(x)}$ is $\{x : Ax^2 + Bx + C > 0\}$, where

$$A = \Delta_\beta^2 - c^2 s^2 \left(\frac{1}{E_{xx,k}} + \frac{1}{E_{xx,l}} \right), \quad B = 2\Delta_\mu \Delta_\beta + 2c^2 s^2 \left(\frac{\bar{x}_k}{E_{xx,k}} + \frac{\bar{x}_l}{E_{xx,l}} \right),$$

$$C = \Delta_\mu^2 - c^2 s^2 \left(\frac{1}{n_k} + \frac{1}{n_l} + \frac{\bar{x}_k^2}{E_{xx,k}} + \frac{\bar{x}_l^2}{E_{xx,l}} \right).$$

If $A > 0$, which happens iff the t test of $\beta_k = \beta_l$ with multiplier c rejects, the region is the whole line with at most an interval removed: δ is significantly positive on one side and significantly negative on the other. If $A < 0$, the region is a bounded interval or empty. With $c = t_{n-2g, \alpha/2}$ each point of the region is significant at level α . With $c = \sqrt{2F_\alpha(2, n-2g)}$ the statement “ $\delta(x) \neq 0$ for every x in the region” holds with probability at least $1 - \alpha$.

Proof. Square both sides of $|\hat{\delta}(x)| > cs\sqrt{v(x)}$ and expand $\hat{\delta}(x)^2 = \Delta_\mu^2 + 2\Delta_\mu \Delta_\beta x + \Delta_\beta^2 x^2$ and $v(x)$ from Eq. 18.3.2 as polynomials in x . Collecting terms gives $Ax^2 + Bx + C$. The coefficient A is positive iff $\Delta_\beta^2 / \{s^2(1/E_{xx,k} + 1/E_{xx,l})\} > c^2$, which is the squared t statistic for $\beta_k - \beta_l$ exceeding c^2 . A quadratic with $A > 0$ is positive outside the interval between its real roots (everywhere if it has none), and with $A < 0$ it is positive between its real roots (nowhere if it has none). The last statement follows from the simultaneous band of item (ii) above: on the event that the band covers $\delta(\cdot)$ everywhere, $\delta(x)$ cannot be zero at any point where the band excludes zero. ■

Example. (Two versions of a reading programme)

In a second synthetic randomized experiment, 48 children with baseline reading fluency between 42 and 118 words per minute are assigned, 24 each, to the current and a new version of a reading programme. The response is the gain in fluency. The new version was designed for weak readers, and the fitted lines of Figure 18.3.1(a) show why that matters. The slope is 0.0156 for the current version and -0.1615 for the new one. The




Figure 18.3.1: The reading programme. (a) Gain against baseline fluency with the separate least squares lines. (b) The estimated difference $\hat{\delta}(x)$ (new minus current) with a pointwise 95% band (dashed) and a simultaneous 95% band (shaded). The dotted line marks the lower end of the Johnson–Neyman region, $x = 93.5$; to its left the difference is significantly positive.

test of parallelism has $F = 11.2$ on 1 and 44 degrees of freedom, $p = 0.0017$. The pooled slope of the parallel lines model would be -0.0638 , a compromise that describes neither group.

The estimated lines cross at $x = 107.7$. At a baseline of 60 words per minute the new version adds 8.46 words per minute (standard error 1.56). At 100 it adds 1.37 (standard error 1.46), which is not significant. With $c = t_{44,0.025}$, the Johnson–Neyman region (Proposition 18.3.2) is $x < 93.5$ together with $x > 152.0$. The second part lies far beyond the largest baseline in the data, and it is an artefact of extending two fitted lines. The useful conclusion is that the new version is better for children who start below about 93 words per minute, which covers 33 of the 48 children in the study. With the simultaneous multiplier $\sqrt{2F_{0.05}(2, 44)}$, the region shrinks slightly, to $x < 90.8$ (and $x > 195.6$).

```
import numpy as np
from scipy import stats

rng = np.random.default_rng(31)
n_per = 24
arm = rng.permutation(np.repeat([0, 1], n_per))    # 0 = current version, 1 = new
version
x = np.round(rng.uniform(40, 120, 2 * n_per))      # baseline fluency
y = np.where(arm == 0, 14 + 0.06 * x, 36 - 0.16 * x) + rng.normal(0, 4, 2 * n_per)
y = np.round(y, 1)
n = len(y)

def line(k):
    """Least squares line within arm k, and the arm's summaries."""
    xs, ys = x[arm == k], y[arm == k]
    Sxx = np.sum((xs - xs.mean()) ** 2)
    b = np.sum((xs - xs.mean()) * (ys - ys.mean())) / Sxx
    a = ys.mean() - b * xs.mean()
    sse = np.sum((ys - a - b * xs) ** 2)
```

```

    return a, b, xs.mean(), Sxx, len(xs), sse

(a0, b0, xb0, S0, n0, e0), (a1, b1, xb1, S1, n1, e1) = line(0), line(1)
nu = n - 4
s2 = (e0 + e1) / nu                                # separate-lines model

def significant_region(crit2):
    """Roots of  $d(x)^2 - \text{crit2} * s2 * v(x) = 0$ , a quadratic in  $x$ ."""
    A = (b1 - b0) ** 2 - crit2 * s2 * (1 / S0 + 1 / S1)
    B = 2 * (a1 - a0) * (b1 - b0) + 2 * crit2 * s2 * (xb0 / S0 + xb1 / S1)
    C = (a1 - a0) ** 2 - crit2 * s2 * (1 / n0 + 1 / n1 + xb0 ** 2 / S0 + xb1 ** 2 / S1)
    return np.sort(np.roots([A, B, C]).real)

t2 = stats.t.ppf(0.975, nu) ** 2                    # pointwise, 5% level
w2 = 2 * stats.f.ppf(0.95, 2, nu)                  # simultaneous over all  $x$  (
    Scheffe,  $q = 2$ )
x_cross = -(a1 - a0) / (b1 - b0)
print(f"lines cross at  $x = \{x\_cross:.1f\}$ ")
print("pointwise 5% region: difference significant outside", significant_region(t2).
    round(1))
print("simultaneous region: difference significant outside", significant_region(w2).
    round(1))

```

[Run this code online](#)

The region is only as good as the linearity of each line, it extrapolates outside the data, and it marks significance, not practical importance: with enough data every value except the crossing point becomes significant.

18.3.4 Several covariates, several factors

With q covariates, the test of a common γ against $\gamma_1, \dots, \gamma_g$ has $(g-1)q$ and $n - g(q+1)$ degrees of freedom. With two factors, slope terms can be built term by term like the means model of [Chapter 16](#). Each such test compares nested column spaces (Theorem 11.1.1).

18.3.5 Exercises

18.3.5.1 A. Check your understanding

Exercise A1.

For $g = 2$, show that [Eq. 18.3.1](#) reduces to $SS_{\text{slopes}} = \frac{E_{xx,1}E_{xx,2}}{E_{xx,1} + E_{xx,2}} (\hat{\beta}_1 - \hat{\beta}_2)^2$, and that the F statistic is the square of the two-sample t statistic for the difference of slopes.

Exercise A2.

In [Example \(Two versions of a reading programme\)](#), the estimated lines cross at $x =$

107.7. Explain why the Johnson–Neyman region never contains the crossing point, whatever the multiplier.

18.3.5.2 B. Practice

Exercise B1.

Give a numerical example of two groups for which $A < 0$ and the Johnson–Neyman region of Proposition 18.3.2 is a bounded interval. Interpret the region.

Exercise B2.

In the tutoring experiment, suppose the true slopes were 0.8, 0.8, 0.8, 1.2 and $\sigma = 6$. Using the $E_{xx,k}$ of the example (their sum is 3643.9; take them equal for the calculation), compute the noncentrality of the test of parallelism and its power at level 0.05 (Theorem 11.5.1).

Exercise B3.

Prove the claim in item (ii): with $c = \sqrt{2F_\alpha(2, n-2g)}$, the band $\hat{\delta}(x) \pm cs\sqrt{v(x)}$ contains $\delta(x)$ for all x simultaneously with probability at least $1 - \alpha$. Is the probability exactly $1 - \alpha$?

Solution. The functions $\delta(x) = \lambda_x^\top \beta$ for $x \in \mathbb{R}$ have λ_x in a two-dimensional space $\mathcal{L}$ spanned by the coefficient vectors of $\mu_k - \mu_l$ and $\beta_k - \beta_l$, both estimable. By Theorem 13.2.2(b) with $q = 2$, with probability exactly $1 - \alpha$ every $\lambda \in \mathcal{L}$ satisfies $|\lambda^\top \hat{\beta} - \lambda^\top \beta| \leq c \text{se}(\lambda^\top \hat{\beta})$. The functions λ_x form a subset of $\mathcal{L}$, so the probability for them is at least $1 - \alpha$. It is in fact exactly $1 - \alpha$. In the coordinates $(\mu_k - \mu_l, \beta_k - \beta_l)$ we have $\lambda_x = (1, x)$, and the studentized ratio $(\lambda_x^\top \hat{\beta} - \lambda_x^\top \beta) / \text{se}(\lambda_x^\top \hat{\beta})$ is unchanged when λ is rescaled and is continuous on $\mathcal{L} \setminus \{0\}$. Every direction of $\mathcal{L}$ except that of $(0, 1)$ is the direction of some λ_x , and $\lambda_x / \|\lambda_x\| \rightarrow \pm(0, 1)$ as $x \rightarrow \pm\infty$. So the supremum of the ratio over $x \in \mathbb{R}$ equals its supremum over $\mathcal{L} \setminus \{0\}$, and the two events coincide. This is the argument of Theorem 12.4.1(b) for a single line.

18.3.5.3 C. Going deeper

Exercise C1.

In a two-group randomized experiment, show that the treatment coefficient in the least squares fit of y on 1, T , $x - \bar{x}$ and $T(x - \bar{x})$, where $\bar{x}$ is the overall mean, equals $(\bar{y}_1 - \hat{\beta}_1(\bar{x}_1 - \bar{x})) - (\bar{y}_0 - \hat{\beta}_0(\bar{x}_0 - \bar{x}))$, with $\hat{\beta}_k$ the within-group slopes. Compare with the adjusted difference of the parallel lines model.

Solution. The model with an interaction is the separate lines model, so its fitted lines are the two within-group least squares lines, $\bar{y}_k + \hat{\beta}_k(x - \bar{x}_k)$. In the given parameterization the line for group k is $(\beta_0 + \beta_T T) + (\beta_x + \beta_{Tx} T)(x - \bar{x})$, so the coefficient of T is the difference of the two lines at $x = \bar{x}$, which is the stated expression. The parallel lines model replaces both $\hat{\beta}_k$ by the pooled slope. The two estimates agree when the slopes

are equal or when $\bar{x}_0 = \bar{x}_1 = \bar{x}$.