# Introduction to Numerical Methods

#### <u>In this lecture...</u>



• The justification for pricing by Monte Carlo simulation



• Grids and discretization of derivatives

• The explicit finite-difference method

By the end of this lecture you will be able to

- implement the Monte Carlo method for simulating asset paths and pricing options
- implement the explicit finite-difference method for pricing options

#### **Introduction**

More often than not we must solve option-pricing problems by numerical means.

It is rare to be able to find closed-form solutions for prices unless both the contract and the model are very simple.

The most useful numerical techniques are Monte Carlo simulations and finite-difference methods.

grid based

Computational Finance - Numerical methods applied to problem in finance

#### **Monte Carlo Simulations**

#### Relationship between derivative values and simulations

Theory says:

pected payoff at expiry under a *risk-neutral* random walk for the underlying.

The risk-neutral random walk for S is

$$dS = mSdt + \sigma SdX$$

$$dS = rSdt + \sigma SdX.$$

This is simply our usual lognormal random walk but with the risk-free rate instead of the real growth rate.

#### **Justification:**

Binomial method

Rist • Black-Scholes Equation similar to backward Kolmogorov equation

SeS • Martingale theory

We can therefore write



option value  $=e^{-r(T-t)}E$  [payoff(S)]

provided that the expectation is with respect to the risk-neutral random walk, not the *real* one.

#### Monte Carlo recipe

#### The algorithm:

underlying state variable

- $\checkmark$ 1. Simulate the risk-neutral random walk starting at today's value of the asset  $S_0$  over the required time horizon. This gives one realization of the underlying price path.
- $\sqrt{2}$ . For this realization calculate the option payoff.
- ✓3. Perform many more such realizations over the time horizon.
- 4. Calculate the average payoff over all realizations.
- 5. Take the present value of this average, this is the option value.

Average: 
$$\frac{1}{n}$$
  $\sum_{i=1}^{n}$   $\max_{i=1}^{n}$   $\sum_{i=1}^{n}$   $\sum_{i=1}^{$ 

#### How do we simulate the asset?

 $dS = (r-D)Sdt + \sigma SdX,$   $dr = (r-V)dt + \sigma dX,$ 

Two ways:



- 1. **If** the s.d.e. for the asset path is integrable **and** the contract is not path dependent (or American) **then** simulate in 'one giant leap'
- 2. **Otherwise** you will have to simulate time step by time step, the entire path

#### One giant leap: A method that works in special cases

For the lognormal random walk we are lucky that we can find a simple, and exact, time stepping algorithm.

We can write the risk-neutral stochastic differential equation for  ${\cal S}$  in the form

$$d(\log S) = \left(r - \frac{1}{2}\sigma^2\right)dt + \sigma dX.$$

This can be integrated exactly to give

$$S(t) = S(0) \exp\left(\left(r - \frac{1}{2}\sigma^2\right)t + \sigma\int_0^t dX\right).$$
 i.e. 
$$\sum_{n=1}^{\infty} \frac{1}{2}\sigma^2 \int_0^t dx = \frac{1}{2}\sigma^2 \int_0^t$$

$$\phi \sim \eta(=,1)$$

Because this expression is exact and simple it is the best time stepping algorithm to use... but only if we have a payoff that only depends on the final asset value, i.e. is European and path independent.

We can then simulate the final asset price in one giant leap, using a time step of T if both of these are true

• the s.d.e. is integrable and

• the contract is European and not path dependent

Recep from mod I

#### Simulating the entire path: A method that always works

Price paths are simulated using a discrete version of the stochastic differential equation for S.

An obvious choice is to use

$$S_{T} = S_{C} + \sigma S \sqrt{\delta t} \phi,$$

$$S_{C+} = S_{C} \left( 1 + \ell d + \delta \phi \right) d + \delta d +$$

where  $\phi$  is from a standardized Normal distribution.

• This way of simulating the time series is called the **Euler**— **method**. This method has an error of  $O(\delta t)$ .

#### Errors $\times$ 2

There are two (at least) sources of error in the Monte Carlo method:

- If the size of the time step is  $\delta t$  then we may introduce errors of  $O(\delta t)$  by virtue of the discrete approximation to continuous events +  $\dot{m}e$   $\rightarrow$   $te\rho$
- Because we are only simulating a finite number of an infinite number of possible paths, the error due to using N realizations of the asset price paths is  $O(N^{-1/2})$ .

### Generating Normal variables — Recap

fast but inaccrete

• **Quick 'n' dirty**: A useful distribution that is easy to implement on a spreadsheet, and is fast, is the following *approximation* to the Normal distribution:

where the  $\psi_i$  are independent random variables, drawn from a uniform distribution over zero to one.

There are other methods such as **Box–Muller**, more later.

Polar Marsaglia

acceptance rejection

#### Other issues

· Greeks; method has to se repeated

Early exercise (and other decisions)

Los free boundars problems


Vega

N(2+92)-N(2)

(F) = V(+6+ )-1(S,1)

St

#### **Advantages of Monte Carlo simulations**





- Correlations can be easily modeled, and it is easy to price options on many assets (high-dimensional contracts)
- It is computationally quite efficient in high dimensions
- There is plenty of software available, at the very least there are spreadsheet functions that will suffice for most of the time
  - To get a better accuracy, just run more simulations

• The effort in getting some answer is very low

The models can often be changed without much work <</p>

• Complex path dependency can often be easily incorporated

Many contracts can be priced at the same time

People accept the technique, and will believe your answers

Lew scope for error.

## Disadvantages of Monte Carlo simulations

christian chooses

- The method is very slow, you need a lot of simulations to r= r(t) ( r(t)-1-0get an accurate answer
- Finding the greeks can be hard have consuming.
- The method does not cope well with early exercise, de

Numerical Integration technique Simpson's rule
Trapezoidal

Milstei- Schene

#### **Finite Difference Methods**

Runge Kutta


Monte Carlo simulations can be very slow to converge to the answer, and they do not give us the greeks without further effort.

There is a method that is very similar to the binomial tree method which is the method of choice for certain types of problem.

Grids FDM came to finance from C. F. D.

Recall the shape of the binomial tree. . .

I Create a grid over which we II to discretize the replace each term in the B.J. E with approx s PDE - difference eg (directe antivous time) The shape of the tree is determined by the asset volatility.



The finite-difference grid.

Discretising time stock.

Differentiation using the grid:  $\frac{\partial V}{\partial \xi}$ ,  $\frac{\partial V}{\partial S}$ ,  $\frac{\partial^2 V}{\partial S^2}$   $\sim 900 \times 10^{-5}$ 

**Notation:** time step  $\delta t$  and asset step  $\delta S$ . The grid is made up of the points at asset values

where  $0 \le i \le I$  and  $0 \le k \le K$ .

We will be solving for the asset value going from zero up to the asset value  $I \delta S$ .


The Black–Scholes equation is to be solved for  $0 \le S < \infty$  so that  $I \delta S$  is our approximation to infinity.

Write the option value at each of these grid points as  $\frac{2}{2} = -\frac{3}{2}$ 

Write the option value at each of these grid points as 
$$\frac{1}{2} = \frac{1}{2}$$

$$\sqrt{\left(\int_{-\infty}^{\infty} T_{-} + V_{i}^{k} - V_{i}^{k} -$$

The superscript is the time variable and the subscript the



Let us now approx. The derivative Approximating  $\theta$  terms in the B.S.  $\epsilon$ .


The definition of the first time derivative of V is simply

backward diff. 
$$\frac{\partial V}{\partial t} = \lim_{h \to 0} \frac{V(S, t) - V(S, t - h)}{h}.$$

It follows naturally that we can approximate the time derivative from our grid of values using

$$\frac{\partial V}{\partial t}(S,t) \approx \frac{V_i^k - V_i^{k+1}}{\delta t}.$$
 of minus coming from T-kSt

This is our approximation to the option's theta.



Approximating the theta.

How accurate is this approximation?

We can expand the option value at asset value S and time  $t-\delta t$ in a Taylor series about the point S, t as follows.

Type 
$$V(S, t - \delta t) = V(S, t) - \delta t \frac{\partial V}{\partial t}(S, t) + O(\delta t^2).$$


In terms of values at grid points this is just

$$V_i^k = V_i^{k+1} + \delta t \frac{\partial V}{\partial t}(S, t) + O(\delta t^2). \blacktriangleleft$$


Which, upon rearranging, is 
$$\sqrt{(\int_{V(s,t)}^{t})} = \sqrt{(\int_{V(s,t)}^{t})} + O(\delta t).$$
• The error is  $O(\delta t)$ .

Approximating 
$$\triangle = \frac{\partial V}{\partial J}$$

Examine a cross section of the grid at one of the time steps.



These three approximations are



These are called a **forward difference**, a **backward difference** and a **central difference** respectively.

One of these approximations is better than the others. (Let one)

From a Taylor series expansion of the option value about the point  $S + \delta S$ , t we have

$$V(S + \delta S, t) = V(S, t) + \delta S \frac{\partial V}{\partial S}(S, t) + \frac{1}{2} \delta S^2 \frac{\partial^2 V}{\partial S^2}(S, t) + O(\delta S^3) + O(\delta S^4)$$

Similarly,

$$V(S - \delta S, t) = V(S, t) - \delta S \frac{\partial V}{\partial S}(S, t) + \frac{1}{2} \delta S^2 \frac{\partial^2 V}{\partial S^2}(S, t) + O(\delta S^2) + O(\delta S^2)$$

$$0 - 2 : V(S + \Delta I) - V(I - \delta I) = 2 \int_{\Delta I} \frac{\partial V}{\partial S} + O(\delta I^2) + O(\delta I^2) + O(\delta I^2)$$

$$0 + 2 : V(S + \Delta I) + V(I - \delta I) = 2 V(I) + \frac{1}{2} \frac{\partial^2 V}{\partial S^2} + O(\delta I^2) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S, t) - \delta S \frac{\partial V}{\partial S}(S, t) + \frac{1}{2} \frac{\partial^2 V}{\partial S^2}(S, t) + O(\delta I^2)$$

$$0 + 2 : V(S + \Delta I) + V(I - \delta I) = 2 V(I) + \frac{1}{2} \frac{\partial^2 V}{\partial S^2} + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S + \Delta I) - V(I - \delta I) + O(\delta I^2)$$

$$0 + 2 : V(S + \Delta I) + V(I - \delta I) = 2 \int_{\Delta I} V(I + \delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S + \Delta I) - V(I - \delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S + \Delta I) - V(I - \delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S + \Delta I) - V(I - \delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S + \Delta I) - V(I - \delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S + \Delta I) - V(I - \delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S + \Delta I) - V(I - \delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S + \Delta I) - V(I - \delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S + \Delta I) - V(I - \delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S + \Delta I) - V(I - \delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S + \Delta I) - V(I - \delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S + \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S + \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S + \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S + \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S + \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S - \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S - \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S - \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S - \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S - \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S - \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S - \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S - \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S - \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S - \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S - \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S - \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S - \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S - \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S - \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S - \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S - \Delta I) + O(\delta I^2)$$

$$V(S - \delta S, t) = V(S - \Delta I) + O(\delta I^2)$$


$$V(S - \delta$$

From these we get

$$\frac{\partial V}{\partial S}(S,t) = \frac{V_{i+1}^k - V_{i-1}^k}{2\delta S} + O(\delta S^2).$$


• The central difference has an error of  $O(\delta S^2)$ , the error in the forward and backward differences are both much larger,  $O(\delta S)$ .

The central difference calculated at S requires knowledge of the option value at  $S + \delta S$  and  $S - \delta S$ .



#### **Approximating** □

Gamma is the sensitivity of the delta to the underlying.



Calculate the delta half way between i and i+1, and the delta half way between i-1 and i . . . and difference them!

Forward difference 
$$=\frac{V_{i+1}^k - V_i^k}{\delta S}$$
.


Backward difference 
$$=\frac{V_i^k - V_{i-1}^k}{\delta S}$$
.

Therefore the natural approximation for the gamma is

$$\frac{\partial^2 V}{\partial S^2}(S,t) \approx \frac{\frac{V_{i+1}^k - V_i^k}{\delta S} - \frac{V_i^k - V_{i-1}^k}{\delta S}}{\delta S}$$

$$= \frac{V_{i+1}^{k} - 2V_{i}^{k} + V_{i-1}^{k}}{\delta S^{2}} + \mathcal{O}(\delta S^{2})$$

The error in this approximation is also  $O(\delta S^2)$ .



#### Final conditions and payoffs

We know that at expiry the option value is just the payoff function. At expiry we have

$$V(S,T) = \mathsf{Payoff}(S)$$
 at  $t = \int$ 

or, in our finite-difference notation,

$$V_i^0 = \text{Payoff}(i \, \delta S).$$

$$U_i^0 = \text{Payoff}(i \, \delta S).$$

The right-hand side is a known function.

For example, if we are pricing a call option we have

$$V_i^0 = \underbrace{\max(i\,\delta S - E, 0)}_{S}.$$

This final condition will get our finite-difference scheme started.

$$\frac{3V}{3V} \approx \frac{V_{i} - V_{i}}{5V_{i}}; \frac{3V}{3V} \approx \frac{V_{i+1} - V_{i+1}}{285}$$

$$\frac{3V}{3V} \approx \frac{V_{i-1} - 2V_{i} + V_{i+1}}{285}$$

$$\frac{3V_{i}}{3V_{i}} \approx \frac{V_{i-1} - 2V_{i} + V_{i+1}}{3V_{i}}; \frac{3V_{i} + V_{i}}{3V_{i}}; \frac{3V_{i}}{3V_{i}}; \frac{3V_{i$$

The explicit finite-difference method

Singlest of all


The Black-Scholes equation is

The Black-Scholes equation is 
$$\frac{\partial V}{\partial t} + \frac{1}{2}\sigma^2 S^2 \frac{\partial^2 V}{\partial S^2} + rS \frac{\partial V}{\partial S} - rV = 0. \qquad \text{support}$$
 Write this as 
$$\frac{\partial V}{\partial t} + a(S,t) \frac{\partial^2 V}{\partial S^2} + b(S,t) \frac{\partial V}{\partial S} + c(S,t) V = 0$$

so that we can examine more general problems.

Using the above approximations

$$\frac{V_{i}^{k} - V_{i}^{k+1}}{\delta t} + o_{i}^{k} \left( \frac{V_{i+1}^{k} - 2V_{i}^{k} + V_{i-1}^{k}}{\delta S^{2}} \right) + b_{i}^{k} \left( \frac{V_{i+1}^{k} - V_{i-1}^{k}}{2\delta S} \right) \\ + e_{i}^{k} V_{i}^{k} = O(\delta t, \delta S^{2}).$$



k.

(That's why this is called the **explicit finite-difference method**.)

$$V_{i-1}^{k+1} = V_{i}^{k} + \frac{1}{2}\sigma_{i}^{2} \int_{a}^{b} \left(V_{i-1}^{k} - 2V_{i+1}^{k} + V_{i+1}^{k}\right) + \frac{(r-D)i}{2}i\delta_{i}\left(V_{i-1}^{k} - V_{i+1}^{k}\right) - rV_{i}^{k}\delta_{i}$$


$$V_{i-1}^{k} : \frac{1}{2}\sigma_{i}^{2} \int_{a}^{b} \left(V_{i-1}^{k} - V_{i+1}^{k}\right) + \frac{(r-D)i}{2}i\delta_{i}\left(V_{i-1}^{k} - V_{i+1}^{k}\right) - rV_{i}^{k}\delta_{i}$$

$$V_{i-1}^{k} : \frac{1}{2}\sigma_{i}^{2} \int_{a}^{b} \left(V_{i-1}^{k} - V_{i+1}^{k}\right) + \frac{(r-D)i}{2}i\delta_{i}\left(V_{i-1}^{k} - V_{i+1}^{k}\right) - rV_{i}^{k}\delta_{i}$$

$$V_{i+1}^{k} : \frac{1}{2}\left(\sigma_{i}^{2} + (r-D)i\right) \int_{a}^{b} \left(V_{i-1}^{k} - V_{i+1}^{k}\right) + \frac{(r-D)i}{2}i\delta_{i}\left(V_{i-1}^{k} - V_{i+1}^{k}\right) + \frac{(r-D)i}{2}i\delta_{i}\left(V_{i-1}^{k} - V_{i+1}^{k}\right) - rV_{i}^{k}\delta_{i}$$

$$V_{i+1}^{k} : \frac{1}{2}\left(\sigma_{i}^{2} + (r-D)i\right) \int_{a}^{b} \left(V_{i-1}^{k} - V_{i+1}^{k}\right) + \frac{(r-D)i}{2}i\delta_{i}\left(V_{i-1}^{k} - V_{i-1}^{k}\right) + \frac{(r-D)i}{2}i\delta_{i}\left(V_{i-1}^{k} - V_{i+1}^{k}\right) + \frac{(r-D)i}{2}i\delta_{i}\left(V_{i-1}^{k} - V_{i-1}^{k}\right) + \frac{(r-D)i}{2}i\delta_{i}\left(V_{i-1}^{k}\right) + \frac{(r-D)i}{2}i\delta_{i}\left(V_{i-1}^{k} - V_{i-1}^{k}\right) + \frac{(r-D)i}{2}i\delta_{i}\left(V_{i-1}^{k} - V_{i-1}^{k}\right) + \frac{(r-D)i}{2}i\delta_{$$

The relationship between the option values in the algorithm is shown in the figure below.



#### Points to note:


• The time derivative uses the option values at 'times' k and k+1, whereas the other terms all use values at k.


 The gamma term is a central difference, in practice one never uses anything else.


• The delta term uses a central difference. There are often times when a one-sided derivative is better. We'll see examples later.


• The asset- and time-dependent functions a, b and c have been valued at  $S_i=i\,\delta S$  and  $t=T-k\,\delta t$  with the obvious notation.


• The error in the equation is  $O(\delta t, \delta S^2)$ .














## **Boundary conditions**

We must specify the option values at the extremes of the region, at S=0 and at  $S=I\,\delta S$ . They will depend on our option.

### Example 1: Call option at S = 0

At S=0 we know that the value is always zero, therefore

$$V_0^k = 0.$$

## Example 2: Call option for large S

For large S the call value asymptotes to  $S - Ee^{-r(T-t)}$ . Thus

$$V_I^k = I \, \delta S - E e^{-rk \, \delta t}.$$

## Example 3: Put option at S = 0

At 
$$S = 0$$
  $V = Ee^{-r(T-t)}$ . I.e.

$$V_0^k = Ee^{-rk\,\delta t}.$$

## Example 4: Put option for large S

The put option becomes worthless for large S and so

$$V_I^k = 0.$$

More sophisticated requiring more thought

Example 5\*: General condition at S=0

A useful boundary condition to apply at S=0 is that the diffusion

and drift terms 'switch off.'

and drift terms 'switch off.'

i.e put 
$$S=0$$
 in  $S$ . If  $S$  is  $S$  in  $S$  in

at i=0  $\alpha_0 = \gamma_0 = 0$   $\beta_0 = 1-r\partial t$ 



Consider Vikti-divint Bivint + Bivint i= I When the option has a payoff that is linear in the underlying for large S then

$$\frac{\partial^2 V}{\partial S^2}(S,t) \to 0 \text{ as } S \to \infty.$$

The finite-difference representation is

$$V_I^k = 2V_{I-1}^k - V_{I-2}^k.$$
at  $i=T$ 

$$\sim V_{I-1}$$

$$\sim V_{I-1}$$

$$\sim V_{I-1}$$

$$\sim V_{I-1}$$

$$\sim V_{I-1}$$

$$\sim V_{I-1}$$

$$\sim V_{I-2}$$


$$\sim V_{I-1}$$


$$\sim V_{I-2}$$


$$\sim V_{I-1}$$


$$\sim V_{I-2}$$


$$\sim V_{I-1}$$














Certificate in Quantitative Finance

## Other issues



• Early exercise (and other decisions) very 5502 for contable early exercise

1 FDM
29 Mach.

## The advantages of the explicit method

- It is very easy to program and hard to make mistakes
- When it does go unstable it is usually obvious
- It copes well with coefficients that are asset and/or time dependent
- it copes very well with early exercise
- It can be used for modern option-pricing models

## The disadvantages of the explicit method

• There are restrictions on the time step

Bis problem


It is slower than Monte Carlo in high dimensions

terrisle

V(S(t)

exponent

7 10/



## **Summary**

Please take away the following important ideas

There are two main numerical methods for pricing derivatives

Monte Carlo methods exploit the relationship between option prices and expectations

• The finite-difference method solved a discretized version of the Black-Scholes equation

Patricle

Computation

Fluid