Chapter 3. **Higher-Order Differential Equations**

Chuan-Ching Sue

Dept. of Computer Science and Information Engineering, National Cheng Kung University

Bernoulli Equation

• 非線性→線性

原式 :
$$\frac{dz}{dx} + p(x)z = q(x)$$

Bernoulli Equation

例:
$$x^2 \cos y \frac{dy}{dx} = 2x \sin y - 1$$
 $\Rightarrow z = \sin y$ 則 $\frac{dz}{dx} = \cos y \cdot \frac{dy}{dx}$

$$\frac{dz}{dx} - \frac{2}{x} dz = \frac{-1}{x^2}$$

$$z = CI^{-1} + I^{-1} \int Ir dx$$
對於型式(1)
$$f(y)$$
同常等於 $y^2, y^3, \dots, \sin(y) \cdots e^y$

3

Bernoulli Equation

型式(2)

Bernoulli

型式(3)

Riccati

$$\frac{dy}{dx} + p(x)y = q(x) + y^{2}r(x)$$
 若 y_{1} 為上式之一特解 則令 $y = y_{1} + \frac{1}{7}$ 得Z的線性D.E.

Bernoulli Equation

• [A]:
$$y' + (2x - 1)y = x^2 - x + 1 + y^2$$

$$p = 2x - 1$$

$$q = x^2 - x + 1$$

$$r = 1$$

$$find \ y_1 \Rightarrow guess? \ 1, x, x^2, \sin x, \cos x, \cdots$$

$$\therefore y_1 = x$$

$$\Rightarrow y = y_1 + \frac{1}{z} \text{ Apd. E. Apd. E. Apd. E. Apd. In } z = Ce^{-x} - 1$$

$$y = x + \frac{1}{ce^{-x} - 1}$$

E

Review

Chuan-Ching Sue

Dept. of Computer Science and Information Engineering,
National Cheng Kung University

$$M(x,y)dx + N(x,y)dy = 0$$

$$E \stackrel{?}{=} : \frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

$$I(x,y)M(x,y)dx + I(x,y)N(x,y)dy = 0$$

$$M_1 \qquad N_1$$

$$\frac{\partial M_1}{\partial y} = \frac{\partial N_1}{\partial x}$$

$$M(x,y)\frac{\partial I(x,y)}{\partial y} + I(x,y)\frac{\partial M(x,y)}{\partial y} = N(x,y)\frac{\partial I(x,y)}{\partial x} + I(x,y)\frac{\partial N(x,y)}{\partial x}$$

$$M(x,y)\frac{\partial I(x,y)}{\partial y} - N(x,y)\frac{\partial I(x,y)}{\partial x} = +I(x,y)\frac{\partial N(x,y)}{\partial x} - I(x,y)\frac{\partial M(x,y)}{\partial y}$$

Review

$$P(x, y, z) \frac{\partial z}{\partial x} + Q(x, y, z) \frac{\partial z}{\partial y} = R(x, y, z)$$

$$\frac{dx}{P} = \frac{dy}{Q} = \frac{dz}{R} (\text{thinh})$$

$$u(x, y, z) = \alpha$$

一階P.D.E.

$$\upsilon(x,y,z) = \beta$$

$$\varphi(u,v) = 0 or C$$

$$v = f(u)$$

$$\frac{dx}{-N(x,y)} = \frac{dy}{M(x,y)} = \frac{dI}{I\left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}\right)}$$

希望 I(x)

$$\frac{dx}{-N(x,y)} = \frac{dI}{I\left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}\right)}$$

$$\frac{\left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}\right)}{-N} dx = \frac{dI}{I} \quad if \ f(x) = \frac{\left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}\right)}{-N}$$

$$I = e^{\int f(x)dx} = I(x)$$
(ii)

$$\frac{\left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}\right)}{M} dy = \frac{dI}{I} \qquad f(y)dy = \frac{dI}{I}$$

$$I = e^{\int f(y)dy}$$

Review

希望
$$I(x+y)$$

$$\frac{dx + dy}{-N + M} = \frac{dI}{I\left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}\right)}$$

$$\frac{\left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}\right)}{-N + M} d(x + y) = \frac{dI}{I}$$

$$I = e^{\int f(x+y)d(x+y)}$$

(iv)

希望 I(xy)

$$\frac{ydx + xdy}{-yN + xM}d(x, y) = \frac{dI}{I}$$

$$I = e^{\int f(x,y)dxy}$$

(3/

(v)

希望
$$I(x^a y^b)$$

$$d(x^a y^b) = ?$$

$$\frac{d(x^a y^b)}{dx} = ax^{a-1} y^b + bx^a y^{b-1} \frac{dy}{dx}$$

$$d(x^a y^b) = ax^{a-1} y^b dx + bx^a y^{b-1} dy = x^{a-1} y^{b-1} (aydx + bxdy)$$

$$\frac{aydx + bxdy}{-ayN + bxM} = \frac{dI}{I\left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}\right)}$$

$$\frac{\left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}\right) d(x^a y^b)}{(-ayN + bxM)x^{a-1}y^{b-1}} = \frac{\partial I}{I}$$

11

$$if \quad f(x^{a}y^{b}) = \frac{\left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}\right)}{(-ayN + bxM)x^{a-1}y^{b-1}}$$

$$I = e^{\int f(x^{a}y^{b})d(x^{a}y^{b})}$$

$$\frac{\left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}\right)}{(-ayN + bxM)} \cdot \frac{1}{x^{a}y^{b}} d(x^{a}y^{b}) = \frac{dI}{I}$$

$$\frac{\left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}\right)}{xy}$$

$$\frac{\left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}\right)}{xy} = 1 \Rightarrow I = x^{a}y^{b}$$

$$\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} = \frac{(-ayN + bxM)}{xy} = \frac{-aN}{x} + \frac{bM}{y}$$

•
$$[5]: (4xy + 6y^2) dx + (2x^2 + 6xy) dy = 0$$

$$\frac{\partial M}{\partial y} = 4x + 12y, \frac{\partial N}{\partial x} = 4x + 6y, \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} = -6y$$

$$\frac{\partial H}{\partial y} = -6y$$

$$\frac{\partial M}{\partial y} = -6y$$

13

• § 2.2分離變數

§ 2.2分離變數
例4:
$$\cos x(e^{2y} - y) \frac{dy}{dx} = e^y \sin 2x$$

 $(e^x \sin 2x) dx - (\cos x)(e^{2y} - y) dy = 0$
 $\frac{\partial M}{\partial y} = e^y \sin 2x, \frac{\partial N}{\partial x} = \sin x(e^{2y} - y)$
 $\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} = \sin x(e^{2y} - y) - e^y \sin 2x$
 $I = ?$

15

$$e^{y} + ye^{-y} + e^{-y} + 2\cos x = C$$

$$\frac{\partial u}{\partial x} = -2\sin x, \frac{\partial u}{\partial y} = e^{y} + e^{-y} + y(-e^{-y}) + (-e^{-y}) = e^{y} - ye^{-y}$$

$$-2\sin x dx + e^{-y}(e^{2y} - y) dy = 0$$

$$I = -e^{y}\cos x$$

$$\frac{e^{2y} - y}{e^{y}} dy = \frac{\sin 2x}{\cos x} dx$$

$$\int (e^{y} - ye^{-y}) dy = 2\int \sin x dx + C$$

$$e^{y} + ye^{-y} + e^{-y} = -2\cos x + C$$

$$u = C$$

WIELES TO BE THE PARTY OF THE P

• § 2.3線性微分方程式

例1:
$$\frac{dy}{dx}$$
-3y = 6

$$y = CI^{-1} + I^{-1} \int Ir dx$$

$$I = e^{\int -3dx} = e^{-3x}$$

$$y = Ce^{3x} + e^{3x} \int e^{-3x} 6dx$$

$$= Ce^{3x} - 2$$

17

例2:
$$x \frac{dy}{dx} - 4y = x^6 e^x \Rightarrow \frac{dy}{dx} - 4 \frac{y}{x} = x^5 e^x$$

$$I = e^{\int \frac{-4}{x} dx} = e^{-4 \ln x} = e^{\ln x^{-4}} = x^{-4}$$

$$y = CI^{-1} + I^{-1} \int Ir dx = Cx^4 + x^4 \int x^{-4} x^5 e^x dx$$

$$= Cx^4 + x^5 e^x - x^4 e^x$$

$$(5) 3: (x^2 - 9) \frac{dy}{dx} + xy = 0 \Rightarrow \frac{dy}{dx} + \frac{1}{x^2 - 9} xy = 0$$

$$I = e^{\int \frac{x}{x^2 - 9} dx} = e^{\int \frac{1}{2} \frac{x}{x^2 - 9} d(x^2 - 9)} = e^{\frac{1}{2} \ln|x^2 - 9|} = (x^2 - 9)^{\frac{1}{2}}$$

$$y = CI^{-1} = \frac{C}{\sqrt{x^2 - 9}}$$

• § 2.4正合方程式

例1:
$$2xydx + (x^2-1)dy = 0$$

$$\frac{\partial u}{\partial x} = 2xy, \frac{\partial u}{\partial y} = x^2 - 1$$

$$u = x^2y + f(y), u = x^2y - y + g(x)$$

$$f(y) = -y, g(x) = 0$$

$$u = x^2y - y$$

10

$$\begin{aligned}
& \frac{\partial M}{\partial y} = x, \frac{\partial N}{\partial x} = 4x \\
& \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} = 3x \\
& \frac{3xdy}{M} = \frac{dI}{I} \Rightarrow \frac{3dy}{y} = \frac{dI}{I} \\
& I = e^{\int_{y}^{3} dy} = e^{3\ln y} = y^{3}
\end{aligned}$$

$$xy(y^{3})dx + (2x^{2} + 3y^{2} - 20)(y^{3})dy = 0$$

$$xy^{4}dx + (2x^{2}y^{3} + 3y^{5} - 20y^{3})dy$$

$$\frac{\partial u}{\partial x} = xy^{4}, \frac{\partial u}{\partial y} = 2x^{2}y^{3} + 3y^{5} - 20y^{3}$$

$$u = \frac{1}{2}x^{2}y^{4} + f(y), u = \frac{1}{2}x^{2}y^{4} + \frac{1}{2}y^{6} - 5y^{4} + g(x)$$

$$f(y) = \frac{1}{2}y^{6} - 5y^{4}, g(x) = 0$$

$$u = \frac{1}{2}x^{2}y^{4} + \frac{1}{2}y^{6} - 5y^{4} = C$$

21

Review

• § 2.5取代法

$$\boxed{\beta 1} : (x^2 + y^2)dx + (x^2 - xy)dy = 0$$

$$\frac{\partial M}{\partial y} = 2y, \frac{\partial N}{\partial x} = 2x - y$$

$$\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} = 2x - 3y$$

$$\Rightarrow \frac{y}{x} = u, y = ux \Rightarrow dy = udx + xdu$$

$$(x^2 + u^2x^2)dx + (x^2 - ux^2)(udx + xdu) = 0$$

$$(x^2 + u^2x^2 + ux^2 - u^2x^2)dx + (x^3 - ux^3)du = 0$$

$$x^2(1 + u)dx + x^3(1 - u)du = 0$$

$$(1 + u)dx + x(1 - u)du = 0$$

$$\frac{\partial M}{\partial u} = 1, \frac{\partial N}{\partial x} = 1 - u$$

$$\frac{1 - u}{1 + u} du + \frac{1}{x} dx = 0$$

$$\frac{1}{x} dx = -\left(\frac{1 - u}{1 + u}\right) du$$

$$\ln x = -\left(-1 + \frac{2}{1 + u}\right) du$$

$$\ln x = -\left(-u + 2\ln|1 + u|\right) + C$$

23

$$|x| = \frac{y}{x}$$

$$\ln x = u - 2\ln\left|1 + \frac{y}{x}\right| + C$$

$$\ln x + 2\ln\left|1 + \frac{y}{x}\right| = \frac{y}{x} + \ln C$$

$$\frac{x(1 + \frac{y}{x})^2}{C'} = e^{\frac{y}{x}}$$

$$(x + y)^2 = C'xe^{\frac{y}{x}}$$

P.D.E

「何」:
$$x \frac{\partial z}{\partial x} + \frac{\partial z}{\partial y} - 3z = 0$$
, Initial Condition $z(x,1) = x^2 e^{x^2}$

$$\frac{dx}{x} = \frac{dy}{y} = \frac{dz}{3z}$$

$$\frac{dx}{x} = \frac{dy}{y} \Rightarrow \frac{y}{x} = \alpha = u$$

$$\frac{dx}{x} = \frac{dz}{3z} \Rightarrow \frac{z}{x^3} = \beta = v$$
通解 $\Phi(u, v) = 0, \frac{z}{x^3} = f(\frac{y}{x})$

$$z(x,1) 帶入通解把f求出
$$\therefore z(x,1) = x^2 e^{x^2}$$

$$z = x^3 f(\frac{y}{x})$$$$

2.

Review

$$z(x,1) = x^3 f(\frac{1}{x}) = x^2 e^{x^2}$$

$$f(\frac{1}{x}) = \frac{1}{x} e^{x^2}$$

$$\Rightarrow \mu = \frac{1}{x} || |x = \frac{1}{\mu}|$$

$$f(\mu) = \mu e^{\frac{1}{\mu^2}}$$

$$\therefore z = x^3 f(\frac{y}{x})$$

$$= x^3 \frac{y}{x} e^{\frac{(\frac{y}{y})^2}{x^2}}$$

$$= x^2 v e^{\frac{x^2}{y^2}}$$

學學學學

• Chap3

常O.D.E

法一U.C 3-3, 3-4

法二R.O. 補充

法三D.D. 補充

法四VV 3-5

27

Review

夕!
$$y'' - 6y' + 9y = 6x^2 + 2 - 12e^{3x}$$

 $y = y_h + y_p$
 $\lambda^2 - 6\lambda + 9 = 0 \Rightarrow \lambda = 3,3$
 $y_h = C_1 e^{3x} + C_2 x e^{3x}$
法四: $y_p = y_{p1}, y_{p2}$
 $w = \begin{vmatrix} e^{3x} & x e^{3x} \\ 3e^{3x} & e^{3x} + 3x e^{3x} \end{vmatrix} = e^{6x} + 3x e^{6x} - 3x e^{6x} = e^{6x}$
 $y_{p1} = e^{3x} \int \frac{-(6x^2 + 2)x e^{3x}}{e^{6x}} dx + x e^{3x} \int \frac{(6x^2 + 2)e^{3x}}{e^{6x}} dx$
 $= e^{3x} \int -(6x^2 + 2)x e^{-3x} dx + x e^{3x} \int (6x^2 + 2)e^{-3x} dx$
 $= \frac{14}{3}$ 方法

法三:

$$y_{p1} = \frac{1}{D^2 - 6D + 9} (6x^2 + 2)$$

$$= \frac{1}{9(1 + \frac{D^2 - 6D}{9})} (6x^2 + 2)$$

$$= \frac{1}{9} \left[1 - \frac{D^2 - 6D}{9} + (\frac{D^2 - 6D}{9})^2 - (\frac{D^2 - 6D}{9})^3 + \dots\right] (6x^2 + 2)$$

$$= \frac{1}{9} (6x^2 + 2 - \frac{12 - 72x}{9} + \frac{36 + 12}{81})$$

$$= \frac{1}{9} (6x^2 + 8x + 6) = \frac{2}{3}x^2 + \frac{8}{9}x + \frac{2}{3}$$

20

$$y_{p2} = \frac{1}{D^2 - 6D + 9} (-12e^{3x})$$

$$= e^{3x} \frac{1}{(D+3)^2 - 6(D+3) + 9} (-12)$$

$$= (-12)e^{3x} \frac{1}{D^2} \times 1$$

$$= e^{3x} (-6x^2)$$

$$= -6x^2 e^{3x}$$