

Quiz (III)

Finished by 18:30 on 5/17

Create a matlab script F7xxxxxxx_quiz3.m and link all the programs to this script. Make sure once type the filename 'F7xxxxxxx_quiz3', the results of the following problems will pop-up automatically in order. Remember not to type any 'clear all' or 'close all' command in any of the codes.

1. [F7xxxxxxx_quiz4_prob1.m & Figure 1]

Given the definite integral

$$\int_{1/4}^{1/2} \pi \cos \pi x \, dx$$

Use box counting and Trapezoid method to analyze the error with different h values, say for example from $h = 10^{-100}$ to 10^{-1} .

- (1) Plot the ABSOLUTE ERROR versus h in Log-Log scale with appropriate figure legends to indicate correspondence between the error curve and the method.
- (2) Print the integration with 8 significant digits. Which h is an optimal choice if 8 significant digits are required for the method selected?

2. [F7xxxxxxx_quiz4_prob2.m & Figure 2 (if you use pictorial method to find the optimal h)]

Given the definite integral

$$\int_0^2 e^{-x^2} \, dx$$

- (1) Find the answer with 8 digit precision using Simpson's Method.
- (2) Which h is optimal for the Simpson's Method? Explain how you find the optimal h.

3. Use Monte Carlo Method to estimate the area of the ellipse enclosed by $\frac{x^2}{100} + \frac{y^2}{64} = 1$.

How many points does it require for the result to achieve 2-digit precision?

