

ASSIGNMENT 3

EE24BTECH11003 - Akshara Sarma Chennubhatla

D: MCQs WITH ONE OR MORE THAN ONE
CORRECT

- 3) The probability that at least one of the events A and B occurs is 0.6. If A and B occur simultaneously with probability 0.2, then $\Pr(\bar{A}) + \Pr(\bar{B})$ is (1987 – 2Marks)
- a) 0.4 b) 0.8 c) 1.2 d) 1.4
- e) none
(Here \bar{A} and \bar{B} are the complements of A and B, respectively).
- 4) For two given events A and B, $\Pr(A \cap B)$ (1988 – 2Marks)
- a) not less than $\Pr(A) + \Pr(B) - 1$
b) not greater than $\Pr(A) + \Pr(B)$
c) equal to $\Pr(A) + \Pr(B) - \Pr(A \cup B)$
d) equal to $\Pr(A) + \Pr(B) + \Pr(A \cup B)$
- 5) If E and F are independent events such that $0 < \Pr(E) < 1$ and $0 < \Pr(F) < 1$, then (1989 – 2Marks)
- a) E and F are mutually exclusive
b) E and F^c (the complement of the event F) are independent
c) E^c and F^c are independent
d) $\Pr(E|F) + \Pr(E^c|F^c) = 1$.
- 6) For any two events A and B in a sample space (1991 – 2Marks)
- a) $\Pr(A|B) \geq \frac{\Pr(A) + \Pr(B) - 1}{\Pr(B)}$, $\Pr(B) \neq 0$ is always true
b) $\Pr(A \cap \bar{B}) = \Pr(A) - \Pr(A \cap B)$ does not hold
c) $\Pr(A \cup B) = 1 - \Pr(\bar{A})\Pr(\bar{B})$, if A and B are independent
d) $\Pr(A \cup B) = 1 - \Pr(\bar{A})\Pr(\bar{B})$, if A and B are disjoint.
- 7) E and F are two independent events. The probability that both E and F happen is $\frac{1}{12}$ and the probability that neither E nor F happens is $\frac{1}{2}$. Then, (1993 – 2Marks)
- a) $\Pr(E) = \frac{1}{3}, \Pr(F) = \frac{1}{4}$
b) $\Pr(E) = \frac{1}{2}, \Pr(F) = \frac{1}{6}$
c) $\Pr(E) = \frac{1}{6}, \Pr(F) = \frac{1}{2}$
d) $\Pr(E) = \frac{1}{4}, \Pr(F) = \frac{1}{3}$
- 8) Let $0 < \Pr(A) < 1, 0 < \Pr(B) < 1$ and $\Pr(A \cup B) = \Pr(A) + \Pr(B) - \Pr(A)\Pr(B)$ then (1995S)
- a) $\Pr(A|B) = \Pr(B) - \Pr(A)$
b) $\Pr(A' - B') = \Pr(A') - \Pr(B')$
c) $\Pr(A \cup B)' = \Pr(A')\Pr(B')$
d) $\Pr(A|B) = \Pr(A)$
- 9) If from each of the three boxes containing 3 white and 1 black, 2 white and 2 black, 1 white and 3 black balls, one ball is drawn at random, then the probability that 2 white and 1 black ball will be drawn is (1998 – 2Marks)
- a) $\frac{13}{32}$ b) $\frac{1}{4}$ c) $\frac{1}{32}$ d) $\frac{3}{16}$
- 10) If \bar{E} and \bar{F} are the complementary events of events E and F respectively and if $0 < \Pr(F) < 1$, then (1998 – 2Marks)
- a) $\Pr(E|F) + \Pr(\bar{E}|\bar{F}) = 1$
b) $\Pr(E|F) + \Pr(\bar{E}|F) = 1$
c) $\Pr(\bar{E}|F) + \Pr(E|\bar{F}) = 1$
d) $\Pr(E|\bar{F}) + \Pr(\bar{E}|\bar{F}) = 1$
- 11) There are four machines and it is known that exactly two of them are faulty. They are tested, one by one, in a random order till both the faulty machines are identified. Then the probability that only two tests are needed is (1998 – 2Marks)
- a) $\frac{1}{3}$ b) $\frac{1}{6}$ c) $\frac{1}{2}$ d) $\frac{1}{4}$
- 12) If E and F are events with $\Pr(E) \leq \Pr(F)$ and $\Pr(E \cap F) > 0$, then (1998 – 2Marks)
- a) occurrence of E \Rightarrow occurrence of F
b) occurrence of F \Rightarrow occurrence of E
c) non-occurrence of E \Rightarrow non-occurrence of F
d) none of the above implications holds
- 13) A fair coin is tossed repeatedly. If the tail appears on first four tosses, then the probability of the head appearing on the fifth toss equals (1998 – 2Marks)

- a) $\frac{1}{2}$ b) $\frac{1}{32}$ c) $\frac{31}{32}$ d) $\frac{1}{5}$

14) Seven white balls and three black balls are randomly placed in a row. The probability that no two black balls are placed adjacently equals (1998 – 2Marks)

- a) $\frac{1}{2}$ b) $\frac{7}{15}$ c) $\frac{2}{15}$ d) $\frac{1}{3}$

15) The probabilities that a student passes in Mathematics, Physics and Chemistry are m , p and c , respectively. Of these subjects, the student has a 75% chance of passing in at least one, a 50% chance of passing in at least two, and a 40% chance of passing in exactly two. Which of the following relations are true? (1999 – 3Marks)

- a) $p + m + c = \frac{19}{20}$ b) $p + m + c = \frac{27}{20}$
 c) $pmc = \frac{1}{10}$ d) $pmc = \frac{1}{4}$

16) Let E and F be two independent events. The probability that exactly one of them occurs is $\frac{11}{25}$ and the probability of none of them occurring is $\frac{2}{25}$. If $\Pr(T)$ denotes the probability of occurrence of the event T , then (2011)

- a) $\Pr(E) = \frac{4}{5}, \Pr(F) = \frac{3}{5}$ b) $\Pr(E) = \frac{1}{5}, \Pr(F) = \frac{2}{5}$
 c) $\Pr(E) = \frac{2}{5}, \Pr(F) = \frac{1}{5}$ d) $\Pr(E) = \frac{3}{5}, \Pr(F) = \frac{4}{5}$

17) A ship is fitted with three engines E_1, E_2 and E_3 . The engines function independently of each other with respective probabilities $\frac{1}{2}, \frac{1}{4}$ and $\frac{1}{4}$. For the ship to be operational at least two of its engines must function. Let X denote the event that the ship is operational and let X_1, X_2 and X_3 denote respectively the events that the engines E_1, E_2 and E_3 are functioning. Which of the following is(are) true? (2012)

- a) $\Pr(X_1^c | X) = \frac{3}{16}$
 b) $\Pr(\text{Exactly two engines of the ship are functioning} | X) = \frac{7}{8}$
 c) $\Pr(X | X_2) = \frac{5}{16}$
 d) $\Pr(X | X_1) = \frac{7}{16}$