

# **MATH-505A: Homework # 3**

Due on Friday, September 12, 2014

**Saket Choudhary**  
**2170058637**

## Contents

<b>Exercise # 1.7</b>	<b>3</b>
(1) . . . . .	3
(2) . . . . .	4
(4) . . . . .	4
(5) . . . . .	4
[ . . . . .	5
<b>1</b>	<b>5</b>
(a) . . . . .	5
(b) . . . . .	5
(c) . . . . .	5
(d) . . . . .	5
<b>2</b>	<b>5</b>
(a) . . . . .	5
(b) . . . . .	6
(c) . . . . .	6
(d) . . . . .	6
<b>4</b>	<b>6</b>
(a) . . . . .	6
(b) . . . . .	6
(c) . . . . .	6
5 . . . . .	7
6 . . . . .	7
14 . . . . .	7

**Exercise # 1.7****(1)**

**Given:** Two roads  $r1_{AB}$ ,  $r2_{AB}$  connecting points A and B and  $s1_{BC}$ ,  $s2_{BC}$  connecting B and C. Let  $p(AB)$  denote the probability that path between A  $\rightarrow$  B is open and let  $p(AB^c)$  denote the probability that there is(are) no path open b/w A and B.

**To find:**  $Y = P(AB|AC^c)$ .

Y is equal to the probability that path between A and B is open AND(given that) the path between A and C is closed  $\implies$  Path between B and C is closed AND between A and B is open

$p(AB) = \text{Path b/w A,B is open} = 1 - \text{Path b/w A,B is closed} = 1 - p * p$

$p(AC^c|AB) = \text{Path b/w A,C is closed given A,B is open} = \text{Path b/w B,C is closed given A,B is open.} = p(BC^c)$

Thus

$$p(AB) = 1 - p^2 \quad (1)$$

Also,

$$p(AB) = p(BC) \quad (2)$$

$p(AC^c) = 1 - \text{Probability A,C is open} = 1 - \text{Probability AB is open AND BC is open.}$  Thus,

$$p(AC^c) = 1 - p(AB)p(AC) = 1 - (1 - p^2)^2 \quad (3)$$

$$p(AB \cap AC^c) = p(AC^c|AB)p(AB) = p(BC^c)p(AB) = p^2(1 - p^2) \quad (4)$$

$$p(AB|AC^c) = \frac{P(AB \cap AC^c)}{p(AC^c)} = \frac{p(AC^c|AB)p(AB)}{p(AC^c)} = \frac{p^2(1 - p^2)}{1 - (1 - p^2)^2} \quad (5)$$

**Part 2:** Additional direct road from A to C. Find  $p(AB|AC^c)$ :

$p(AC^c|AB) = \text{Probability that path b/w A,C is closed given path b/w A,B are open} = \text{Probability path b/w A,C(direct) are closed AND path b/w B,C are closed}$

$$p(AC^c|AB) = p * p(BC^c)p(AB) \quad (6)$$

where the extra p in 6 as compared to 4 is because the direct path A,C should be blocked too.

$$p(AC^c) = 1 - (p(1 - p^2)^2 - (1 - p)) \quad (7)$$

where the second term is the probability that direct A  $\rightarrow$  C is blocked, but A  $\longleftrightarrow$  B and B  $\rightarrow$  is open(both). The third term represents the probability that A  $\rightarrow$  C is open directly(does not matter if the other paths are open or blocked),

Thus, for part 2:

$$p(AB|AC^c) = \frac{p^3(1 - p^2)}{1 - (1 - p^2)^2 - (1 - p)} = \frac{p^2(1 - p^2)}{1 - (1 - p^2)^2} \quad (8)$$

(2)

$$p(2K \cap 1A) = \frac{\binom{4}{2} * \binom{4}{1} * \binom{52-4-4}{10}}{\binom{52}{13}} \quad (9)$$

$$p(1A|2K) = \frac{p(1A \cap 2K)}{p(2K)}$$

$$p(2K) = \frac{\binom{4}{2} * \binom{52-4}{11}}{\binom{52}{13}} = \frac{6 * 48! * 13!}{52! * 11!} = \frac{6 * 12 * 13}{49 * 50 * 51 * 52} = 2.52 * 10^{-4} \quad (10)$$

Thus,

$$p(1A|2K) = \frac{\binom{4}{2} * \binom{4}{1} * \binom{44}{10}}{\binom{4}{2} * \binom{48}{11}} = \frac{4 * 11 * 37!}{46 * 47 * 48 * 34!} = 0.0399$$

(4)

**To prove/disprove:**  $p(x|C) > p(y|C) \text{ AND } p(x|C^c) > p(y|C^c) \implies p(x) > p(y)$

$$p(x|C) - p(y|C) > 0 \quad (11)$$

$$p(x|C^c) - p(y|C^c) > 0 \quad (12)$$

$$p(x) = p(x|C)p(C) + p(x|C^c)p(C^c) \quad (13)$$

Also,

$$p(y) = p(y|C)p(C) + p(y|C^c)p(C^c) \quad (14)$$

Consider  $p(x) - p(y)$  :

$$p(x) - p(y) = (p(x|C) - p(y|C))p(C) + (p(x|C^c) - p(y|C^c))p(C^c) \quad (15)$$

From the 12, ?? and 15:

$$p(x) - p(y) > 0 \forall x, y \quad (16)$$

**Thus, x is always preferred over y.**

(5)

Let  $X_i$  represent the  $i^{th}$  card draw**Given:**  $X_k > X_i, \forall i \in [1, k-1] \text{ and } k \in [1, m]$ 

$$p(X_k = m | X_k > X_i) = \frac{p(X_k = m \cap X_k > X_i)}{p(X_k > X_i)} = \frac{p(X_k = m)}{p(X_k > X_i)} = \frac{\frac{1}{m}}{\frac{1}{k}}$$

Where the equality in the last step comes from the fact that the probability of choosing cards such that  $p(X_k > X_i)$  is simply to choose the largest card, i.e.  $k$  among the rest  $i$ .

Thus  $p(X_k = m | X_k > X_i) = \frac{k}{m}$ .

[  
Exercise # 1.8]

## 1

(a)

**Six turns up exactly once** The dice on which 6 should appear is chosen in  $\binom{2}{1}$  way and has just one option(6) while the other has  $6 - 1 = 5$  options. Total outcomes =  $6 * 6 = 36$

Thus:

$$p = \frac{1*5}{36} = \frac{5}{36}$$

(b)

**Both numbers are odd.** 3 out of 6 numbers are odd and the outcome of odd on both dice are independent. Thus,

$$p = \frac{3*3}{36} = \frac{1}{12}$$

(c)

**Sum of scores is 4.**

Possible configurations: (1, 3); (2, 2); (3, 1)

Thus:

$$p = \frac{3}{36} = \frac{1}{12}$$

(d)

**Sum of scores is divisible by 3.**

Possible choices for sum to be divisible by 3:

3, 6, 9, 12:

3: (1, 2), (2, 1)

6: (1, 5), (2, 4), (3, 3), (4, 2), (5, 1)

9: (3, 6), (4, 5), (5, 4), (6, 3)

12: (6, 6)

$$\text{Thus } p = \frac{2+5+4+1}{36} = \frac{1}{3}$$

## 2

(a)

Head appears the first time on  $n^{th}$  throw. Thus a series of  $n - 1$  consecutive tails followed by head.

$$p = \binom{n}{1} \frac{1}{2^{n-1}} * \frac{1}{2} = \frac{n}{2^n}$$

(b)

The number of heads and tails to date are equal:

**Case1:**  $n$  is odd.

Clearly the probability of heads and tails being equal in this case is zero!

**Case2:**  $n$  is even.

A series of H,T such that  $|H| = |T|$ . Since the probability of occurrence of either a H or a T is  $\frac{1}{2}$ , the result will still be:

$$p = \binom{n}{n/2} \frac{1}{2^n}$$

(c)

Exactly two heads have appeared altogether.

$$p = \binom{n}{2} * \frac{1}{2^2} * \frac{1}{2^{n-2}} = \binom{n}{2} * \frac{1}{2^n}$$

(d)

At least two heads have appeared.

This is same as the one minus probability that 0 or 1 heads have appeared so far:

$$p = 1 - p(0 \text{ heads}) - p(1 \text{ head only}) \quad p = 1 - \binom{n}{0} \frac{1}{2^0} \frac{1}{2^n} - \binom{n}{1} \frac{1}{2} \frac{1}{2^{n-1}} = 1 - \frac{1}{2^n} - \frac{n}{2^n}$$

4

(a)

**Biased coin tossed three times:**

Sample space: HHH,HHT,HHT,HTT,THH,THT,TTH,TTT

(b)

**Balls drawn without replacement from 2U,2V balls**

**Sample space:**  $(U1, V1), (U2, V1), (V1, U1), (V1, U2), (U1, V2), (U2, V2), (V2, U1), (V2, U2)$

(c)

**Biased coin tossed till H turns up**

Sample Space:

$\{H\}, \{T, H\}, \{T, T, H\}, \{T, T, T, H\}, \dots, \{T, T, T, \dots, T, H\}$

## 5

**To find:** Probability that exactly one of A,B occurs

Probability that exactly one of A,B occurs is equal to the probability that either **only A** occurs or **only B** occurs. This is simply given by:

$P(A \cup B) - P(A \cap B)$ . That is either of A,B occurs removing the portion when both A,B occur.

Thus, the probability that exactly one of A,B occurs is :

$$p(A \cup B) - p(A \cap B) = p(A) + p(B) - p(A \cap B) - p(A \cap B) = p(A) + p(B) - 2p(A \cap B)$$

## 6

**To Prove:**  $P(A \cup B \cup C) = 1 - P(A^C|B^C \cap C^C)p(B^C|C^C)p(C^C)$   $RHS = 1 - P(A^C|B^C \cap C^C)P(B^C|C^C)p(C^C)$

$$RHS = 1 - P(A^C|B^C \cap C^C) \frac{P(B^C \cap C^C)}{p(C^C)} p(C^C)$$

Thus, expanding further  $p(A^C|B^C \cap C^C) = \frac{p(A^C \cap B^C \cap C^C)}{p(B^C \cap C^C)}$

Thus,

$$RHS = 1 - \frac{p(A^C \cap B^C \cap C^C)}{p(B^C \cap C^C)} * \frac{P(B^C \cap C^C)}{p(C^C)} * p(C^C)$$

$$RHS = 1 - P(A^C \cap B^C \cap C^C) \text{ which is same as } LHS$$

## 14

Consider only two events:  $A, B_j$ .

From the definition of conditional probability:

$$p(A_j|B) = \frac{p(A_j \cap B)}{p(B)} \quad (17)$$

Also:

$$p(B|A_j) = \frac{p(A_j \cap B)}{p(A_j)} \quad (18)$$

Substituting for  $p(A_j \cap B)$  from 17 in 18:

$$p(B|A_j) = \frac{p(A_j|B)p(B)}{p(A_j)} \quad (19)$$

or equivalently,

$$p(A_j|B) = \frac{p(B|A_j)p(A_j)}{p(B)} \quad (20)$$

where  $p(B)$  can be expressed as(using the law of partition):

$$p(B) = p(B|A_1)P(A_1) + P(B|A_2)P(A_2) + \dots + P(B|A_n)P(A_n) \quad (21)$$

Thus  $p(B) = \sum_1^n p(B|A_i)p(A_i)$

Ans hence,

$$p(A_j|B) = \frac{p(B|A_j)p(A_j)}{\sum_1^n p(B|A_i)p(A_i)}$$