

MATH 542 Homework 3

Saket Choudhary
skchoudh@usc.edu

January 29, 2016

Problem 1

Given: $A_{n \times n}$ is idempotent $\implies AA = A$; P is non-singular and $C_{n \times n}$ is orthogonal $\implies CC^T = C^TC = I$

1.a

$$\begin{aligned}(I - A)(I - A) &= I - A - A + AA \\ &= I - 2A + A \text{ using } AA = A \\ &= I - A\end{aligned}$$

Hence $I - A$ is idempotent

1.b

$$\begin{aligned}A(I - A) &= A - AA \\ &= A - A \text{ using } AA = A \\ &= 0\end{aligned}$$

Similarly,

$$\begin{aligned}(I - A)A &= A - AA \\ &= A - A \text{ using } AA = A \\ &= 0\end{aligned}$$

1.c

$$\begin{aligned}(P^{-1}AP)(P^{-1}AP) &= P^{-1}APP^{-1}AP \\ &= P^{-1}AAP \text{ since } PP^{-1} = I \\ &= P^{-1}AP \text{ using } AA = A\end{aligned}$$

Hence $P^{-1}AP$ is idempotent

1.d

$$\begin{aligned}
(C'AC)(C'AC) &= C'ACC'AC \\
&= C'AAC \text{ since } CC' = C'C = I \\
&= C'AC \text{ using } AA = A
\end{aligned}$$

1.e

A is a projection matrix and $C'C = C'C = I$ For $C'AC$ to be a projection matrix $C'ACz = z \forall z \in S$ for some vector space S

For some $z \in S$:

$$\begin{aligned}
C'ACz &= C'Ay \text{ where } y = Az \\
&= C'Ay \\
&= C'y \text{ since } A \text{ is projection matrix} \\
&= C'Cz \\
&= z \text{ since } C'C = I
\end{aligned}$$

Thus, $C'ACz \in S$ and $C'ACz = z$ and hence $C'AC$ is a projection matrix

Problem 2

2.a

$$\begin{aligned}
Q(x_1, x_2, x_3) &= 12x_1^2 + 3x_2^2 + 3x_3^2 + 2x_1x_2 - 10x_1x_3 + 4x_2x_3 \\
&= 3(2x_1 + x_2 + x_3)^2 - (12x_1x_2 + 6x_2x_3 + 12x_1x_3) + 2x_1x_2 - 10x_1x_3 + 4x_2x_3 \\
&= 3(2x_1 + x_2 + x_3)^2 - 10x_1x_2 - 2x_2x_3 - 22x_1x_3 \\
&= (x_1 + x_2)^2 + 2(x_2 + x_3)^2 + (5x_1 - x_3)^2 - 14x_1^2 \\
&= (x_1 + x_2)^2 + 2(x_2 + x_3)^2 + x_3^2 + 11x_1^2 - 10x_1x_3 \\
&= (x_1 + x_2)^2 + 2(x_2 + x_3)^2 + (5x_1 - x_3)^2 - 14x_1^2
\end{aligned}$$

and hence $Q(x_1, x_2, x_3)$ is neither positive definite nor positive semidefinite.

2.b

$$\begin{aligned}
x'Ax &= \begin{pmatrix} x_1 & x_2 & x_3 \end{pmatrix} \begin{pmatrix} 2 & 1 & 2 \\ 1 & 2 & 1 \\ 2 & 1 & 4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \\
&= (2x_1 + x_2 + x_3)x_1 + (x_1 + 2x_2 + x_3)x_2 + (2x_1 + x_2 + 4x_3)x_3 \\
&= 2x_1^2 + 2x_2^2 + 4x_3^2 + 2x_1x_2 + 3x_1x_3 + 2x_2x_3 \\
&= (x_1 + x_2)^2 + (x_2 + x_3)^2 + (x_1 + \frac{3}{2}x_2)^2 + \frac{3}{4}x_2^2 \\
&\geq 0
\end{aligned}$$

A is positive semidefinite

2.c

$$\begin{aligned}
 x'Ax &= \begin{pmatrix} x_1 & x_2 & x_3 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 1 \\ 3 & 1 & -2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \\
 &= (x_1 + 2x_2 + 3x_3)x_1 + (2x_1 + x_2 + x_3)x_2 + (3x_1 + x_2 - 2x_3)x_3 \\
 &= x_1^2 + x_2^2 - 2x_3^2 + 4x_1x_2 + 6x_1x_3 + 2x_2x_3 \\
 &= (x_2 + x_3)^2 + 3(x_1 + x_3)^2 + 2(x_1 + x_2)^2 - 6x_1^2 - 6x_3^2 - 4x_1^2 - 2x_2^2
 \end{aligned}$$

A is neither positive definite nor positive semidefinite.

Problem 3

$$\begin{aligned}
 \begin{pmatrix} a & b \\ c & d \end{pmatrix} \times \begin{pmatrix} a & b \\ c & d \end{pmatrix} &= \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix} \\
 \begin{pmatrix} a^2 + b^2 & ac + bd \\ ac + bd & c^2 + d^2 \end{pmatrix} &= \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}
 \end{aligned}$$

$$\begin{aligned}
 a^2 + b^2 &= 2 \\
 ac + bd &= -1 \\
 c^2 + d^2 &= 2
 \end{aligned}$$

$$\text{Let } a = 1, b = 1 \implies c + d = 1$$

$$\begin{aligned}
 c^2 + d^2 &= 2 \\
 c^2 + (-1 - c)^2 &= 2 \\
 2c^2 + 2c - 1 &= 0 \\
 c &= \frac{-2 \pm \sqrt{4 + 8}}{4} \\
 c &= \frac{-1 \pm \sqrt{3}}{2} \\
 d &= \frac{-1 \pm \sqrt{3}}{2}
 \end{aligned}$$

and hence one possible B is:

$$B = \begin{pmatrix} 1 & 1 \\ \frac{-1-\sqrt{3}}{2} & \frac{-1+\sqrt{3}}{2} \end{pmatrix}$$

Problem 4

$$G = (X'X)^{-}$$

Let $Y = (X'X)$ so that $G = Y^{-}$

$$\begin{aligned} G(X'X)G &= GYG \\ &= Y^{-}YY^{-} \\ &= Y^{-} \end{aligned}$$

Note that Y is symmetric, $Y = Y^T$ (Proof: $Y = X^T X$, $Y^T = (X^T X)^T = X^T X$)

Since Y is symmetric, Y^{-} is symmetric too $\implies G(X'X)G$ is symmetric.

To prove $P = G(X'X)G$ is a generalised inverse of $Y = X'X$ we need to show $YPY = P$

$$\begin{aligned} YPY &= YGYGY \\ &= Y(Y^{-})Y \text{ using } GYG = Y^{-} \text{ from the previous result} \\ &= Y \text{ since } YY^{-} = I \end{aligned}$$

Problem 5

Given: $E[X] = 1$ and $Var(X) = E[X^2] - E[X]^2 = 5$ Using $E[X] = 1$ we get, $E[X^2] = 5 + E[X]^2 = 6$

5.a

$$\begin{aligned} E[(1 + 2X)^2] &= E[1 + 4X + 4X^2] \\ &= E[1] + E[4X] + E[4X^2] \\ &= 1 + 4E[X] + 4E[X^2] \\ &= 1 + 4 + 4(6) \\ &= 30 \end{aligned}$$

5.b

$$\begin{aligned} var(3 + 4X) &= Var(3) + Var(4X) + 2Cov(3, 4X) \\ &= 0 + 16Var(X) + 2(0) \\ &= 80 \end{aligned}$$