

ANKRD11 And KBG Syndrome

<https://medlineplus.gov/genetics/gene/ankrd11/>

You Are Here:

Home

→

Genetics

→

Genes

→

ANKRD11 gene

ANKRD11 gene

ankyrin repeat domain containing 11

To use the sharing features on this page, please enable JavaScript.

Normal Function

The ANKRD11 gene provides instructions for making a protein called ankyrin repeat domain 11 (ANKRD11). As its name suggests, this protein contains multiple regions called ankyrin domains; proteins with these domains help other proteins interact with each other. The ANKRD11 protein interacts with certain proteins called histone deacetylases, which are important for controlling gene activity. Through these interactions, ANKRD11 affects when genes are turned on and off. For example, ANKRD11 brings together histone deacetylases and other proteins called p160 coactivators. This association regulates the ability of p160 coactivators to turn on gene activity. ANKRD11 may also enhance the activity of a protein called p53, which controls the growth and division (proliferation) and the self-destruction (apoptosis) of cells. The ANKRD11 protein is found in nerve cells (neurons) in the brain. During embryonic development, ANKRD11 helps regulate the proliferation of these cells and development of the brain. Researchers speculate that the protein may also be involved in the ability of neurons to change and adapt over time (plasticity), which is

important for learning and memory. ANKRD11 may function in other cells in the body and appears to be involved in normal bone development.

Health Conditions Related to Genetic Changes

KBG syndrome

Several ANKRD11 gene mutations have been found to cause KBG syndrome, a condition characterized by large upper front teeth and other unusual facial features, skeletal abnormalities, and intellectual disability. Most of these mutations lead to an abnormally short ANKRD11 protein, which likely has little or no function. Reduction of this protein's function is thought to underlie the signs and symptoms of the condition. Because ANKRD11 is thought to play an important role in neurons and brain development, researchers speculate that a partial loss of its function may lead to developmental delay and intellectual disability in KBG syndrome. However, the mechanism is not fully known. It is also unclear how loss of ANKRD11 function leads to the skeletal features of the condition. Another type of mutation that affects the ANKRD11 gene, called 16q24.3 microdeletion, deletes genetic material from chromosome 16 in a region designated q24.3. The deleted region typically removes the ANKRD11 and ZNF778 genes, although other nearby genes may also be affected. People with this type of mutation have similar signs and symptoms to those of KBG syndrome, including unusual facial features and intellectual disability. Many also have brain abnormalities and features of autism spectrum disorders, which are characterized by impaired communication and socialization skills. Some researchers think that these microdeletions are different enough from KBG syndrome to be considered a separate disorder, called 16q24.3 microdeletion syndrome.

More About This Health Condition

Other Names for This Gene

ANCO-1 ANCO1 ankyrin repeat domain-containing protein 11 ankyrin repeat-containing cofactor 1
LZ16 nasopharyngeal carcinoma susceptibility protein T13

Additional Information & Resources

Tests Listed in the Genetic Testing Registry

Tests of ANKRD11

Scientific Articles on PubMed

PubMed

Catalog of Genes and Diseases from OMIM

ANKYRIN REPEAT DOMAIN-CONTAINING PROTEIN 11; ANKRD11

Gene and Variant Databases

NCBI Gene

ClinVar

References

Gallagher D, Voronova A, Zander MA, Cancino GI, Bramall A, Krause MP, Abad C,

Tekin M, Neilsen PM, Callen DF, Scherer SW, Keller GM, Kaplan DR, Walz K, Miller FD. Ankrd11 is a chromatin regulator involved in autism that is essential for neural development. *Dev Cell*. 2015 Jan 12;32(1):31-42. doi: 10.1016/j.devcel.2014.11.031. Epub 2014 Dec 31. Citation on PubMed

Kim HJ, Cho E, Park JB, Im WY, Kim HJ. A Korean family with KBG syndrome identified by ANKRD11 mutation, and phenotypic comparison of ANKRD11 mutation and 16q24.3 microdeletion. *Eur J Med Genet*. 2015 Feb;58(2):86-94. doi: 10.1016/j.ejmg.2014.11.003. Epub 2014 Nov 20. Citation on PubMed

Neilsen PM, Cheney KM, Li CW, Chen JD, Cawrse JE, Schulz RB, Powell JA, Kumar R, Callen DF. Identification of ANKRD11 as a p53 coactivator. *J Cell Sci*. 2008 Nov 1;121(Pt 21):3541-52. doi: 10.1242/jcs.026351. Epub 2008 Oct 7. Citation on PubMed

Sacharow S, Li D, Fan YS, Tekin M. Familial 16q24.3 microdeletion involving ANKRD11 causes a KBG-like syndrome. *Am J Med Genet A*. 2012 Mar;158A(3):547-52. doi: 10.1002/ajmg.a.34436. Epub 2012 Feb 3. Citation on PubMed

Sirmaci A, Spiliopoulos M, Brancati F, Powell E, Duman D, Abrams A, Bademci G, Agolini E, Guo S, Konuk B, Kavaz A, Blanton S, Digilio MC, Dallapiccola B, Young J, Zuchner S, Tekin M. Mutations in ANKRD11 cause KBG syndrome, characterized by intellectual disability, skeletal malformations, and macrodontia. *Am J Hum Genet*. 2011 Aug 12;89(2):289-94. doi: 10.1016/j.ajhg.2011.06.007. Epub 2011 Jul 21. Citation on PubMed or Free article on PubMed Central

Walz K, Cohen D, Neilsen PM, Foster J 2nd, Brancati F, Demir K, Fisher R, Moffat M, Verbeek NE, Bjorgo K, Lo Castro A, Curatolo P, Novelli G, Abad C, Lei C, Zhang L, Diaz-Horta O, Young JI, Callen DF, Tekin M. Characterization of ANKRD11 mutations in humans and mice related to KBG syndrome. *Hum Genet*. 2015 Feb;134(2):181-90. doi: 10.1007/s00439-014-1509-2. Epub 2014 Nov 21. Citation on PubMed

Willemsen MH, Fernandez BA, Bacino CA, Gerkes E, de Brouwer AP, Pfundt R, Sikkema-Raddatz B, Scherer SW, Marshall CR, Potocki L, van Bokhoven H, Kleefstra

T. Identification of ANKRD11 and ZNF778 as candidate genes for autism and variable cognitive impairment in the novel 16q24.3 microdeletion syndrome. Eur J Hum Genet. 2010 Apr;18(4):429-35. doi: 10.1038/ejhg.2009.192. Epub 2009 Nov 18. Citation on PubMed or Free article on PubMed Central

Zhang A, Yeung PL, Li CW, Tsai SC, Dinh GK, Wu X, Li H, Chen JD.

Identification of a novel family of ankyrin repeats containing cofactors for p160 nuclear receptor coactivators. J Biol Chem. 2004 Aug 6;279(32):33799-805. doi: 10.1074/jbc.M403997200. Epub 2004 Jun 7. Citation on PubMed

Genomic LocationThe ANKRD11 gene is found on chromosome 16.

Related Health Topics

Genes and Gene Therapy

Genetic Disorders

MEDICAL ENCYCLOPEDIA

Genes

Genetics

Understanding Genetics

What is DNA?

What is a gene?

What is a gene variant and how do variants occur?

Disclaimers

MedlinePlus links to health information from the National Institutes of Health and other federal government agencies. MedlinePlus also links to health information from non-government Web sites. See our disclaimer about external links and our quality guidelines.

The information on this site should not be used as a substitute for professional medical care or advice. Contact a health care provider if you have questions about your health.

Learn how to cite this page

