Single Gene Disorders - Autosomal Dominant Inheritance

https://medlineplus.gov/genetics/condition/familial-mediterranean-fever/

You Are Here:		
Home		
& #8594;		
Canatias		
Genetics		
& #8594;		
Genetic Conditions		
→		
Familial Mediterranean fever		

URL of this page: https://medlineplus.gov/genetics/condition/familial-mediterranean-fever/

Familial Mediterranean fever

To use the sharing features on this page, please enable JavaScript.

Description

Familial Mediterranean fever is an inherited condition characterized by recurrent episodes of painful inflammation in the abdomen, chest, or joints. These episodes are often accompanied by fever and sometimes a rash or headache. Occasionally inflammation may occur in other parts of the body, such as the heart; the membrane surrounding the brain and spinal cord; and in males, the testicles. In about half of affected individuals, attacks are preceded by mild signs and symptoms known as a prodrome. Prodromal symptoms include mildly uncomfortable sensations in the area that will later become inflamed, or more general feelings of discomfort. The first episode of illness in familial Mediterranean fever usually occurs in childhood or the teenage years, but in some cases, the initial attack occurs much later in life. Typically, episodes last 12 to 72 hours and can vary in severity. Episodes generally occur once a month, and in affected women of reproductive age, attacks often correspond with menstruation or ovulation. However, the length of time between episodes can range from days to years. During these periods, affected individuals usually have no signs or symptoms related to the condition. However, without treatment to help prevent attacks and

complications, a buildup of protein deposits (amyloidosis) in the body's organs and tissues may occur, especially in the kidneys, which can lead to kidney failure.

Frequency

Familial Mediterranean fever primarily affects populations originating in the Mediterranean region, particularly people of Armenian, Arab, Turkish, or Jewish ancestry. The disorder affects 1 in 200 to 1,000 people in these populations. It is less common in other populations.

Causes

Familial Mediterranean fever is caused by variants (also known as mutations) in the MEFV gene. The MEFV gene provides instructions for making a protein called pyrin (also known as marenostrin), which is found in white blood cells. This protein is involved in the immune system, helping to regulate the process of inflammation. Inflammation occurs when the immune system sends signaling molecules and white blood cells to a site of injury or disease to fight microbial invaders and facilitate tissue repair. When this process is complete, the body stops the inflammatory response to prevent damage to its own cells and tissues. Variants in the MEFV gene reduce the activity of the pyrin protein, which disrupts control of the inflammation process. An inappropriate or prolonged inflammatory response can result, leading to fever and pain in the abdomen, chest, or joints. Normal variations in the SAA1 gene may modify the course of familial Mediterranean fever. Some evidence suggests that a particular version of the SAA1 gene (called the alpha variant) increases the risk of amyloidosis among people with familial Mediterranean fever.

Learn more about the genes associated with Familial Mediterranean fever

MEFV

SAA1

Inheritance

Familial Mediterranean fever is usually inherited in an autosomal recessive pattern, which means both copies of the MEFV gene in each cell have variants. The parents of an individual with an autosomal recessive condition each carry one copy of the altered gene, but they typically do not show signs and symptoms of the condition. Research shows that about one-third of people with familial Mediterranean fever have a single MEFV gene variant, though most experts believe there is a second variant somewhere else in the gene that has not been detected by genetic testing. In rare cases, this condition appears to be inherited in an autosomal dominant pattern. An autosomal dominant inheritance pattern describes cases in which one copy of the altered gene in each cell is sufficient to cause the disorder. In autosomal dominant inheritance, affected individuals often inherit the variant from one affected parent. However, another mechanism is believed to account for some cases of familial Mediterranean fever that were originally thought to be inherited in an autosomal dominant pattern. A gene variant that occurs frequently in a population may result in a disorder with autosomal recessive inheritance appearing in multiple generations in a family, a pattern that mimics autosomal dominant inheritance. If one parent has familial Mediterranean fever (with variants in both copies of the MEFV gene in each cell) and the other parent is an unaffected carrier (with a variant in one copy of the MEFV gene in each cell), it may appear as if the affected

child inherited the disorder only from the affected parent. This appearance of autosomal dominant
inheritance when the pattern is actually autosomal recessive is called
pseudodominance.
Other Names for This Condition
Benign paroxysmal peritonitis Familial paroxysmal polyserositis FMF MEF Recurrent polyserositis
Reimann periodic disease Siegal-Cattan-Mamou disease Wolff periodic disease
Additional laternation 9 December
Additional Information & Resources
Genetic Testing Information
Genetic Testing Registry: Familial Mediterranean fever
Genetic Testing Registry: Familial mediterranean fever, autosomal dominant

Genetic and Rare Diseases Information Center
Familial Mediterranean fever
Patient Support and Advocacy Resources
National Organization for Rare Disorders (NORD)
Clinical Trials
ClinicalTrials.gov
Catalog of Genes and Diseases from OMIM
FAMILIAL MEDITERRANEAN FEVER, AUTOSOMAL DOMINANT

FAMILIAL MEDITERRANEAN FEVER; FMF
Scientific Articles on PubMed
PubMed
References
Bakkaloglu A. Familial Mediterranean fever. Pediatr Nephrol. 2003
Sep;18(9):853-9. doi: 10.1007/s00467-003-1185-2. Epub 2003 Jun 27. Citation on PubMed
Bhat A, Naguwa SM, Gershwin ME. Genetics and new treatment modalities for
familial Mediterranean fever. Ann N Y Acad Sci. 2007 Sep;1110:201-8. doi:
10.1196/annals.1423.022. Citation on PubMed

Booth DR, Gillmore JD, Lachmann HJ, Booth SE, Bybee A, Soyturk M, Akar S, Pepys MB, Tunca M, Hawkins PN. The genetic basis of autosomal dominant familial Mediterranean fever. QJM. 2000 Apr;93(4):217-21. doi: 10.1093/qjmed/93.4.217. Citation on PubMed

Eisenstein EM, Berkun Y, Ben-Chetrit E. Familial Mediterranean fever: a critical digest of the 2012-2013 literature. Clin Exp Rheumatol. 2013

May-Jun;31(3 Suppl 77):103-7. Epub 2013 Sep 9. Citation on PubMed

Lidar M, Kedem R, Berkun Y, Langevitz P, Livneh A. Familial Mediterranean fever in Ashkenazi Jews: the mild end of the clinical spectrum. J Rheumatol. 2010

Feb;37(2):422-5. doi: 10.3899/jrheum.090401. Epub 2009 Dec 15. Citation on PubMed

Lidar M, Livneh A. Familial Mediterranean fever: clinical, molecular and management advancements. Neth J Med. 2007 Oct;65(9):318-24. Citation on PubMed

Mikula M, Buller A, Sun W, Strom CM. Prevalence of known mutations in the familial Mediterranean fever gene (MEFV) in various carrier screening populations. Genet Med. 2008 May;10(5):349-52. doi: 10.1097/GIM.0b013e3181723cc8.

Citation on PubMed

Rabinovitch E, Harats D, Yaron P, Luvish T, Lidar M, Kedem R, Shaish A, Ben-Dov I, Livneh A. Familial Mediterranean fever gene and protection against asthma. Ann Allergy Asthma Immunol. 2007 Dec;99(6):517-21. doi: 10.1016/S1081-1206(10)60380-8. Citation on PubMed Stoffman N, Magal N, Shohat T, Lotan R, Koman S, Oron A, Danon Y, Halpern GJ, Lifshitz Y, Shohat M. Higher than expected carrier rates for familial Mediterranean fever in various Jewish ethnic groups. Eur J Hum Genet. 2000 Apr;8(4):307-10. doi: 10.1038/sj.ejhg.5200446. Citation on PubMed Yepiskoposyan L, Harutyunyan A. Population genetics of familial Mediterranean fever: a review. Eur J Hum Genet. 2007 Sep;15(9):911-6. doi: 10.1038/sj.ejhg.5201869. Epub 2007 Jun 13. Citation on PubMed Zadeh N, Getzug T, Grody WW. Diagnosis and management of familial Mediterranean fever: integrating medical genetics in a dedicated interdisciplinary clinic. Genet Med. 2011 Mar;13(3):263-9. doi:

10.1097/GIM.0b013e31820e27b1. Citation on PubMed

Related Health Topics

Fever

Genetic Disorders

MEDICAL ENCYCLOPEDIA
Familial Mediterranean fever
Genetics
Secondary systemic amyloidosis
Understanding Constice
Understanding Genetics
What is the prognosis of a genetic condition?
How can gene variants affect health and development?
What does it mean if a disorder seems to run in my family?

What are the different ways a genetic condition can be inherited?

How are genetic conditions treated or managed?

MedlinePlus links to health information from the National Institutes of Health and other federal
government agencies. MedlinePlus also links to health information from non-government Web sites.
See our disclaimer about external links and our quality guidelines.
The information on this site should not be used as a substitute for professional medical care or
advice. Contact a health care provider if you have questions about your health.
Learn how to cite this page