PURA And 5q31

https://medlineplus.gov/genetics/condition/5q313-microdeletion-syndrome/

Home

→

You Are Here:

Genetics

→

Genetic Conditions

&#8594;

5q31.3 microdeletion syndrome

URL of this page: https://medlineplus.gov/genetics/condition/5q313-microdeletion-syndrome/

5q31.3 microdeletion syndrome

To use the sharing features on this page, please enable JavaScript.

Description

5q31.3 microdeletion syndrome is a condition characterized by severely delayed development of speech and motor skills, such as walking. Beginning in infancy, affected individuals also have weak muscle tone (hypotonia), feeding difficulties, and breathing problems. Breathing problems and difficulty swallowing (dysphagia) can be life-threatening.5q31.3 microdeletion syndrome is also characterized by distinctive facial features. Such features include a narrow forehead, widely spaced eyes (hypertelorism), an open mouth with an upper lip that points outward (called a tented lip), a high arch in the roof of the mouth (high-arched palate), a small lower jaw (micrognathia), and a lack of facial expression. Some of these features, such as an open mouth with a tented lip and an expressionless face, are thought to be due to hypotonia.Recurrent seizures (epilepsy) and seizure-like episodes (which can include muscle jerking, twitching, and stiffening), are common in 5q31.3 microdeletion syndrome. Many individuals with 5q31.3 microdeletion syndrome have brain abnormalities, several of which are caused by reduced production of myelin or delayed maturation of myelin. Myelin is the protective covering that insulates nerves and ensures the rapid transmission of


nerve impulses.

Frequency

5q31.3 microdeletion syndrome is a very rare disorder. At least eight individuals with the condition have been described in the medical literature.

Causes

5q31.3 microdeletion syndrome is caused by a chromosomal change in which a small piece of chromosome 5 is deleted in each cell. The deletion occurs on the long (q) arm of the chromosome at a position designated q31.3. The size of the deletion can range from several thousand to several million DNA building blocks (base pairs). The deleted region typically contains at least three genes. The loss of one of these genes, PURA, is thought to lead to most of the characteristic features of the condition. The protein produced from the PURA gene, called Pur-alpha (Purα), has multiple roles in cells, including controlling the activity of genes (gene transcription) and aiding in the copying (replication) of DNA. This protein is especially important for normal brain development; it helps direct the growth and division of nerve cells (neurons) and may be involved in the formation or maturation of myelin. A loss of one copy of the PURA gene is thought to alter normal brain development and impair the function of neurons, leading to developmental delay, hypotonia, seizures, and other neurological problems in people with 5q31.3 microdeletion syndrome. Some studies suggest that loss of another nearby gene increases the severity of the signs and symptoms. It is unclear how the loss of other genes in the deleted region contributes to the development of 5q31.3 microdeletion syndrome.



Severe neonatal hypotonia-seizures-encephalopathy syndrome due to 5q31.3 microdeletion
Additional Information & Resources
Genetic Testing Information
Genetic Testing Registry: PURA-related severe neonatal hypotonia-seizures-encephalopathy
syndrome
Patient Support and Advocacy Resources
National Organization for Rare Disorders (NORD)
Scientific Articles on PubMed

References

Bonaglia MC, Zanotta N, Giorda R, D'Angelo G, Zucca C. Long-term follow-up of a patient with 5q31.3 microdeletion syndrome and the smallest de novo 5q31.2q31.3 deletion involving PURA. Mol Cytogenet. 2015 Nov 14;8:89. doi: 10.1186/s13039-015-0193-9. eCollection 2015. Citation on PubMed or Free article on PubMed Central

Brown N, Burgess T, Forbes R, McGillivray G, Kornberg A, Mandelstam S, Stark Z. 5q31.3 Microdeletion syndrome: clinical and molecular characterization of two further cases. Am J Med Genet A. 2013 Oct;161A(10):2604-8. doi: 10.1002/ajmg.a.36108. Epub 2013 Aug 15. Citation on PubMed Hosoki K, Ohta T, Natsume J, Imai S, Okumura A, Matsui T, Harada N, Bacino CA, Scaglia F, Jones JY, Niikawa N, Saitoh S. Clinical phenotype and candidate genes for the 5q31.3 microdeletion syndrome. Am J Med Genet A. 2012 Aug;158A(8):1891-6. doi: 10.1002/ajmg.a.35439. Epub 2012 Jun 18. Citation on PubMed Shimojima K, Isidor B, Le Caignec C, Kondo A, Sakata S, Ohno K, Yamamoto T. A new microdeletion syndrome of 5q31.3 characterized by severe developmental delays, distinctive facial features, and delayed myelination. Am J Med Genet A. 2011 Apr;155A(4):732-6. doi: 10.1002/ajmg.a.33891. Epub 2011 Mar 15. Erratum In:

Am J Med Genet A. 2011 Nov;155A(11):2903. Citation on PubMed White MK, Johnson EM, Khalili K. Multiple roles for Puralpha in cellular and viral regulation. Cell Cycle. 2009 Feb 1;8(3):1-7. doi: 10.4161/cc.8.3.7585. Epub 2009 Feb 10. Citation on PubMed or Free article on PubMed Central Related Health Topics **Developmental Disabilities Genetic Disorders**

MEDICAL ENCYCLOPEDIA Genetics Hypotonia Intellectual disability **Understanding Genetics** What is the prognosis of a genetic condition? How can gene variants affect health and development? What does it mean if a disorder seems to run in my family? What are the different ways a genetic condition can be inherited? How are genetic conditions treated or managed?

Disclaimers
MedlinePlus links to health information from the National Institutes of Health and other federal
government agencies. MedlinePlus also links to health information from non-government Web sites.
See our disclaimer about external links and our quality guidelines.
The information on this site should not be used as a substitute for professional medical care or
advice. Contact a health care provider if you have questions about your health.
Learn how to cite this page