GRIN2A related syndrome

https://medlineplus.gov/genetics/gene/grin2a/

You Are Here:			
Home			
& #8594;			
Genetics			
& #8594;			
Genes			
& #8594;			
GRIN2A gene			

URL of this page: https://medlineplus.gov/genetics/gene/grin2a/

GRIN2A gene

glutamate ionotropic receptor NMDA type subunit 2A

To use the sharing features on this page, please enable JavaScript.

Normal Function

The GRIN2A gene provides instructions for making a protein called GluN2A (formerly known as NR2A). This protein is found in nerve cells (neurons) in the brain and spinal cord, including regions of the brain involved in speech and language. The GluN2A protein is one component (subunit) of a subset of NMDA receptors. There are several types of NMDA receptors, made up of different combinations of protein components. NMDA receptors are glutamate-gated ion channels; when brain chemicals called glutamate and glycine attach to the receptor, a channel opens, allowing positively charged particles (cations) to flow through. The flow of cations generates currents that activate (excite) neurons to send signals in the brain. NMDA receptors are involved in normal brain development, changes in the brain in response to experience (synaptic plasticity), learning, and memory. They also appear to play a role during deep (slow-wave) sleep. The GluN2A subunit of NMDA receptors determines where in the brain the receptor is located and how it functions. It also provides the site to which glutamate binds.

Health Conditions Related to Genetic Changes

Epilepsy-aphasia spectrum

More than 50 mutations in the GRIN2A gene have been identified in some people with conditions that fall along the epilepsy-aphasia spectrum. This group of conditions is characterized by abnormal electrical activity in the brain, usually during slow-wave sleep; a loss of speech and language skills and sometimes other developmental skills; and in many cases, recurrent seizures (epilepsy). Landau-Kleffner syndrome (LKS) and epileptic encephalopathy with continuous spike-and-wave during sleep (ECSWS) are at the severe end of the spectrum, while childhood epilepsy with centrotemporal spikes (CECTS) is at the mild end. Several other conditions have signs and symptoms of intermediate severity. Many GRIN2A gene mutations lead to production of a nonfunctional GluN2A protein or prevent the production of any protein at all. These mutations likely lead to a reduced number of NMDA receptors containing the GluN2A subunit. Researchers suspect that, as a result, signaling occurs through other types of NMDA receptors that more easily excite neurons, leading to excessive signaling in the brain. Other mutations lead to production of abnormal GluN2A proteins that likely alter how the NMDA receptors function, possibly increasing signaling. Excessive activity of neurons in the brain can lead to seizures and other abnormal brain activity and may result in death of the neurons. \$\%\pi\$4194; \$\%\pi\$160; Changes in GluN2A appear to specifically affect signaling in regions of the brain involved in speech and language and disrupt brain activity during slow-wave sleep, leading to several of the signs and symptoms of this group of conditions. It is not clear why some GRIN2A gene mutations lead to a relatively mild condition and others cause more severe signs and symptoms.

More About This Health Condition

Opioid addiction

MedlinePlus Genetics provides information about Opioid addiction

More About This Health Condition

Other disorders

GRIN2A gene mutations have been found in people with neurological disorders that have features similar to epilepsy-aphasia spectrum disorders (described above) but lacking consistent language problems. These shared features can include recurrent seizures (epilepsy), often of a type that originates from abnormal activity in the rolandic region of the brain (rolandic epilepsy); intellectual disability; and developmental delay. Some people with mutations in chromosome 16 that delete the GRIN2A gene as well as other nearby genes also have unusual facial features. The varying effects of different GRIN2A gene mutations and how they contribute to different neurological disorders are under study.

Other Names for This Gene

EPND GluN2A glutamate receptor ionotropic, NMDA 2A isoform 1 precursor glutamate receptor				
ionotropic, NMDA 2A isoform 2 precursor glutamate receptor, ionotropic, N-methyl D-aspartate 2A				
LKS N-methyl D-aspartate receptor subtype 2A N-methyl-D-aspartate receptor channel, subunit				
epsilon-1 N-methyl-D-aspartate receptor subunit 2A NMDAR2A NR2A				
Additional Information & Resources				
Tests Listed in the Genetic Testing Registry				
Tests of GRIN2A				
Scientific Articles on PubMed				
PubMed				

Catalog of Genes and Diseases from OMIM			
GLUTAMATE RECEPTOR, IONOTROPIC, N-METHYL-D-ASPARTATE, SUBUNIT 2A; GRIN2A			
Gene and Variant Databases			
NCBI Gene			
ClinVar			
References			
Burnashev N, Szepetowski P. NMDA receptor subunit mutations in			
neurodevelopmental disorders. Curr Opin Pharmacol. 2015 Feb;20:73-82. doi:			

neurodevelopmental disorders. Curr Opin Pharmacol. 2015 Feb;20:73-82. doi: 10.1016/j.coph.2014.11.008. Epub 2014 Dec 11. Citation on PubMed Carvill GL, Regan BM, Yendle SC, O'Roak BJ, Lozovaya N, Bruneau N, Burnashev N, Khan A, Cook J, Geraghty E, Sadleir LG, Turner SJ, Tsai MH, Webster R, Ouvrier R, Damiano JA, Berkovic SF, Shendure J, Hildebrand MS, Szepetowski P, Scheffer

IE, Mefford HC. GRIN2A mutations cause epilepsy-aphasia spectrum disorders. Nat Genet. 2013 Sep;45(9):1073-6. doi: 10.1038/ng.2727. Epub 2013 Aug 11. Citation on PubMed or Free article on PubMed Central

Lemke JR, Lal D, Reinthaler EM, Steiner I, Nothnagel M, Alber M, Geider K, Laube B, Schwake M, Finsterwalder K, Franke A, Schilhabel M, Jahn JA, Muhle H, Boor R, Van Paesschen W, Caraballo R, Fejerman N, Weckhuysen S, De Jonghe P, Larsen J, Moller RS, Hjalgrim H, Addis L, Tang S, Hughes E, Pal DK, Veri K, Vaher U, Talvik T, Dimova P, Guerrero Lopez R, Serratosa JM, Linnankivi T, Lehesjoki AE, Ruf S, Wolff M, Buerki S, Wohlrab G, Kroell J, Datta AN, Fiedler B, Kurlemann G, Kluger G, Hahn A, Haberlandt DE, Kutzer C, Sperner J, Becker F, Weber YG, Feucht M, Steinbock H, Neophythou B, Ronen GM, Gruber-Sedlmayr U, Geldner J, Harvey RJ, Hoffmann P, Herms S, Altmuller J, Toliat MR, Thiele H, Nurnberg P, Wilhelm C, Stephani U, Helbig I, Lerche H, Zimprich F, Neubauer BA, Biskup S, von Spiczak S. Mutations in GRIN2A cause idiopathic focal epilepsy with rolandic spikes. Nat Genet. 2013 Sep;45(9):1067-72. doi: 10.1038/ng.2728. Epub 2013 Aug 11. Citation on PubMed

Lesca G, Rudolf G, Bruneau N, Lozovaya N, Labalme A, Boutry-Kryza N, Salmi M, Tsintsadze T, Addis L, Motte J, Wright S, Tsintsadze V, Michel A, Doummar D, Lascelles K, Strug L, Waters P, de Bellescize J, Vrielynck P, de Saint Martin A, Ville D, Ryvlin P, Arzimanoglou A, Hirsch E, Vincent A, Pal D, Burnashev N, Sanlaville D, Szepetowski P. GRIN2A mutations in acquired epileptic aphasia and related childhood focal epilepsies and encephalopathies with speech and language dysfunction. Nat Genet. 2013 Sep;45(9):1061-6. doi: 10.1038/ng.2726. Epub 2013 Aug 11. Citation on PubMed

Paoletti P. Molecular basis of NMDA receptor functional diversity. Eur J
Neurosci. 2011 Apr;33(8):1351-65. doi: 10.1111/j.1460-9568.2011.07628.x. Epub
2011 Mar 14. Citation on PubMed

Reutlinger C, Helbig I, Gawelczyk B, Subero JI, Tonnies H, Muhle H,

Finsterwalder K, Vermeer S, Pfundt R, Sperner J, Stefanova I, Gillessen-Kaesbach

G, von Spiczak S, van Baalen A, Boor R, Siebert R, Stephani U, Caliebe A.

Deletions in 16p13 including GRIN2A in patients with intellectual disability,

various dysmorphic features, and seizure disorders of the rolandic region.

Epilepsia. 2010 Sep;51(9):1870-3. doi: 10.1111/j.1528-1167.2010.02555.x. Citation on PubMed

Turner SJ, Mayes AK, Verhoeven A, Mandelstam SA, Morgan AT, Scheffer IE.

GRIN2A: an aptly named gene for speech dysfunction. Neurology. 2015 Feb

10;84(6):586-93. doi: 10.1212/WNL.000000000001228. Epub 2015 Jan 16. Citation on PubMed or

Free article on PubMed Central


Wyllie DJ, Livesey MR, Hardingham GE. Influence of GluN2 subunit identity on

NMDA receptor function. Neuropharmacology. 2013 Nov;74:4-17. doi:

10.1016/j.neuropharm.2013.01.016. Epub 2013 Jan 31. Citation on PubMed or Free article on

PubMed Central

Genomic LocationThe GRIN2A gene is found on chromosome 16.
Related Health Topics
Genes and Gene Therapy
Genetic Disorders
MEDICAL ENCYCLOPEDIA
Genes
Genetics



Learn how to cite this page