

Assignment 6 : Papoulis Textbook

Akshitha Kola

June 3, 2022

Outline

1 Question

2 Solution

Question

Chapter 3 example 3.17

A and B plays a series of games where the probability of winning p in a single play for A is unfairly kept at less than $\frac{1}{2}$. However, A gets to choose in advance the total number of plays. To win the whole game one must score more than half the plays. If the total number of plays is to be even, how many plays should A choose?

Solution

As the total number of plays to be played is even, let the total number of plays be $2n$. Let X_k be the number of plays A wins out of $2n$ plays. Then

$$P(X_k) = \binom{2n}{k} p^k q^{2n-k}$$

Let P_{2n} denote the event that A wins in $2n$ plays.

$$P_{2n} = P\left(\bigcup_{k=n+1}^{2n} X_k\right) \quad (1)$$

$$= \sum_{k=n+1}^{2n} P(X_k) \quad (2)$$

$$= \sum_{k=n+1}^{2n} \binom{2n}{k} p^k q^{2n-k} \quad (3)$$

As the required number of plays is $2n$ $P_{2n-2} \leq P_{2n} \leq P_{2n+2}$

$$P_{2n+2} = \sum_{k=n+2}^{2n+2} \binom{2n+2}{k} p^k q^{2n+2-k} = (p+q)^{2n+2} = (p+q)^{2n}(p+q)^2 \quad (4)$$

$$= \left\{ \sum_{k=n+1}^{2n} \binom{2n}{k} p^k q^{2n-k} \right\} (p^2 + 2pq + q^2) \quad (5)$$

$$(6)$$

From the above equation we get a identity after equating the powers of p and q.

$$P_{2n+2} = P_{2n} + \binom{2n}{n} p^{n+2} q^n - \binom{2n}{n+1} p^{n+1} q^{n+1} \quad (7)$$

If $2n$ is optimum, then from the inequality we get

$$\binom{2n}{n+1} p^{n+1} q^{n+1} \leq \binom{2n}{n} p^{n+2} q^n \quad (8)$$

$$\implies nq \leq (n+1)p \implies n(q-p) \leq p \implies n \leq \frac{p}{1-2p} \quad (9)$$

and

$$\binom{2n-2}{n-1} p^{n+1} q^{n-1} \leq \binom{2n-2}{n} p^n q^n \quad (10)$$

$$\implies np \leq (n-1)q \implies n(q-p) \geq q \implies n \geq \frac{q}{1-2p} \quad (11)$$

From (9) and (11) we get

$$\frac{1}{1-2p} - 1 \leq 2n \leq \frac{1}{1-2p} + 1 \quad (12)$$

\therefore This means $2n$ is the even number nearest to $\frac{1}{1-2p}$