

Issues in assessing reliability in pain neuroimaging

Tor D. Wager^{1,2*}, Choong-Wan Woo^{1,2},

¹Department of Psychology and Neuroscience, University of Colorado, Boulder

²Institute of Cognitive Science, University of Colorado, Boulder

* Please address correspondence to:

Tor D. Wager

Department of Psychology and Neuroscience

University of Colorado Boulder

345 UCB, Boulder, CO 80305

Email: Tor.Wager@Colorado.Edu

Telephone: 303-492-7487

Letzen et al. [5] have thoughtfully responded to our comment on the limitations of reliability measures in brain imaging and beyond. We find much to agree with, and a few points that require additional clarification. We agree that inter-individual reliability is a cornerstone of measurement theory, and there has been too little emphasis in neuroimaging studies on measurement properties.

We also agree that there are many valid and reliable pain-related assessments that go far beyond intensity ratings, including measures associated with diverse aspects and consequences of pain [3,7,10]. These assessments are valuable, and, neuroimaging studies would do well to identify predictive signatures for biological component processes underlying these diverse aspects of pain.

Despite these points of agreement, we would like to emphasize several additional points related to reliability and neuroimaging. First, it does not make sense to talk about the reliability of neuroimaging as a whole, any more than it does to talk about the reliability of behavior as a whole. Brain imaging provides myriad measures, some reliable, others not. Most are irrelevant for pain. When assessing the reliability of brain measures related to pain, we must carefully select measures that are maximally pain-related. This is true even when considering *a priori* regions of interest. For example, the anterior mid-cingulate cortex (aMCC) is pain-related in general, but it contains on the order of 550 million neurons, and there are literally hundreds of ways to define a summary measure of activity across the region as a whole. Some will yield signals that are pain-related, and others signals that are completely unrelated [13]. Thus, before assessing reliability, we should define measures that are as closely related to the construct of interest as possible. That is why it is particularly relevant to assess the reliability of brain signatures such as the Neurologic Pain Signature (NPS) [12]—which appears to be approximately as reliable as pain self-reports [15].

Second, most pain neuroimaging studies have not focused on individual differences. Reliability places a cap on the validity of assessing inter-individual variability—for example who feels more or less pain—but not on the validity of measures assessing why pain exists at all, or what brain and behavioral processes might be driving it. Most pain imaging

studies have been designed to identify brain processes that are consistent across individuals, and so have striven to maximize homogeneity. The net result, as Letzen et al. [5] point out, is reduced reliability—but these studies are still valuable for their contribution to understanding the nature of pain processing. Neuroimaging studies should examine both consistent effects and inter-individual variability in more diverse community and patient samples.

Third, we disagree with Letzen et al. that “If neuroimaging markers are truly representative of the symptom in question [pain], then they should be similarly influenced by the same factors that influence self-report.” As Letzen et al. [5] point out, pain is a multidimensional experience [3,8], and diverse aspects of functionality and wellbeing are also important [10]. In addition, people are diverse in the ways they reflect on and describe pain. Thus, it is unlikely that one brain marker can, even in principle, capture pain reports in all contexts. It is more likely that pain is supported by a family of brain states, and shades and hues in pain are emergent properties that arise from a family of basic neurophysiological ‘ingredient’ processes (cf. [6]). It is also unlikely that one brain marker will ever be able to capture effects of every treatment or variable that affects pain report—as our earlier work has shown [12,14]. Some “factors that influence self-report” influence nociception, whereas others affect emotion, decision-making about experience, or social self-presentation.

Furthermore, neuroimaging and behavioral measures are likely to have *complementary* sources of bias and noise. All kinds of decisions (not just those about pain) are subject to cognitive biases, including anchor-and-adjust decision biases [11], self-consistency biases [9], social conformity effects [16], and others. Some pain-related neuroimaging measures (including the NPS [12]) may be less subject to these forms of decision bias. They are, however, subject to their own sources of error, including sensitivity to factors such as caffeine intake [4], vascular aging [2], and ‘off-target’ activity that is not pain-related. To the extent that they have complementary sources of error but load on the same construct (“pain”), measuring brain responses and behavior will be superior to either alone [1]. Thus, using converging measures to ‘triangulate’ on pain and its consequences may prove advantageous.

In sum, rather than looking for one brain measure that tracks all aspects of self-reported pain in all contexts, neuroimaging measures can be acknowledged for what they are: A source of information about the *component neurophysiological ingredients* that give rise to pain. These *component processes* may combine in various ways to contribute to what is reported as pain, as letters combine to form words (cf. [17]). The presence or absence of individual neurophysiological components may prove to be informative about the *nature* of a person's pain, beyond its mere presence or intensity. They may have implications for health independent of a person's subjective pain experience, and may ultimately be considered as endpoints in their own right.

Acknowledgements

This work was funded by NIDA R01DA035484-01 (T.D.W., principal investigator).

References

- [1] Campbell DT, Fiske DW. Convergent and discriminant validation by the multitrait-multimethod matrix. Psychol. Bull. 1959;56:81–105.
- [2] D'Esposito M, Deouell LY, Gazzaley A. Alterations in the BOLD fMRI signal with ageing and disease: a challenge for neuroimaging. Nat. Rev. Neurosci. 2003;4:863–872.
- [3] Knotkova H, Clark WC, Keohan ML, Kuhl JP, Winer RT, Wharton RN. Validation of the Multidimensional Affect and Pain Survey (MAPS). J. Pain Off. J. Am. Pain Soc. 2006;7:161–169.
- [4] Laurienti PJ, Field AS, Burdette JH, Maldjian JA, Yen Y-F, Moody DM. Dietary Caffeine Consumption Modulates fMRI Measures. NeuroImage 2002;17:751–757.
- [5] Letzen JE, Boissoneault J, Sevel LS, Robinson ME. Reply to: What reliability can and cannot tell us about pain report and pain neuroimaging. PAIN n.d.
- [6] Lindquist K a, Wager TD, Kober H, Bliss-Moreau E, Barrett LF. The brain basis of emotion: a meta-analytic review. Behav. Brain Sci. 2012;35:121–43.

- [7] Melzack R. The McGill Pain Questionnaire: major properties and scoring methods. *Pain* 1975;1:277–299.
- [8] Melzack R, Casey KL. Sensory, motivational, and central control determinants of pain: a new conceptual model. *The Skin Senses*. 1968.
- [9] Scharfe E, Bartholomew K. Do you remember?: Recollections of adult attachment patterns. *Pers. Relatsh.* 1998;5:219–234.
- [10] Turk DC, Dworkin RH, Revicki D, Harding G, Burke LB, Cella D, Cleeland CS, Cowan P, Farrar JT, Hertz S, Max MB, Rappaport BA. Identifying important outcome domains for chronic pain clinical trials: an IMMPACT survey of people with pain. *Pain* 2008;137:276–285.
- [11] Tversky a, Kahneman D. Judgment under Uncertainty: Heuristics and Biases. *Science* 1974;185:1124–31.
- [12] Wager TD, Atlas LY, Lindquist M a, Roy M, Woo C-W, Kross E. An fMRI-based neurologic signature of physical pain. *N. Engl. J. Med.* 2013;368:1388–97.
- [13] Woo C-W, Koban L, Kross E, Lindquist M a, Banich MT, Ruzic L, Andrews-Hanna JR, Wager TD. Separate neural representations for physical pain and social rejection. *Nat. Commun.* 2014;5:5380.
- [14] Woo C-W, Roy M, Buhle JT, Wager TD. Distinct Brain Systems Mediate the Effects of Nociceptive Input and Self-Regulation on Pain. *PLoS Biol.* 2015;13:e1002036.
- [15] Woo C-W, Wager TD. What reliability can and cannot tell us about pain report and pain neuroimaging. *Pain* 2016;157:511–513.
- [16] Wood W. Attitude Change: Persuasion and Social Influence. *Annu. Rev. Psychol.* 2000;51:539–570.
- [17] Zaki J, Wager TD, Singer T, Keysers C, Gazzola V. The anatomy of suffering: Understanding the relationship between nociceptive and empathic pain. *Trends Cogn. Sci.* n.d.