

We induct over the structure of typing derivations of $\Gamma \vdash t : \tau$, assuming $\omega : \Gamma' \triangleright \Gamma$ holds. In each case, we construct the new derivation Δ' from the derivation Δ giving $\Gamma \vdash t : \tau$ and show that $\Delta \circ \llbracket \omega : \Gamma' \triangleright \Gamma \rrbracket_M = \Delta'$

0.0.1 Variable Terms

Case Var and Weaken We case split on the weakening ω .

If $\omega = \iota$ Then $\Gamma' = \Gamma$, and so $\Gamma' \vdash x : A$ holds and the derivation Δ' is the same as Δ

$$\Delta' = \Delta = \Delta \circ \text{Id}_\Gamma = \Delta \circ \llbracket \iota : \Gamma \triangleright \Gamma \rrbracket_M \quad (1)$$

If $\omega = \omega' \pi$ Then $\Gamma' = (\Gamma'', x' : A')$ and $\omega' : \Gamma'' \triangleright \Gamma$. So by induction, there is a tree, Δ_1 deriving $\Gamma'' \vdash x : A$, such that

$$\Delta_1 = \Delta \circ \llbracket \omega' : \Gamma'' \triangleright \Gamma \rrbracket_M \quad \text{By Induction} \quad (2)$$

, and hence by the weaken rule, we have

$$(\text{Weaken}) \frac{\Gamma'' \vdash x : A}{\Gamma'', x' : A' \vdash x : A} \quad (3)$$

This preserves denotations:

$$\Delta' = \Delta_1 \circ \pi_1 \quad \text{By Definition} \quad (4)$$

$$= \Delta \circ \llbracket \omega' : \Gamma'' \triangleright \Gamma \rrbracket_M \circ \pi_1 \quad \text{By induction} \quad (5)$$

$$= \Delta \circ \llbracket \omega' \pi_1 : \Gamma' \triangleright \Gamma \rrbracket_M \quad \text{By denotation of weakening} \quad (6)$$

If $\omega = \omega' \times$ Then

$$\Gamma' = \Gamma''', x' : B \quad (7)$$

$$\Gamma = \Gamma'', x' : A' \quad (8)$$

$$B \leq A \quad (9)$$

If $x = x'$ Then $A = A'$.

Then we derive the new derivation, Δ' as so:

$$(\text{Sub-type}) \frac{(\text{var}) \frac{\Gamma''', x : B \vdash x : B}{B \leq A}}{\Gamma' \vdash x : A} \quad (10)$$

This preserves denotations:

$$\Delta' = \llbracket B \leq A \rrbracket_M \circ \pi_2 \quad \text{By Definition} \quad (11)$$

$$= \pi_2 \circ (\llbracket \omega' : \Gamma''' \triangleright \Gamma'' \rrbracket_M \times \llbracket B \leq A \rrbracket_M) \quad \text{By the properties of binary products} \quad (12)$$

$$= \Delta \circ \llbracket \omega : \Gamma' \triangleright \Gamma \rrbracket_M \quad \text{By Definition} \quad (13)$$

Case $x \neq x'$ Then

$$\Delta = (\text{Weaken}) \frac{() \frac{\Delta_1}{\Gamma'' \vdash x:A}}{\Gamma \vdash x:A} \quad (14)$$

By induction with $\omega : \Gamma''' \triangleright \Gamma''$, we have a derivation Δ_1 of $\Gamma''' \vdash x:A$

We have the weakened derivation:

$$\Delta' = (\text{Weaken}) \frac{() \frac{\Delta'_1}{\Gamma''' \vdash x:A}}{\Gamma' \vdash x:A} \quad (15)$$

This preserves denotations:

By induction, we have

$$\Delta'_1 = \Delta_1 \circ \llbracket \omega : \Gamma''' \triangleright \Gamma'' \rrbracket_M \quad (16)$$

So we have:

$$\Delta' = \Delta'_1 \circ \pi_1 \quad \text{By denotation definition} \quad (17)$$

$$= \Delta_1 \circ \llbracket \omega' : \Gamma''' \triangleright \Gamma'' \rrbracket_M \quad \text{By induction} \circ \pi_1 \quad (18)$$

$$= \Delta_1 \circ \pi_1 \circ (\llbracket \omega' : \Gamma''' \triangleright \Gamma'' \rrbracket_M \times \llbracket A' \leq B \rrbracket_M) \quad \text{By product properties} \quad (19)$$

$$= \Delta \circ \llbracket \omega : \Gamma' \triangleright \Gamma \rrbracket_M \quad \text{By definition} \quad (20)$$

0.0.2 Value Terms

From this point onwards, since we no-longer case split over the weakening relations, we write the denotation $\llbracket \omega : \Gamma' \triangleright \Gamma \rrbracket_M$, simply as ω .

Case Constant The constant typing rules, $()$, **true**, **false**, \mathcal{C}^A , all proceed by the same logic. Hence I shall only prove the theorems for the case \mathcal{C}^A .

$$(\text{Const}) \frac{\Gamma 0k}{\Gamma \vdash \mathcal{C}^A:A} \quad (21)$$

By inversion, we have $\Gamma 0k$, so we have $\Gamma' 0k$.

Hence

$$(\text{Const}) \frac{\Gamma' 0k}{\Gamma' \vdash \mathcal{C}^A:A} \quad (22)$$

Holds.

This preserves denotations:

$$\Delta' = \llbracket \mathcal{C}^A \rrbracket_M \circ \langle \rangle_{\Gamma'} \quad \text{By definition} \quad (23)$$

$$= \llbracket \mathcal{C}^A \rrbracket_M \circ \langle \rangle_{\Gamma} \circ \omega \quad \text{By the terminal property} \quad (24)$$

$$= \Delta \quad \text{By Definition} \quad (25)$$

$$(26)$$

Case Lambda By inversion, we have a derivation Δ_1 giving

$$\Delta = (\text{Fn}) \frac{() \frac{\Delta_1}{\Gamma, x:A \vdash C : \mathbb{M}_\epsilon B}}{\Gamma \vdash \lambda x : A. C : A \rightarrow \mathbb{M}_\epsilon B} \quad (27)$$

Since $\omega : \Gamma' \triangleright \Gamma$, we have:

$$\omega \times : (\Gamma, x : A) \triangleright (\Gamma, x : A) \quad (28)$$

Hence, by induction, using $\omega \times : (\Gamma, x : A) \triangleright (\Gamma, x : A)$, we derive Δ'_1 :

$$\Delta' = (\text{Fn}) \frac{() \frac{\Delta'_1}{\Gamma', x:A \vdash C : \mathbb{M}_\epsilon B}}{\Gamma', x : A \vdash \lambda x : A. C : A \rightarrow \mathbb{M}_\epsilon B} \quad (29)$$

This preserves denotations:

$$\Delta' = \text{cur}(\Delta'_1) \quad \text{By Definition} \quad (30)$$

$$= \text{cur}(\Delta_1 \circ (\omega \times \text{Id}_\Gamma)) \quad \text{By the denotation of } \omega \times \quad (31)$$

$$= \text{cur}(\Delta_1) \circ \omega \quad \text{By the exponential property} \quad (32)$$

$$= \Delta \circ \omega \quad \text{By Definition} \quad (33)$$

Case Sub-typing

$$(\text{Sub-type}) \frac{\Gamma \vdash v : A \quad A \leq B}{\Gamma \vdash v : B} \quad (34)$$

by inversion, we have a derivation Δ_1

$$() \frac{\Delta_1}{\Gamma \vdash v : A} \quad (35)$$

So by induction, we have a derivation Δ'_1 such that:

$$(\text{Sub-type}) \frac{() \frac{\Delta'_1}{\Gamma' \vdash v : A} \quad A \leq B}{\Gamma' \vdash v : B} \quad (36)$$

This preserves denotations:

$$\Delta' = \llbracket A \leq B \rrbracket_M \circ \Delta'_1 \quad \text{By Definition} \quad (37)$$

$$= \llbracket A \leq B \rrbracket_M \circ \Delta_1 \circ \omega \quad \text{By induction} \quad (38)$$

$$= \Delta \circ \omega \quad \text{By Definition} \quad (39)$$

$$(40)$$

0.0.3 Computation Terms

Case Return We have the sub-derivation Δ_1 such that

$$\Delta = (\text{Return}) \frac{() \frac{\Delta_1}{\Gamma \vdash v : A}}{\Gamma \vdash \text{return } v : \mathbb{M}_1 A} \quad (41)$$

Hence, by induction, with $\omega : \Gamma' \triangleright \Gamma$, we find the derivation Δ'_1 such that:

$$\Delta' = (\text{Return}) \frac{() \frac{\Delta'_1}{\Gamma' \vdash v:A}}{\Gamma' \vdash \text{return } v : \mathbb{M}_1 A} \quad (42)$$

This preserves denotations:

$$\Delta' = \eta_A \circ \Delta'_1 \quad \text{By definition} \quad (43)$$

$$= \eta_A \circ \Delta_1 \circ \omega \quad \text{By induction of } \Delta_1, \Delta'_1 \quad (44)$$

$$= \Delta \circ \omega \quad \text{By Definition} \quad (45)$$

Case Apply By inversion, we have derivations Δ_1, Δ_2 such that

$$\Delta = (\text{Apply}) \frac{() \frac{\Delta_1}{\Gamma \vdash v_1:A \rightarrow \mathbb{M}_\epsilon B} \quad () \frac{\Delta_2}{\Gamma \vdash v_2:A}}{\Gamma \vdash v_1 \ v_2 : \mathbb{M}_\epsilon B} \quad (46)$$

By induction, this gives us the respective derivations: Δ'_1, Δ'_2 such that

$$\Delta' = (\text{Apply}) \frac{() \frac{\Delta'_1}{\Gamma' \vdash v_1:A \rightarrow \mathbb{M}_\epsilon B} \quad () \frac{\Delta'_2}{\Gamma' \vdash v_2:A}}{\Gamma' \vdash v_1 \ v_2 : \mathbb{M}_\epsilon B} \quad (47)$$

This preserves denotations:

$$\Delta' = \text{app} \circ \langle \Delta'_1, \Delta'_2 \rangle \quad \text{By Definition} \quad (48)$$

$$= \text{app} \circ \langle \Delta_1 \circ \omega, \Delta_2 \circ \omega \rangle \quad \text{By induction on } \Delta_1, \Delta_2 \quad (49)$$

$$= \text{app} \circ \langle \Delta_1, \Delta_2 \rangle \circ \omega \quad (50)$$

$$= \Delta \circ \omega \quad \text{By Definition} \quad (51)$$

Case If By inversion, we have the sub-derivations $\Delta_1, \Delta_2, \Delta_3$, such that:

$$\Delta = (\text{If}) \frac{() \frac{\Delta_1}{\Gamma \vdash v:\text{Bool}} \quad () \frac{\Delta_2}{\Gamma \vdash C_1:\mathbb{M}_\epsilon A} \quad () \frac{\Delta_3}{\Gamma \vdash C_2:\mathbb{M}_\epsilon A}}{\Gamma \vdash \text{if}_{\epsilon,A} v \text{ then } C_1 \text{ else } C_2 : \mathbb{M}_\epsilon A} \quad (52)$$

By induction, this gives us the sub-derivations $\Delta'_1, \Delta'_2, \Delta'_3$ such that

$$\Delta' = (\text{If}) \frac{() \frac{\Delta'_1}{\Gamma' \vdash v:\text{Bool}} \quad () \frac{\Delta'_2}{\Gamma' \vdash C_1:\mathbb{M}_\epsilon A} \quad () \frac{\Delta'_3}{\Gamma' \vdash C_2:\mathbb{M}_\epsilon A}}{\Gamma' \vdash \text{if}_{\epsilon,A} v \text{ then } C_1 \text{ else } C_2 : \mathbb{M}_\epsilon A} \quad (53)$$

And

$$\Delta'_1 = \Delta_1 \circ \omega \quad (54)$$

$$\Delta'_3 = \Delta_2 \circ \omega \quad (55)$$

$$\Delta'_3 = \Delta_3 \circ \omega \quad (56)$$

This preserves denotations. Since $\omega : \Gamma' \rightarrow \Gamma$,
Let $(T_\epsilon A)^\omega : T_\epsilon A^\Gamma \rightarrow T_\epsilon A^{\Gamma'}$ be as defined in ExSh 3 ⁽¹⁾ That is:

¹<https://www.cl.cam.ac.uk/teaching/1819/L108/exercises/L108-exercise-sheet-3.pdf>

$$(T_\epsilon A)^\omega = \text{cur}(\text{app} \circ (\text{Id}_{T_\epsilon A} \times w)) \quad (57)$$

. And hence, we have:

$$\text{cur}(f \circ (\text{Id} \times \omega)) = (T_\epsilon A)^\omega \circ \text{cur}(f) \quad (58)$$

$$\Delta' = \text{app} \circ (([\text{cur}(\Delta'_2 \circ \pi_2), \text{cur}(\Delta'_3 \circ \pi_2)] \circ \Delta'_1) \times \text{Id}_{\Gamma'}) \circ \delta_{\Gamma'} \quad \text{By Definition} \quad (59)$$

$$= \text{app} \circ (([\text{cur}(\Delta_2 \circ \omega \circ \pi_2), \text{cur}(\Delta_3 \circ \omega \circ \pi_2)] \circ \Delta'_1) \times \text{Id}_{\Gamma'}) \circ \delta_{\Gamma'} \quad \text{By Induction} \quad (60)$$

$$= \text{app} \circ (([\text{cur}(\Delta_2 \circ \pi_2 \circ (\text{Id}_1 \times \omega)), \text{cur}(\Delta_3 \circ \pi_2 \circ (\text{Id}_1 \times \omega))] \circ \Delta_1 \circ \omega) \times \text{Id}_{\Gamma'}) \circ \delta_{\Gamma'} \quad \text{By product property} \quad (61)$$

$$= \text{app} \circ (((T_\epsilon A)^\omega \circ \text{cur}(\Delta_2 \circ \pi_2), (T_\epsilon A)^\omega \circ \text{cur}(\Delta_3 \circ \pi_2)) \circ \Delta_1 \circ \omega) \times \text{Id}_{\Gamma'}) \circ \delta_{\Gamma'} \quad \text{By } (T_\epsilon A)^\omega \text{ property} \quad (62)$$

$$= \text{app} \circ (((T_\epsilon A)^\omega \circ [\text{cur}(\Delta_2 \circ \pi_2), \text{cur}(\Delta_3 \circ \pi_2)] \circ \Delta_1 \circ \omega) \times \text{Id}_{\Gamma'}) \circ \delta_{\Gamma'} \quad \text{Factor out transformation} \quad (63)$$

$$= \text{app} \circ ((T_\epsilon A)^\omega \times \text{Id}_{\Gamma'}) \circ (([\text{cur}(\Delta_2 \circ \pi_2), \text{cur}(\Delta_3 \circ \pi_2)] \circ \Delta_1) \times \text{Id}_{\Gamma'}) \circ (\omega \times \text{Id}_{\Gamma'}) \circ \delta_{\Gamma'} \quad \text{Factor out Identity pairs} \quad (64)$$

$$= \text{app} \circ (\text{Id}_{(T_\epsilon A)} \times \omega) \circ (([\text{cur}(\Delta_2 \circ \pi_2), \text{cur}(\Delta_3 \circ \pi_2)] \circ \Delta_1) \times \text{Id}_{\Gamma'}) \circ (\omega \times \text{Id}_{\Gamma'}) \circ \delta_{\Gamma'} \quad \text{By defintion of app, } (T_\epsilon A)^\omega \quad (65)$$

$$= \text{app} \circ (([\text{cur}(\Delta_2 \circ \pi_2), \text{cur}(\Delta_3 \circ \pi_2)] \circ \Delta_1) \times \text{Id}_{\Gamma'}) \circ (\omega \times \omega) \circ \delta_{\Gamma'} \quad \text{Push through pairs} \quad (66)$$

$$= \text{app} \circ (([\text{cur}(\Delta_2 \circ \pi_2), \text{cur}(\Delta_3 \circ \pi_2)] \circ \Delta_1) \times \text{Id}_{\Gamma'}) \circ \delta_{\Gamma} \circ \omega \quad \text{By Definition of the diagonal morphism.} \quad (67)$$

$$= \Delta \circ \omega \quad (68)$$

Case Bind By inversion, we have derivations Δ_1, Δ_2 such that:

$$\Delta = (\text{Bind}) \frac{() \frac{\Delta_1}{\Gamma \vdash C_1 : \mathbb{M}_{\mathbb{E}_1} A} \quad () \frac{\Delta_2}{\Gamma, x : A \vdash C_2 : \mathbb{M}_{\epsilon_2} B}}{\Gamma \vdash \text{do } x \leftarrow C_1 \text{ in } C_2 : \mathbb{M}_{\epsilon_1 \cdot \epsilon_2} B} \quad (69)$$

If $\omega : \Gamma' \triangleright \Gamma$ then $\omega \times : \Gamma', x : A \triangleright \Gamma, x : A$, so by induction, we can derive Δ'_1, Δ'_2 such that:

$$\Delta' = (\text{Bind}) \frac{() \frac{\Delta'_1}{\Gamma' \vdash C_1 : \mathbb{M}_{\mathbb{E}_1} A} \quad () \frac{\Delta'_2}{\Gamma', x : A \vdash C_2 : \mathbb{M}_{\epsilon_2} B}}{\Gamma' \vdash \text{do } x \leftarrow C_1 \text{ in } C_2 : \mathbb{M}_{\epsilon_1 \cdot \epsilon_2} B} \quad (70)$$

This preserves denotations:

$$\Delta' = \mu_{\epsilon_1, \epsilon_2, B} \circ T_{\epsilon_1} \Delta'_2 \circ \mathbf{t}_{\epsilon_1, \Gamma', A} \circ \langle \text{Id}_{\Gamma'}, \Delta'_1 \rangle \quad \text{By definition} \quad (71)$$

$$= \mu_{\epsilon_1, \epsilon_2, B} \circ T_{\epsilon_1} (\Delta_2 \circ (\omega \times \text{Id}_A)) \circ \mathbf{t}_{\epsilon_1, \Gamma', A} \circ \langle \text{Id}_{\Gamma'}, \Delta_1 \circ \omega \rangle \quad \text{By induction on } \Delta'_1, \Delta'_2 \quad (72)$$

$$= \mu_{\epsilon_1, \epsilon_2, B} \circ T_{\epsilon_1} \Delta_2 \circ \mathbf{t}_{\epsilon_1, \Gamma, A} \circ \langle \omega, \Delta_1 \circ \omega \rangle \quad \text{By tensor strength} \quad (73)$$

$$= \mu_{\epsilon_1, \epsilon_2, B} \circ T_{\epsilon_1} \Delta_2 \circ \mathbf{t}_{\epsilon_1, \Gamma, A} \circ \langle \text{Id}_{\Gamma}, \Delta_1 \rangle \circ \omega \quad \text{By product property} \quad (74)$$

$$= \Delta \quad \text{By definition} \quad (75)$$

Case Sub-effect

$$\text{(Sub-effect)} \frac{\Gamma \vdash C : \mathbb{M}_{\epsilon_1} A \quad A \leq B \quad \epsilon_1 \leq \epsilon_2}{\Gamma \vdash C : \mathbb{M}_{\epsilon_2} B} \quad (76)$$

by inversion, we have a derivation Δ_1

$$() \frac{\Delta_1}{\Gamma \vdash C : \mathbb{M}_{\epsilon_1} A} \quad (77)$$

So by induction, we have a derivation Δ'_1 such that:

$$\text{(Sub-effect)} \frac{() \frac{\Delta'_1}{\Gamma' \vdash C : \mathbb{M}_{\epsilon_1} A} \quad A \leq B \quad \epsilon_1 \leq \epsilon_2}{\Gamma' \vdash C : \mathbb{M}_{\epsilon_2} B} \quad (78)$$

This preserves denotations:

Let

$$g = \llbracket A \leq B \rrbracket_M : A \rightarrow B \quad (79)$$

$$h = \llbracket \epsilon_1 \leq \epsilon_2 \rrbracket_M : T_{\epsilon_1} \rightarrow T_{\epsilon_2} \quad (80)$$

Then

$$\Delta' = h_B \circ T_{\epsilon_1} g \circ \Delta'_1 \quad \text{By Definition} \quad (81)$$

$$= h_B \circ T_{\epsilon_1} g \circ \Delta_1 \circ \omega \quad \text{By Induction} \quad (82)$$

$$= \Delta \circ \omega \quad \text{By Definition} \quad (83)$$