



Monoclonal antibody ; Rituximab

Presented by : Salma Mohamed



Agenda

01

Mechanism of action

05

Emetic potential

02

Therapeutic application

06

Risk points & managing toxicity

03

Preparation & Administration

07

Drug-Drug interactions

04

Administration guidelines

08

Cautions & Warning

1-MOA

Rituximab is an anti **CD20** chimeric antibody

- CD20 is a surface **transmembrane protein marker** expressed on **B cells** during differentiation from the pre-B cell until the plasma cell stage.
- CD20 is believed to function as a **calcium channel** and play **a role in the maturation and activation of B cells**
- ❑ Once rituximab is bound to CD20 positive cells, **cell death** is induced by various mechanisms, including :
 - antibody-dependent cell-mediated cytotoxicity (ADCC),
 - complement-mediated-cytotoxicity(CDC)
 - antibody-dependent phagocytosis (ADP),
 - direct effects of binding of rituximab to CD20.



Rituximab combination

Combination of rituximab and hyaluronidase

Rituximab-hyaluronidase human is used in **subcutaneous form**.

i.e. Hyaluronidase causes a reversible increase in permeability of subcutaneous tissue by causing depolymerization of hyaluronan,

leading to an **increase in the absorption rate of rituximab**.



2-Therapeutic applications

P e d i a t r i c s



1- Leukemia, acute B-cell; CD20+; advanced stage, previously untreated:

❑ **Infants ≥ 6 months, Children, and Adolescents:**

- IV infusion: **375 mg/m²/dose** in combination with systemic (LMB) chemotherapy regimen

2- Acute lymphoblastic leukemia, mature B-cell, CD 20+ (B-ALL); relapsed/refractory:

❑ **≥ 5 years and Adolescents:**

- IV infusion: **375 mg/m²** administered **on days 1 and 3** of **courses 1 & 2**
- **On day 1 only** of **course 3** ; used in combination with ifosfamide, carboplatin, and etoposide (ICE)

Lymphome Malin B (LMB)

LMB 2001 was a FAB (**French-American-British**) /LMB 96-based protocol.

❑ Treatment was based on **St Jude's/Murphy's stage** (Staging system for childhood NHL)

All patients received a prephase of

low-dose cyclophosphamide, vincristine, and prednisone (COP).

Administer :

- **2 doses** during each **induction course** (day 1 and day 2),
- **1 dose** on **day 1** of each of the **2 consolidation cycles** (6 doses total)



2- Therapeutic applications

P e d i a t r i c s



3- Non-Hodgkin lymphoma; CD20+, diffuse large B-cell, Burkitt lymphoma, Burkitt-like lymphoma; previously untreated, advanced stage

Infants ≥ 6 months, Children, and Adolescents:

- IV infusion: **375 mg/m²/dose** in combination with systemic (LMB) chemotherapy regimen;
- Administer:
 - ✓ **2 doses** during each **induction course** (day -2 and day 1),
 - ✓ **1 dose** on **day 1** of each of the **2 consolidation cycles** (6 doses total).

Ataxia patients LMB protocol modifications

Group A

Completely resected stage I, & completely resected abdominal stage II

2 courses R-COPAD

D0	D1	D2	D3	D4	D5	D6
Rituximab	Pred CTX VCR Adria	Pred CTX	Pred CTX	Pred	Pred	VCR

Rituximab: 375mg/m²

Prednisone (Pred): 60 mg/m²/day (divided into bid doses) orally (or IV) on days 1-5 inclusive then reduced to zero over 3days.

Cyclophosphamide(CTX): 125 mg/m² 12 hourly as 15-minute infusion on D1-3 inclusive (250 mg/ m² /day) The first dose should be given on day 1 prior to the start of the doxorubicin infusion. Hydration should be maintained at a rate of 3000 mls/m²/day (125 mls/m²/hr). Continue hydration until 12 hours after the last dose of cyclophosphamide.

Adriamycin(Adria): 30mg/m² as 6-hour infusion, after the first dose of Cyclophosphamide.

Vincristine (VCR): 1mg/m² IV bolus D1 and D6. (Maximum single dose 2.0 mg).

Notes:

*CBC, serum electrolytes, BUN, creatinine, bilirubin, SGOT/SGPT before each cycle.

*CR evaluation after COPAD (II) including CSF exam, BM exam and imaging studies of 1ry tumor site should be done after count recovery.

*Re-evaluation will be done by conventional methods [Chest X-ray, Abd US], serum LDH after 2 courses of chemotherapy.

*Avoid ionizing radiation [CT scan, PET CT]

*If patients are not in remission after two courses of COPAD (as confirmed by biopsy), they will then advance to induction in Group C (2nd COPADM8).

Induction
R-COPADM3-I > R-COPADM3-II

D0	D1	D2	D3	D4	D5	D6
Rituximab	Pred HDMTX VCR	Pred CTX Adria IT	Pred CTX	Pred CTX IT*	Pred	IT

Rituximab: 375mg/m²

Prednisone (Pred): 60 mg/m²/day (divided into BID doses) orally on days 1 -5 inclusive then reduced to zero over 3 days.

Cyclophosphamide(CTX): 125 mg/m² 12 hourly as 15-minute infusion on D2-4 inclusive (250 mg/ m² /day) The first dose should be given on day 1 prior to the start of the doxorubicin infusion. Hydration should be maintained at a rate of 3000 mls/m²/day (125 mls/m²/hr). Continue hydration until 12 hours after the last dose of cyclophosphamide.

Adriamycin (Adria): 30mg/m² as 6-hour infusion, after the first dose of Cyclophosphamide.

Vincristine (VCR): 1mg/m² IV bolus D1 (Maximum single dose 2.0 mg).

High Dose Methotrexate (HDMTX): 1g/m² iv infusion over 4 hours with rescue Ca leucovorin according to protocol.

IT (HC + MTX) *: Additional Intrathecal on D4 for patients with **CNS disease (Group C)**.

Calcium Leucovorin: 15 mg/m² orally every 6 hours for a total of 12 doses (or as required depending on methotrexate levels).

Consolidation

CYMI

CYMII

D0	D1	D2	D3	D4	D5	D6	D7
Rituximab	HDMTX	Ara-C IT ₁	Ara-C	Ara-C IT*	Ara-C	Ara-C	IT ₂

Rituximab: 375mg/m²

High Dose Methotrexate (HDMTX): 1g/m² 1g/m² iv infusion over 4 hours with rescue Ca leucovorin according to protocol .

Calcium Leucovorin: 15 mg/m² orally every 6 hours for a total of 12 doses (or as required depending on methotrexate levels).

Cytarabine (Ara-C): 100 mg/m² in 1000 mls/m² dextrose saline as infusion over 24 hours. Repeat daily from Day 2-6 inclusive.

IT₁ (HC + MTX): Each dose adjusted.

IT₂ (HC + Ara-C): Each dose adjusted.

IT (HC + MTX) *: Additional Intrathecal on D4 for patients with **CNS disease (Group C)**.

Further reduction of cytotoxic chemotherapy should be suggested guided by the patient general

Notes:

- *The interval between 1st and 2nd consolidation phase should be reduced to minimum. Each course should be given upon hematological recovery (ANC=1000, Pt=100,000).
- *CBC, Lytes, BUN, Creatinine, Bilirubin, SGOT/SGPT, Uric acid and Phosphorus should be done twice weekly during chemotherapy.
- *CR evaluation after CYM I; including CSF exam, BM exam, imaging studies of 1ry tumor site should be done after count recovery.
- *Re-evaluation will be done by conventional methods [Chest X-ray, Abd US], serum LDH after 2 courses of chemotherapy.
- *Avoid ionizing radiation [CT scan, PET- CT].

2-Therapeutic applications

P e d i a t r i c s

4- Non-Hodgkin lymphoma B-cell, CD 20+; relapsed and refractory:

Children ≥ 11 years and Adolescents:

- IV infusion: **375 mg/m²** administered **on days 1 and 3** of **courses 1 & 2**
- **On day 1 only** of **course 3** ; used in combination with ifosfamide, carboplatin, and etoposide (ICE)

Clinical considerations !!!!

For oncology uses, a **uricostatic agent** (eg, allopurinol) and **aggressive hydration** are recommended for patients at risk for tumor lysis syndrome



Preparation & Administration



3- Preparation


- Rituximab should be diluted in an infusion bag of either 0.9% sodium chloride or 5% dextrose in water.
- No other drugs should be mixed with it, and patients should be closely monitored for infusion reactions.

4- Administration guidelines

- Rituximab and its biosimilars are approved for use by intravenous infusion.
- They should not be administered as IV bolus or push.
- Patients should be given **acetaminophen** and **antihistamine** before each infusion.

5-Emetic potential

- ❑ Pediatrics and Adults: **Minimal** (<10%)
- ❑ Dose-adjusted EPOCH and rituximab (DA-EPOCH-R), a **treatment regimen** used for subsets of patients with aggressive non-Hodgkin B-cell lymphoma, is considered **highly emetogenic** as it contains the combination of doxorubicin and cyclophosphamide.
- ❑ The DA-EPOCH multiday regimen includes :
 - ❑ **etoposide, vincristine, doxorubicin** : infused continuously over days 1 to 4,
 - cyclophosphamide given on day 5,
 - 60 mg/m² of prednisone taken by mouth twice daily on days 1 to 5



6-Management of adverse effects

Risk points	Managing toxicity
Bowel obstruction/perforation : <ul style="list-style-type: none">▪ Association between depletion of B-cells and severe colitis.	<ul style="list-style-type: none">• Tell your healthcare provider right away if you have any stomach-area pain during treatment with Rituxan
Cytopenia : <ul style="list-style-type: none">• Rituximab is associated with lymphopenia, leukopenia, neutropenia, thrombocytopenia, and anemia;• Duration of cytopenias may be prolonged and may extend months beyond treatment.	<ul style="list-style-type: none">✓ Monitor CBC
Renal toxicity: <p>May cause fatal renal toxicity in patients with non-Hodgkin lymphomas (NHL). & Renal toxicity also occurred due to tumor lysis syndrome.</p>	<ul style="list-style-type: none">➤ Monitor kidney functions

Risk points	Managing toxicity
<p>Tumor lysis syndrome :</p> <ul style="list-style-type: none"> ▪ Leading to acute renal failure requiring dialysis (sometimes fatal) may occur within 12 to 24 hours following the first dose when used as a single agent in the treatment of NHL. ▪ Hyperkalemia, hypocalcemia, hyperuricemia, and/or hyperphosphatemia may occur. 	<ul style="list-style-type: none"> ❑ Administer prophylaxis : <u>antihyperuricemic therapy</u> and <u>aggressive hydration</u> in patients at high risk ❑ Correct electrolyte abnormalities and administer supportive care as indicated.
<p>Serious Infusion-related reactions</p> <p>Reactions may include : angioedema, bronchospasm, cough, hypotension, hypoxia, throat irritation, urticaria, and in more severe cases, acute respiratory distress syndrome, pulmonary infiltrates, acute myocardial infarction, ventricular fibrillation, and cardiogenic shock</p>	<p>Management of Rituximab anaphylaxis :</p> <p>Hypersensitivity reactions to first-dose rituximab treatment can be reduced by <u>priming intravenous lines with Standardized premedications</u>, including :</p> <ul style="list-style-type: none"> ➤ acetaminophen, ➤ diphenhydramine ➤ dexamethasone

Risk points	Managing toxicity
<p>Hypogammaglobulinemia (LOW AB) and Infection ; (dose related)</p> <p>Serious, including fatal, bacterial, fungal, and new or reactivated viral infections can occur during and following the completion of rituximab-based therapy</p> <p>Infections have been reported in some patients with prolonged hypogammaglobulinemia. New or reactivated viral infections have included CMV viremia, herpes simplex infection, parvovirus B19 seroconversion, varicella zoster infection, West Nile virus, and hepatitis C and B.</p>	<p>❑ Management :</p> <p><u>Trimethoprim-sulfamethoxazole prophylaxis</u> prevents severe/life-threatening infections following rituximab in antineutrophil cytoplasm antibody-associated vasculitis</p>



7-Drug interaction :

Amphotericin B & Cisplatin : both increase nephrotoxicity and/or ototoxicity (category c)



8- WHAT ARE WARNINGS AND PRECAUTIONS FOR RITUXIMAB?

1- Mucocutaneous reactions (severe):

Severe, including fatal, mucocutaneous reactions reported including paraneoplastic pemphigus, **Stevens-Johnson syndrome**, lichenoid **dermatitis**, vesiculobullous dermatitis, and **toxic epidermal necrolysis**

2- Progressive multifocal leukoencephalopathy:

- John Cunningham virus infection resulting in progressive multifocal leukoencephalopathy and death has been reported in patients treated with rituximab

3- Reactivation of hepatitis B:

- Screen all patients for HBV infection before initiating drug by measuring hepatitis B surface **antigen** (HBsAg) and hepatitis B core antibody (anti-HBc)
- Consult with hepatitis experts regarding monitoring and use of HBV **antiviral** therapy when screening identifies patients at risk of HBV reactivation due to evidence of prior HBV infection

Difference between Rituximab & Brentuximab ?

- Brentuximab vedotin is an **immunotoxin targeting CD30-expressing cells**, including those in Hodgkin lymphoma
- It is approved for treatment of **relapse/refractory Hodgkin lymphoma**
- First line treatment of Hodgkin lymphoma.
- Compared to conventional chemotherapy, Both rituximab and brentuximab vedotin are better tolerated with less toxicity.



Adverse effects :

Immediate effects after having brentuximab (infusion related)

You might have a **rash, shortness of breath, difficulty breathing, a cough, a tight chest, fever, back pain, chills, headache**, feeling sick or being sick. These can happen within a few minutes or up to several hours after having brentuximab. This can be life threatening.



Thank you !

Any questions ?

References

1. Yelvington BJ. Subcutaneous Rituximab in Follicular Lymphoma, Chronic Lymphocytic Leukemia, and Diffuse Large B-Cell Lymphoma. *J Adv Pract Oncol*. 2018 Jul-Aug;9(5):530-534. [[PMC free article](#)] [[PubMed](#)]
2. Maloney DG, Smith B, Rose A. Rituximab: mechanism of action and resistance. *Semin Oncol*. 2002 Feb;29(1 Suppl 2):2-9. [[PubMed](#)]
3. Pierpont TM, Limper CB, Richards KL. Past, Present, and Future of Rituximab-The World's First Oncology Monoclonal Antibody Therapy. *Front Oncol*. 2018;8:163. [[PMC free article](#)] [[PubMed](#)]
4. Griffin TC, Weitzman S, Weinstein H, et al. A study of rituximab and ifosfamide, carboplatin, and etoposide chemotherapy in children with recurrent/refractory B-cell (CD20+) non-Hodgkin lymphoma and mature B-cell acute lymphoblastic leukemia: a report from the Children's Oncology Group. *Pediatr Blood Cancer*. 2009;52(2):177-181.[[PubMed 18816698](#)]
5. Kasi PM, Tawbi HA, Oddis CV, Kulkarni HS. Clinical review: serious adverse events associated with the use of rituximab - a critical care perspective. *Crit Care*. 2012;16(4):231. doi:10.1186/cc11304[[PubMed 22967460](#)]
6. Salles G, Barrett M, Foà R, et al. Rituximab in B-cell hematologic malignancies: a review of 20 years of clinical experience. *Adv Ther*. 2017;34(10):2232-2273. doi:10.1007/s12325-017-0612-x[[PubMed 28983798](#)]
7. Shah S, Geetha D. Place in therapy of rituximab in the treatment of granulomatosis with polyangiitis and microscopic polyangiitis. *Immunotargets Ther*. 2015;4:173-183. doi:10.2147/ITT.S55516[[PubMed 27471722](#)]

References

- 8- Kasi PM, Tawbi HA, Oddis CV, Kulkarni HS. Clinical review: serious adverse events associated with the use of rituximab - a critical care perspective. *Crit Care*. 2012;16(4):231. doi:10.1186/cc11304[[PubMed 22967460](#)]
- 9- Lenz HJ. Management and preparedness for infusion and hypersensitivity reactions. *Oncologist*. 2007;12(5):601-609. doi:10.1634/theoncologist.12-5-601
- 10- Makatsori M, Kiani-Alikhan S, Manson AL, et al. Hypogammaglobulinaemia after rituximab treatment-incidence and outcomes. *QJM*. 2014;107(10):821-828. doi:10.1093/qjmed/hcu094
- 11-Christou EAA, Giardino G, Worth A, Ladomenou F. Risk factors predisposing to the development of hypogammaglobulinemia and infections post-Rituximab. *Int Rev Immunol*. 2017;36(6):352-359. doi:10.1080/08830185.2017.1346092[[PubMed 28800262](#)]
- 12- Kosmidis ML, Dalakas MC. Practical considerations on the use of rituximab in autoimmune neurological disorders. *Ther Adv Neurol Disord*. 2010 Mar;3(2):93-105.
- 13-Yelvington BJ. Subcutaneous Rituximab in Follicular Lymphoma, Chronic Lymphocytic Leukemia, and Diffuse Large B-Cell Lymphoma. *J Adv Pract Oncol*. 2018 Jul-Aug;9(5):530-534.
- 14-Jennette JC, Falk RJ, Bacon PA, Basu N, Cid MC, Ferrario F, Flores-Suarez LF, Gross WL, Guillevin L, Hagen EC, et al. 2012 revised International Chapel Hill consensus conference nomenclature of vasculitides. *Arthritis Rheum*. 2013;65(1):1–11.

References

- 15-** Bartlett NL, Chen R, Fanale MA, et al. Retreatment with brentuximab vedotin in patients with CD30-positive hematologic malignancies. *J. Hematol. Oncol.* 2014;**7**:24. doi: 10.1186/1756-8722-7-24. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
- 16-** Younes A, Gopal AK, Smith SE, et al. Results of a pivotal phase II study of brentuximab vedotin for patients with relapsed or refractory Hodgkin's lymphoma. *J. Clin. Oncol.* 2012;**30**:2183–2189. doi: 10.1200/JCO.2011.38.0410. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]