



Entresto

(sacubitril / valsartn)



Contents



01

Place in therapy

02

**Pharmacological
category**

03

**Mechanism of
action**

07

Drug interactions

04

Administration

05

Pediatric dose

06

Toxicity

08

**Patient Education
guide**



01

Place in therapy

Heart Failure



Entresto (sacubitril / valsartan)



“Used mainly in Heart Failure ”.

Place in therapy:

- ❑ Patients with chronic symptomatic class **II-III HFrEF**
- ❑ In replacement to current **ACEI/ARB**
- ❑ In addition to beta-blockers
- ❑ **Reduced morbidity/mortality. (BG therapy)**




- ❑ In a recent case series, **four patients** with **chemotherapy-related acute cardiac failure** with severely reduced ejection fraction were successfully treated with sacubitril/valsartan.
- ❑ Sacubitril/valsartan was also demonstrated to be valuable in **anthracycline-related cardiac toxicity**
- ❑ In a recent clinical trial, sacubitril/valsartan emerged as a promising treatment option in patients with **refractory CTRCD**.

02


Pharmacological category

Angiotensin receptor-
Neprilysin inhibitor (ARNI)





03




Mechanism of action

How ARNI works ?



Heart Failure and Compensatory Mechanisms



NPS ↑

- 1- Elevated BNP and NT-pro BNP
- 2- Natriuretic diuresis (BV)
- 3- Dec BP ↓

BR

- 1- SNS activation
- 2- HR ↑
- 3- BP ↑
- 3- CO ↑


RAAS

- 1- Renin sec ↑
- 2- Na/H₂O retention
- 3- BP ↑
- 4- Co ↑

Cardiac Remolding

- 1- Ventricular wall distention
- 2- Myocyte growth
- 3- Hypertrophy

NPS : **decreases blood pressure (BP)** , **lowers the sympathetic tone** , **and reduces aldosterone levels**.
antagonistically to the RAAS and has **favorable effects on the pathogenesis of heart failure**



Sacubitril/valsartan is a combination product.
Where it's a pro-drug that, upon activation,
acts as ARNI :



Sacubitril

(Npyralisin inhibitor)

- Block the action of neprilysin,
- Thus preventing the breakdown of natriuretic peptides,
- Prolonged duration of the favorable effects of these peptides

However, because neprilysin breaks down angiotensin II & Bradykinin, So inhibiting neprilysin will **accumulate Ang. II. & Bradykinin**

Valsartan

(ARBS)

- Blocking the RAAS system.
- Dec BP and CO

For this reason, a **neprilysin inhibitor cannot be used alone**; it must always be combined with an ARB to **block the effect of the excess angiotensin II**

Monitoring Parameter



- ✓ **NOTE for Monitoring Parameter :**
- ✓ **Because sacubitril/valsartan therapy inhibits the breakdown of brain natriuretic peptide (BNP) :**

Therefore :

- **BNP will be elevated in patients taking this drug. BNP will not be a reliable marker of heart failure exacerbations in these patients.**
- **NT-pro-BNP is not a substrate for neprilysin and is therefore not affected by sacubitril. Therefore, NT-pro-BNP should be utilized in patients on sacubitril/valsartan when **a heart failure exacerbation is suspected****

04

Administration



24/26 mg

(sacubitril 24 mg
and valsartan 26 mg)



49/51 mg

(sacubitril 49 mg
and valsartan 51 mg)



97/103 mg

(sacubitril 97 mg
and valsartan 103 mg)

Entresto Administration ?



- ❑ Sacubitril/valsartan is available as an oral tablet in three dosage strengths containing :
 - ✓ **Sacubitril** (24 mg, 49 mg, or 97 mg)
 - ✓ **Valsartan** (26 mg, 51 mg, or 103 mg).
- ❑ Valsartan component in this combination has a higher bioavailability than regular valsartan tablets :
- **Valsartan 26 mg, 51 mg, and 103 mg** in the brand-name combination are equivalent to valsartan 40 mg, 80 mg, and 160 mg in other formulations, respectively

Entresto Administration ?



- ☐ **Twice a day (BID)** and maybe administered without regard to meals.
- ☐ Allow at least a **36-hour washout period** when switching from an **ACEI** before starting sacubitril/valsartan.
- ☐ Patients **must be tolerated an ACEI or an ARB** before being started on sacubitril/valsartan.
- ☐ Clinicians can replace sacubitril/valsartan **oral suspension** at the recommended tablet dosage in patients unable to swallow tablets.

05

Pediatric dose



Pediatric dose :



Table 1: Recommended Dose Titration

	Titration Step Dose (twice daily)		
	Starting	Second	Final
Pediatric Patients Less than 40 kg [†]	1.6 mg/kg	2.3 mg/kg	3.1 mg/kg
Pediatric Patients At least 40 kg, less than 50 kg	24/26 mg	49/51 mg	72/78 mg [‡]
Pediatric Patients At least 50 kg	49/51 mg	72/78 mg [‡]	97/103 mg

- ❑ **Mild to moderate** Kidney impairment (eGFR ≥ 30 mL/minute/1.73 m²):

No dosage adjustment necessary.

- ❑ **Severe impairment** (eGFR < 30 mL/minute/1.73 m²):
Initial:

Reduce the usual starting dose by 50%, then follow the recommended dose escalation to titrate dose.



The background of the slide is a collage of various pharmaceutical products. It includes several blister packs containing capsules and tablets. The capsules are in various colors: orange, blue, red and white, and yellow. Some are oval-shaped, while others are round. The blister packs are made of a shiny, metallic material. The overall image is a close-up, slightly angled view of these medications, creating a textured and colorful background.

06 Toxicity

Adverse Effects

- Hypotension,
- Hyperkalemia,
- Renal failure,
- Cough,

➤ **Note :**

Sacubitril/valsartan was associated with :

- ✓ lower risk of elevation in serum potassium or serum creatinine and a lower risk of cough than enalapril.
- ✓ More patients experienced angioedema in the sacubitril/valsartan arm than in the enalapril; however, this outcome was not statistically significant




Contraindications

- ❑ Hypersensitivity
- ❑ History of angioedema due to an ACEI or ARBs
- ❑ In diabetic patients receiving the renin inhibitor, aliskiren, specifically, the valsartan (any ARB), is contraindicated with aliskiren due to an **increased risk of hypotension, hyperkalemia, and renal impairment.**
- ❑ Patients who have received an ACE-inhibitors within 36 hours due to increased risk of angioedema,

Box warning

- ❖ Drugs that work directly on the renin-angiotensin system, injury and/or death to the developing fetus. **When pregnancy is confirmed, discontinue sacubitril/valsartan as soon as possible**



Toxicity of Entresto



Limited literature is available concerning toxicity in human subjects. However, a **single dose of 583 mg sacubitril/617 mg valsartan** in healthy volunteers and **multiple doses of 437 mg sacubitril and 463 mg valsartan** for **14 days** were studied. **Hypotension resulting from overdose requires prompt treatment.**

- ❑ In pharmacokinetics, sacubitril is converted to **LBQ657** (active metabolized compound)
- ❑ All three compounds (sacubitril, LBQ657, and valsartan) are **highly bound (94% to 97%) to plasma protein.**
- ❑ Hence, it is **unlikely to be removed by hemodialysis.**



07

Drug Interactions



Drug interactions



1- Aliskiren:

- ❑ May enhance the **hyperkalemic effect** of Angiotensin II Receptor Blockers.
- ❑ May enhance the **hypotensive effect** of Angiotensin II Receptor Blockers.
- ❑ May enhance the **nephrotoxic effect** of Angiotensin II Receptor Blockers.
- **Management:** Aliskiren use with ACEIs or ARBs in patients with diabetes is contraindicated
- ❖ **Risk D: Consider therapy modification**

2- Amifostine:

- ❑ Blood Pressure Lowering Agents may enhance the hypotensive effect of Amifostine.
- **Management:** When used at chemotherapy doses, hold blood pressure lowering medications for 24 hours before amifostine administration.
- ❖ **Risk D: Consider therapy modification**


3- Angiotensin-Converting Enzyme Inhibitors: (ACEI)

- ❑ May enhance the **adverse/toxic effect of Sacubitril.** (bradykinin ↑)
- ❑ Specifically, the risk of **angioedema** may be increased with this combination.
- ❖ **Risk X: Avoid combination**




08

Patient Education



Patient Education

- ✓ Entresto is usually given **twice daily**.
- ✓ May be taken with or without food.
- ✓ Report any signs or symptoms of **angioedema** :
(swelling of the face or throat, difficulty breathing)
immediately to your doctor.
- ✓ May cause a fall in blood pressure that may be noticed as **light-headedness**; call your doctor and ask for advice, symptoms usually resolve with continued therapy.
- ✓ Ensure you **do not become dehydrated**.



Patient Education



- ✓ **Do not use salt substitutes containing potassium** without first consulting your doctor.
- ✓ Entresto is usually used in addition to other drugs to lower blood pressure, and **all lifestyle recommendations given to you** by your doctor (such as losing weight if overweight, smoking cessation, partaking in regular exercise, and limiting sodium intake) should also be followed.
- ✓ **In addition**, other conditions, such as high cholesterol levels or diabetes if present, also need to be **controlled**.
- ✓ Do not take any other medications including those bought over-the-counter without checking with your doctor or pharmacist.

Thanks

Do you have any questions?

Dr. Salma Mohamed Abdelmoneim

salma.abdelmoneim11@gmail.com

01111821906



CREDITS: This presentation template was created by **Slidesgo**, including icons by **Flaticon** and infographics & images by **Freepik**



References

- 1.Sauer AJ, Cole R, Jensen BC, Pal J, Sharma N, Yehya A, Vader J. Practical guidance on the use of sacubitril/valsartan for heart failure. Heart Fail Rev. 2019 Mar;24(2):167-176. [[PMC free article](#)] [[PubMed](#)]
- 2.Eadie AL, Brunt KR, Herder M. Exploring the Food and Drug Administration's review and approval of Entresto (sacubitril/valsartan). Pharmacol Res Perspect. 2021 May;9(3):e00794. [[PMC free article](#)] [[PubMed](#)]
- 3.Vicent L, Esteban-Fernández A, Gómez-Bueno M, De-Juan J, Díez-Villanueva P, Iniesta ÁM, Ayesta A, González-Saldívar H, Rojas-González A, Bover-Freire R, Iglesias D, García-Aguado M, Perea-Egido JA, Martínez-Sellés M. Sacubitril/Valsartan in Daily Clinical Practice: Data From a Prospective Registry. J Cardiovasc Pharmacol. 2019 Feb;73(2):118-124. [[PubMed](#)]
- 4.Canale ML, Coviello K, Solarino G, Del Meglio J, Simonetti F, Venturini E, Camerini A, Maurea N, Bisceglia I, Tessa C, Casolo G. Case Series: Recovery of Chemotherapy-Related Acute Heart Failure by the Combined Use of Sacubitril Valsartan and Wearable Cardioverter Defibrillator: A Novel Winning Combination in Cardio-Oncology. Front Cardiovasc Med. 2022;9:801143. [[PMC free article](#)] [[PubMed](#)]
- 5 Gregoriotti V, Fernandez TL, Costa D, Chahla EO, Daniele AJ. Use of Sacubitril/valsartan in patients with cardio toxicity and heart failure due to chemotherapy. Cardiooncology. 2020 Nov 05;6(1):24. [[PMC free article](#)] [[PubMed](#)]

References

6. Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM, Deswal A, Drazner MH, Dunlay SM, Evers LR, Fang JC, Fedson SE, Fonarow GC, Hayek SS, Hernandez AF, Khazanie P, Kittleson MM, Lee CS, Link MS, Milano CA, Nwacheta LC, Sandhu AT, Stevenson LW, Vardeny O, Vest AR, Yancy CW. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2022 May 03;145(18):e895-e1032. [[PubMed](#)]
7. Du AX, Westerhout CM, McAlister FA, Shanks M, Oudit GY, Paterson DI, Hanninen M, Thomas J, Ezekowitz JA. Titration and Tolerability of Sacubitril/Valsartan for Patients With Heart Failure in Clinical Practice. *J Cardiovasc Pharmacol*. 2019 Mar;73(3):149-154. [[PubMed](#)]
8. Hubers SA, Brown NJ. Combined Angiotensin Receptor Antagonism and Neprilysin Inhibition. *Circulation*. 2016 Mar 15;133(11):1115-24. [[PMC free article](#)] [[PubMed](#)]
9. In brief: Sacubitril/valsartan (Entresto) expanded indication. *Med Lett Drugs Ther*. 2021 May 03;63(1623):65. [[PubMed](#)]
10. Cada DJ, Baker DE, Leonard J. Sacubitril/Valsartan. *Hosp Pharm*. 2015 Nov;50(11):1025-36. [[PMC free article](#)] [[PubMed](#)]

References

11. Drugs and Lactation Database (LactMed®) [Internet]. National Institute of Child Health and Human Development; Bethesda (MD): Feb 28, 2019. Sacubitril. [[PubMed](#)]
12. Entresto (sacubitril and valsartan) [Package Insert] Updated 02/2022. Novartis Pharmaceuticals Corporation <https://www.drugs.com/pro/entresto.html>
- 13-Nicolas D, Kerndt CC, Reed M. StatPearls [Internet]. StatPearls Publishing; Treasure Island (FL): May 26, 2022. Sacubitril/Valsartan. [[PubMed](#)]
- 14-Fala L. Entresto (Sacubitril/Valsartan): First-in-Class Angiotensin Receptor Neprilysin Inhibitor FDA Approved for Patients with Heart Failure. Am Health Drug Benefits. 2015 Sep;8(6):330-4. [[PMC free article](#)] [[PubMed](#)]
- 15.McCullough PA, Mehta HS, Barker CM, Van Houten J, Mollenkopf S, Gunnarsson C, Ryan M, Cork DP. Mortality and guideline-directed medical therapy in real-world heart failure patients with reduced ejection fraction. Clin Cardiol. 2021 Sep;44(9):1192-1198. [[PMC free article](#)] [[PubMed](#)]