



Patient Name:	Wt.
Patient ID:	Ht.
Pathology:	BSA
Diagnosis: Stage:	Age

R-DHAP

Salvage Chemotherapy for Relapsed or Refractory for Hodgkin Lymphoma and Non-Hodgkin Lymphoma

Date:

***Premedication 30 minutes prior to Rituximab Perfelgan 1000 mg, Chlorphenaramine 10 mg & Hydrocortisone 100 mg (Day 0)**

• Rituximab mg (375mg/m²) + ml NS over 4 hours (Day 0)

• Dexamethasone (40 mg/day) once daily over 30 minutes (Day 1 to Day 4)

Anti-Emetics:

• Palonosetron_____mcg (20 mcg/kg) max 1.5mg IV+ 50 ml NS IV infusion over 20 min, half an hour before Chemotherapy q48hr (Day1 & Day3).

• ****Aprepitant:** D1 _____mg orally then _____ mg D2 & D3 orally
Start: 1h before chemo; Max: 125 mg on day 1, 80 mg/day on days 2-3.

Prehydration :

..... ml Glucose 5%, Sodium chloride 0.45% (125 mL/ m²/hr) + Mannitol

20% (2ml/kg) Over 6 hr. (Day 1)

• Cisplatin mg (100 mg/ m²) + Sodium chloride 0.9% (125 mL/ m²) +

Mannitol (2ml/kg) over 6 hr (Day 1)

Post-hydration:

..... ml Glucose 5%, Sodium chloride 0.45% (125 mL/ m²/hr) + potassium chloride

..... ml (20meq/L) + ml Mg (10 mmol/L) over 12 hr. (Day 1)

• High Dose Cytarabine mg +..... ml G/NS IV Q12 over 3 hrs (Day 2 x 2 Doses)

• Dexamethasone Eye drops 2 drops in each eye twice daily for 7 days

****Dosing: IF age ≥ 6 months old, >6 kg, give 3 mg/kg PO x1 on day 1, then 2 mg/kg PO on D2-3, Oral suspension.
IF age ≥ 12 years old, give: 125 mg PO x1 on day 1, then 80 mg PO on D2-3, Oral cap.**