


Data Assimilation Research Testbed Tutorial



Section 5: Comprehensive Filtering Theory: Non-Identity Observations and the Joint Phase Space

Version 1.0: June, 2005

A More General Context for Filtering with Geophysical Models

Dynamical system governed by (stochastic) Difference Equation:

$$dx_t = f(x_t, t) + G(x_t, t)d\beta_t, \quad t \geq 0 \quad (1)$$

Observations at discrete times:

$$y_k = h(x_k, t_k) + v_k; \quad k = 1, 2, \dots; \quad t_{k+1} > t_k \geq t_0 \quad (2)$$

Observational error white in time and Gaussian (nice, not essential).

$$v_k \rightarrow N(0, R_k) \quad (3)$$

Complete history of observations is:

$$Y_\tau = \{y_l; t_l \leq \tau\} \quad (4)$$

Goal: Find probability distribution for state at time t:

$$p(x, t | Y_t) \quad (5)$$

A More General Context for Filtering with Geophysical Models

State between observation times obtained from Difference Equation.
Need to update state given new observation:

$$p(x, t_k | Y_{t_k}) = p(x, t_k | y_k, Y_{t_{k-1}}) \quad (6)$$

Apply Bayes rule:

$$p(x, t_k | Y_{t_k}) = \frac{p(y_k | x_k, Y_{t_{k-1}}) p(x, t_k | Y_{t_{k-1}})}{p(y_k | Y_{t_{k-1}})} \quad (7)$$

Noise is white in time (3) so:

$$p(y_k | x_k, Y_{t_{k-1}}) = p(y_k | x_k) \quad (8)$$

Integrate numerator to get normalizing denominator:

$$p(y_k | Y_{t_{k-1}}) = \int p(y_k | x) p(x, t_k | Y_{t_{k-1}}) dx \quad (9)$$

A More General Context for Filtering with Geophysical Models

Probability after new observation:

$$p\left(x, t_k | Y_{t_k}\right) = \frac{p(y_k | x) p(x, t_k | Y_{t_{k-1}})}{\int p(y_k | \xi) p(\xi, t_k | Y_{t_{k-1}}) d\xi} \quad (10)$$

Exactly analogous to earlier derivation except that x and y are vectors.

EXCEPT, no guarantee we have prior sample for each observation.

SO, let's make sure we have priors by 'extending' state vector.

A More General Context for Filtering with Geophysical Models

Extending the state vector to joint state-observation vector.

Recall: $y_k = h(x_k, t_k) + v_k; k = 1, 2, \dots; \quad t_{k+1} > t_k \geq t_0 \quad (2)$

Applying h to x at a given time gives expected values of observations.

Get prior sample of obs. by applying h to each sample of state vector x .

Let $z = [x, y]$ be the combined vector of state and observations.

A More General Context for Filtering with Geophysical Models

NOW, we have a prior for each observation:

$$p(z, t_k | Y_{t_k}) = \frac{p(y_k | z) p(z, t_k | Y_{t_{k-1}})}{\int p(y_k | \xi) p(\xi, t_k | Y_{t_{k-1}}) d\xi} \quad (10.\text{ext})$$

Dealing with Many Observations

One more issue: how to deal with many observations in set y_k ?

Let y_k be composed of s subsets of observations: $y_k = \{y_k^1, y_k^2, \dots, y_k^s\}$

Observational errors for obs. in set i independent of those in set j .

$$\text{Then: } p(y_k | z) = \prod_{i=1}^s p(y_k^i | z)$$

Can rewrite (10.ext) as series of products and normalizations.

Dealing with Many Observations

One more issue: how to deal with many observations in set y_k ?

Implication: can assimilate observation subsets sequentially.

If subsets are scalar (individual obs. have mutually independent error distributions), can assimilate each observation sequentially.

If not, have two options:

1. Repeat everything above with matrix algebra.
2. Do singular value decomposition; diagonalize obs. error covariance.
Assimilate observations sequentially in rotated space.
Rotate result back to original space.


Good news: Most geophysical obs. have independent errors!

How an Ensemble Filter Works for Geophysical Data Assimilation

1. Use model to advance **ensemble** (3 members here) to time at which next observation becomes available.


Ensemble state estimate after using previous observation (**analysis**).

Ensemble state at time of next observation (**prior**).



How an Ensemble Filter Works for Geophysical Data Assimilation


2. Get prior ensemble sample of observation, $y=h(x)$, by applying forward operator h to each ensemble member.



Theory: observations from instruments with uncorrelated errors can be done sequentially.


How an Ensemble Filter Works for Geophysical Data Assimilation

3. Get **observed value** and **observational error distribution** from observing system.




How an Ensemble Filter Works for Geophysical Data Assimilation

4. Find **increment** for each prior observation ensemble (this is a scalar problem for uncorrelated observation errors).




How an Ensemble Filter Works for Geophysical Data Assimilation

5. Use ensemble samples of y and each state variable to linearly regress observation increments onto state variable increments.



How an Ensemble Filter Works for Geophysical Data Assimilation

6. When all ensemble members for each state variable are updated, have a new analysis. Integrate to time of next observation...



Non-Identity Observation Operators in Lorenz-63:

Try observing $\text{mean}(x, y)$, $\text{mean}(y, z)$, $\text{mean}(z, x)$
Same error variance and frequency as previously

In *models/lorenz_63/work*:

Edit *input.nml*

Change *obs_sequence_in_name* in *filter.nml* to
obs_seq.out.average.

Execute *./filter* program to produce a new assimilation.
Look at the error statistics and time series with matlab.
Record the error and spread values and compare to identity case.

Error is much larger!

Identity observations remove all regression error; **can be very misleading.**