

R M M

ROMANIAN MATHEMATICAL MAGAZINE

www.ssmrmh.ro



If $0 < m \leq a_i \leq M, i \in \overline{1, n}, n \in \mathbb{N}, n > 2$ then:

$$\sum_{i=1}^n \frac{a_i}{\sqrt{\sum_{\substack{j=1 \\ j \neq i}}^n a_j^2}} \leq \frac{n}{2\sqrt{n-1}} \left(\frac{m}{M} + \frac{M}{m} \right)$$

Proposed by Seyran Ibrahimov-Masilli-Azerbaijan

Solution 1 by Michael Sterghiou-Greece, Solution 2 by Florică Anastase-Romania

Solution 1 by Michael Sterghiou-Greece

$$\sum_{i=1}^n \frac{a_i}{\sqrt{\sum_{\substack{j=1 \\ j \neq i}}^n a_j^2}} \leq \frac{n}{2\sqrt{n-1}} \left(\frac{m}{M} + \frac{M}{m} \right); \quad (1)$$

$f(t) = \sqrt{t}$ is concave on $(0, \infty)$

$$\sum_{i=1}^n \sqrt{\frac{a_1^2}{x - a_i^2}} \leq n \cdot \sqrt{\frac{\left(\frac{p}{n}\right)^2}{x - \left(\frac{p}{n}\right)^2}}; \quad (2)$$

R M M

ROMANIAN MATHEMATICAL MAGAZINE

www.ssmrmh.ro

Where $x = \sum_{i=1}^n a_i^2$ and $p = \sum_{i=1}^n a_i$ with $x \geq \frac{p^2}{n}$.

$$(2) \Rightarrow LHS_{(1)} \stackrel{Jensen}{\leq} n \cdot \sqrt{\frac{p^2}{n^2 x - p^2}} \leq n \cdot \sqrt{\frac{p^2}{n^2 \cdot \frac{p^2}{n} - p^2}} = \frac{n}{\sqrt{n-1}}; \quad (3)$$

$$\text{But } \frac{m}{M} + \frac{M}{m} \stackrel{AM-GM}{\geq} 2 \Rightarrow \frac{1}{2} \left(\frac{m}{M} + \frac{M}{m} \right) \geq 1 \text{ and from (3)}$$

Since $LHS_{(1)} \leq \frac{n}{\sqrt{n-1}}$ it follows that:

$$\sum_{i=1}^n \frac{a_i}{\sqrt{\sum_{\substack{j=1 \\ j \neq i}}^n a_j^2}} \leq \frac{n}{2\sqrt{n-1}} \left(\frac{m}{M} + \frac{M}{m} \right)$$

Solution 2 by Florică Anastase-Romania

$$\text{Let } S = \sum_{i=1}^n a_i$$

From Cauchy-Schwartz Inequality we have:

$$\begin{cases} a_1^2 + a_2^2 + \dots + a_{n-1}^2 \geq \frac{(a_1 + a_2 + \dots + a_{n-1})^2}{n-1} \\ \dots \\ a_2^2 + a_3^2 + \dots + a_n^2 \geq \frac{(a_2 + a_3 + \dots + a_n)^2}{n-1} \end{cases} \Rightarrow$$

$$\begin{cases} \frac{a_n}{\sqrt{a_1^2 + a_2^2 + \dots + a_{n-1}^2}} \leq \frac{a_n \sqrt{n-1}}{S - a_n} \\ \dots \\ \frac{a_1}{\sqrt{a_2^2 + a_3^2 + \dots + a_n^2}} \leq \frac{a_1 \sqrt{n-1}}{S - a_1} \end{cases}$$

$$\sum_{i=1}^n \frac{a_i}{\sqrt{\sum_{\substack{j=1 \\ j \neq i}}^n a_j^2}} \leq \sqrt{n-1} \left(\frac{a_1}{S - a_1} + \frac{a_2}{S - a_2} + \dots + \frac{a_n}{S - a_n} \right) \stackrel{(1)}{\leq} \frac{n}{\sqrt{n-1}}$$

R M M

ROMANIAN MATHEMATICAL MAGAZINE

www.ssmrmh.ro

$$n \left(\frac{a_1}{S-a_1} + \frac{a_2}{S-a_2} + \dots + \frac{a_n}{S-a_n} \right) \leq \frac{S}{S-a_1} + \frac{S}{S-a_2} + \dots + \frac{S}{S-a_n} \Leftrightarrow$$

$$(n-1) \left(\frac{a_1}{S-a_1} + \frac{a_2}{S-a_2} + \dots + \frac{a_n}{S-a_n} \right) \leq \frac{S-a_1}{S-a_1} + \frac{S-a_2}{S-a_2} + \dots + \frac{S-a_n}{S-a_n} \Leftrightarrow$$

$$\frac{a_1}{S-a_1} + \frac{a_2}{S-a_2} + \dots + \frac{a_n}{S-a_n} \leq \frac{n}{n-1} \Rightarrow (1) \text{ true.}$$

$$\text{But } \frac{m}{M} + \frac{M}{m} \stackrel{AM-GM}{\geq} 2 \Rightarrow \frac{1}{2} \left(\frac{m}{M} + \frac{M}{m} \right) \geq 1; \quad (2)$$

From (1), (2) it follows that:

$$\sum_{i=1}^n \frac{a_i}{\sqrt{\sum_{\substack{j=1 \\ j \neq i}}^n a_j^2}} \leq \frac{n}{2\sqrt{n-1}} \left(\frac{m}{M} + \frac{M}{m} \right)$$