



## **4MM013 - Computational Mathematics**

## Mathematics Assignment-1

Full Marks: 10

University ID : 2329767

Submitted by : Alan Shah

Submitted on : 4/13/2023

1. State the definition of a function and a composite function.


[2Marks]

Let f and g be functions defined as follows:  
f: R o R, f(x) = 
$$\frac{x-3}{x+1}$$
, f(2) =? and  
g: R o R, g(x) =  $\frac{1}{x}$ ,  $x \neq 0$ 

g: R 
$$\rightarrow$$
 R, g(x) =  $\frac{1}{x}$ ,  $x \neq 0$ 

Calculate  $(f \circ g)(x)$  and  $(g \circ f)(x)$ .


|    | 16: id:  |
|----|--|
|    | Uni id:  |
| 1  | Mon Oshah 2329767  |
| 4. | The state of the s |
| 7  | ena; rule 4 0 10 10  |
| 7  | Timelian is matter that definer dedorming  |
| -  | Function: Tunction is a vittle that tefner dedecuring the value of the depotent variables to values of the   |
| -  | and variables be values of the   |
| 1  | independent variables. Til con ber depar   |
| -  | depicited in overeal methods to de including a lable, a bounula on a grouph.   |
| -  | industrial a hope a framuela an a grouph.  |
| 1  | more day of the same   |
| 7- | Compaide Roselien: At is a function which is within another function one Punction is continued to a the input do another function:   |
| -  | and Profession is  |
| -  | Och Hit led so the word do   |
|    | angell as Resident   |
|    |  |
| -  | P: R → R, P(n) n-3 PG) =?  |
| -  | 21+1   |
| -  | $g(n) = \frac{1}{n}, n \neq 0$   |
| -  |  |
|    | fag (n) = ? . gaf (n)=?  |
| -  | 0 0  |
| -  | P(2) = ? = -1/3  |
|    | f(2) = 2-3 = -1/2  |
|    | 2+1  |
|    | Ozan.  |
|    | gof (n) = ?  |
| _  | $Q_{q}(P(n)) = Q(n-3) = V_{21} = 1$  |
|    | 0 ( ari ) n-3  |
|    | $g(R(n)) = g(n-3) = \sqrt{n} = 1$ $g(R(n)) = g(n-3) = \sqrt{n} = 1$ $\frac{n-3}{n+1}$  |
|    |  |
|    |  |
| -  |  |
|    |  |
|    |  |




2. Solve the following using the inverse matrix method:

[ 2 Marks]

$$6x - y = 0$$






3. Calculate the inverse of the following functions: a.  $f(x) = \frac{6+x}{7}$ b.  $f(x) = \frac{3}{2-x}$ 


a. 
$$f(x) = \frac{6+x}{7}$$

b. 
$$f(x) = \frac{3}{3}$$

[ 2 Marks]



4. Sketch the graph of the following functions: [2 Marks]  $f(x) = 2(x-1)^2 + 3$ , -2 < x < 2 In the interval of x=0.5



5.

a. Define gradient of a function. State the gradient and intercept of :

$$2y + 8 = 6x$$

[2 Marks]

b. Solve the following equations:

$$x^2 + \frac{17}{6}x + 2 = 0$$

| g. | Hon Olah 2329767   |
|----|--|
| 1  | d  |
|    | The gradient of a Rundien is the direction of rate of the draction of the function:  Gradient of intercept of:  or, of : 3n-4  comparing the a eqn (1) with y = mn+c  or, y = 3n-4  comparing the a eqn (1) with y = mn+c  or, y = 3n-4  comparing the a eqn (1) with y = mn+c |
|    | 2 6 6 11 - 63  |
|    | intercept)   |
| 6) | Oblin.   |
|    | topony n2 + 17/6 n + z = 0   |
|    | Comparing about eg' with an +sn'   |
|    | Comparing above egt with and +bn+0=0   |
|    | 1000   |
|    | 22 -b± 18-40C  |
|    | - La   |
|    | 11-12  |
|    | = -176 + 1 (176)2-4x1x2  |
|    |  |
|    | = 13 t t   |
|    |  |
|    | 2  |
|    | Toking tue: Taking -ve:  |
|    | Taking tve: Taking -ve: 176 - 473 - 176 - 176  |
|    |  |
|    | -3/2   |
|    | : 22 - 1/3 or - 3/2  |
|    |  |

The End