

First Principles: A Mathematics Handbook

Alan D. Smith

First Principles: A Mathematics Handbook

© 2025 Alan D. Smith

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without prior written permission of the author, except as permitted by law.

This book is intended for educational purposes. While every effort has been made to ensure accuracy, the author assumes no responsibility for errors, omissions, or any consequences arising from the use of this material.

First edition, 2025

Published by Amazon KDP

ISBN: 9798297344082

Set in LaTeX using Pandoc and XeLaTeX.

Cover design by the author.

*For my brother, Samuel,
whose strength of character, humour, and curiosity
continue to inspire my work and life.*

Contents

Introduction	1
Purpose of This Guide	1
How to Use This Handbook	1
Notation and Conventions	1
1 Exponents and Logarithms	3
1.1 Exponents	3
1.2 Logarithms	3
1.2.1 Logarithmic scales	4
1.3 Euler's number and the natural logarithm	4
2 Sequences and Series	5
2.1 Sigma and Product Notation	5
2.2 Arithmetic Sequences and Series	6
2.3 Geometric Sequences and Series	7
3 Functions	9
3.1 Testing if a relation is a function	9
3.2 Domain and Range	10
3.3 Linear functions	10
3.4 Quadratic functions	10
3.4.1 Line of symmetry	10
3.4.2 Discriminant	10
3.4.3 The Quadratic formula	10
4 Finance	11
4.1 Important Concepts	11
4.2 Parameters and Conventions	11
4.3 Interest Rates	12
4.3.1 Nominal and Effective Rates	12
4.4 Compound Interest	12
4.5 Continuous Compounding	13

5	Sets	15
5.1	Types of Sets	15
5.2	Subsets	15
5.3	Standard number sets	16
5.4	Set operations	17
5.4.1	Union	17
5.4.2	Intersection	17
5.4.3	Mutual exclusivity	17
6	Combinatorics	19
6.1	Basic Counting Principles	19
6.1.1	Multiplication Principle	19
6.1.2	Addition Principle	19
6.2	Factorials	20
6.3	Permutations	20
6.3.1	Permutations with Identical Objects	21
6.4	Combinations	22
6.5	Partitions of Sets	22
6.6	Partitions of Objects (Stars and Bars)	22
6.7	Binomial Expansion	23
6.7.1	Pascal's Triangle	23
6.8	Multinomial Theorem	24
7	Sectors and Radians	25
8	Geometry	27
9	Trigonometry	29
10	Statistics	31
11	Probability	33
12	Random Variables and Probability Distributions	35
13	Hypothesis Testing	37
14	Differentiation	39
15	Integration	41
16	Differential Equations	43
17	Complex Numbers	45
18	Vectors	47

19 Matrices	49
19.1 Eigenvalues and Eigenvectors	49
19.2 Power Formula	49
20 Graph Theory	51
20.1 Adjacency Matrices	51
20.2 Transition Matrices	51
21 Modelling	53
21.1 Logistic Functions	53
Appendix 1. Formula Reference	55
Appendix 2. Exercise Solutions	65
Appendix 3. References and Notes	67
Appendix 4. Worked Examples Template Page (TEMP)	69

Introduction

Purpose of This Guide

This handbook aims to be a concise yet comprehensive reference for key mathematical first principles, covering topics essential for high school mathematics, university preparation, and STEM-related fields. It is designed for fast navigation, allowing readers to look up topics easily, find clear explanations, quickly learn new topics or revise specific ones.

How to Use This Handbook

Each chapter focuses on a single topic, beginning with definitions and essential formulas. Mathematical notation follows a standardised convention throughout. Where topics are built upon, cross-references to other sections are included for further exploration or for revisiting relevant material.

Notation and Conventions

- Definitions are shown in *italics*.
- Variables are italicised (e.g. x, y, z).
- Constants such as e or π retain their standard mathematical meaning.
- Essential formulas are highlighted in boxes (e.g. $e^{i\pi} + 1 = 0$).
- All angles are expressed in radians unless stated otherwise.

Chapter 1

Exponents and Logarithms

1.1 Exponents

An exponent is written as a^n , where a is the *base*, and n is the *power* (also known as *index* or *exponent*).

Exponents follow a set of algebraic rules:

Rules of Exponents

$$a^m \cdot a^n = a^{m+n}$$

$$a^m \div a^n = a^{m-n}$$

$$(a^m)^n = a^{mn}$$

$$a^0 = 1$$

$$a^1 = a$$

$$a^{-m} = \frac{1}{a^m}$$

$$a^{\frac{1}{m}} = \sqrt[m]{a}$$

$$a^{\frac{n}{m}} = (\sqrt[m]{a})^n$$

$$(ab)^n = a^n \cdot b^n$$

$$\left(\frac{a}{b}\right)^n = \frac{a^n}{b^n}$$

1.2 Logarithms

The logarithmic function $\log_a b = x$ is the solution to the exponential equation $a^x = b$. In other words, $\log_a b$ gives the power to which the base a needs to be raised in order to produce b . Therefore,

$a^x = b \iff \log_a b = x$

Base restrictions: Unless stated otherwise, $a > 0$, $a \neq 1$, and $b > 0$.
(An exception would be for complex logarithms, for example.)

- The logarithm with base e is written as $\ln x$ (natural logarithm).
- The logarithm with base 10 is typically written as $\log x$ (common logarithm).

Laws of logarithms

$$\log_a x + \log_a y = \log_a (xy)$$

$$\log_a x - \log_a y = \log_a \left(\frac{x}{y}\right)$$

$$n \log_a x = \log_a (x^n)$$

1.2.1 Logarithmic scales

When dealing with incredibly small or large numbers, it becomes difficult to distinguish them, so they can be scaled using logarithms.

1.3 Euler's number and the natural logarithm

Euler's number ($e \approx 2.718 \dots$) is an irrational constant, which arises in situations involving growth or decay, such as compound interest or radioactive decay.

A definition for e can be found in finance, through continuous compounding:

$$e = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n$$

where n is the number of compounding periods per unit time, and e is the resulting constant as n grows infinitely large (Euler's number). See **4. Finance** for more information.

The *natural logarithm*, written as $\ln x$, is the logarithm with base e :

$\ln x = \log_e x$

Note: e^x and $\ln x$ have unique properties in calculus. See **Chapter [[14. Differentiation]]**, and **Chapter [[15. Integration]]** for more information.

Chapter 2

Sequences and Series

A numerical *sequence* is an ordered list of numbers which can often be described by a formula or recurrence relation. Each number within the sequence is called a *term*. The general term within a sequence is called the n^{th} term.

A numerical *series* is the sum of a range of terms in a sequence. A *finite series* describes a series which has n terms ($u_1, u_2, u_3, \dots, u_n$, where u_n is the n^{th} term of the sequence), and thus also has a finite sum. An *infinite series* describes a series which has infinite terms (u_1, u_2, u_3, \dots), which can *converge* to a finite sum or *diverge* to infinity.

2.1 Sigma and Product Notation

Sigma notation can be useful when describing a series concisely. For example, the sum:

$$u_1 + u_2 + u_3 + u_4 + u_5 + \dots + u_{100}$$

can be rewritten using sigma notation as:

$$\sum_{n=1}^{100} u_n$$

In general,

$$\boxed{\sum_{n=a}^b u_n}$$

represents the sum of all terms from $n = a$ to $n = b$.

Sigma notation also follows certain important algebraic properties:

Addition Property:

$$\sum_{n=1}^k (a_n + b_n) = \sum_{n=1}^k a_n + \sum_{n=1}^k b_n$$

Constant Multiplication Property:

Given c is a constant:

$$\sum_{n=1}^k c \cdot u_n = c \cdot \sum_{n=1}^k u_n$$

Product notation is similar to sigma notation. It also represents a repeated operation, but it denotes multiplication rather than summation. For example, the product:

$$u_1 \cdot u_2 \cdot u_3 \cdots u_{100}$$

can be written concisely as:

$$\prod_{n=1}^{100} u_n$$

In general, the product of all terms from $n = a$ to $n = b$ can be written as:

$$\prod_{n=a}^b u_n$$

2.2 Arithmetic Sequences and Series

An *arithmetic sequence* (sometimes called an arithmetic *progression*) is a sequence where each term differs from the previous term by a constant value, called the *common difference*, typically denoted as d .

The general formula for the n^{th} term of an arithmetic sequence is given by:

$$u_n = u_1 + (n - 1)d$$

where: - u_n is the n^{th} term, - d is the common difference, - n is the position of the term (e.g., for the 5th term, $n = 5$).

The sum of a finite *arithmetic series* is given by:

$$S_n = \frac{n}{2}(u_1 + u_n) \quad \text{OR} \quad S_n = \frac{n}{2}(2u_1 + (n-1)d)$$

where S_n is the sum of the first n terms.

2.3 Geometric Sequences and Series

A *geometric sequence* (sometimes called a *geometric progression*) is a sequence where each term is the product of the previous term and a constant value, called the *common ratio*, typically denoted as r .

The general formula for the n^{th} term of a geometric sequence is given by:

$$u_n = u_1 r^{n-1}$$

The sum of a finite *geometric series* is given by:

$$S_n = \frac{u_1(r^n - 1)}{r - 1} \quad \text{OR} \quad S_n = \frac{u_1(1 - r^n)}{1 - r} \quad \text{where } r \neq 1.$$

where r is the common ratio.

If $|r| < 1$, the geometric series converges as $n \rightarrow \infty$, resulting in an infinite geometric series with sum:

$$S_\infty = \frac{u_1}{1 - r} \quad \text{where } |r| < 1.$$

Chapter 3

Functions

A *relation* between two variables x and y , is any set of points which are on the (x, y) plane.

There exist four different types of relations:

- One to one
- One to many
- Many to one
- Many to many

A *function* is defined as a mapping onto a single value. Therefore one to one, and many to one relations are considered to be functions.

3.1 Testing if a relation is a function

1. **Algebraic Method:** If we substitute any value of x and it results in a singular y -value, then it can be considered a function.
2. **Graphical Method (Vertical Line Test):** If we are given a plot of a function on the (x, y) plane, where the x -axis is parallel with the horizontal, and the y -axis is parallel with the vertical, then if we are able to draw a vertical line anywhere on the graph, and it only intersects the plot once, it is a function. If the vertical line intersects the plot more than once, then it cannot be considered a function.

3.2 Domain and Range

3.3 Linear functions

Linear functions are functions which can be written in the form:

$$\boxed{y = mx + c} \quad \text{OR} \quad \boxed{y = m(x - x_1) + y_1}$$

3.4 Quadratic functions

3.4.1 Line of symmetry

For functions in the form $f(x) = ax^2 + bx + c$, the axis of symmetry of is

$$\boxed{x = -\frac{b}{2a}}$$

3.4.2 Discriminant

$$\boxed{\Delta = b^2 - 4ac}$$

3.4.3 The Quadratic formula

The solutions to $ax^2 + bx + c = 0$ are:

$$\boxed{x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}} \quad \text{where } a \neq 0$$

Chapter 4

Finance

4.1 Important Concepts

- *Interest* is money paid at a regular rate in order to use lent money, or to delay the repayment of debt.
- *Inflation* is the increase in the price of goods and services with respect to time. Inflation means that a fixed amount of capital will be able to purchase fewer goods and services as prices increase.
- *Depreciation* is the decrease in the value of an asset with respect to time.
- The *principal* is the original loan amount that is borrowed.
- The *principal portion* of a payment is the portion that reduces the outstanding loan balance (principal).
- The *interest portion* is the payment covering the interest charged for that period.
- *Amortisation* is a process whereby a loan is repaid through a series of regular payments. In an amortised loan, interest for each period is calculated on the outstanding loan balance, therefore as regular payments reduce the principal, the interest portion of each payment decreases over time, whilst the principal portion increases.

4.2 Parameters and Conventions

In order to work with financial formulas, it is crucial to first understand the standard notation on which they depend.

Calculator Label	Symbol in This Book	Meaning
N	N	Total number of compounding periods
I%	r	Nominal annual interest rate (%)
	r_{eff}	Effective annual interest rate (%)
PV	V_0	Present value of the loan/investment
PMT	M	Payment per period
FV	V_f	Future value of loan/investment
P/Y	n_p	Number of payments per year
C/Y	n_c	Number of compounding periods per year

Note: This topic is primarily solved using calculators or software tools which allow for the automation of iterative calculations. Regardless, it is important to recognise the meaning of each parameter and their relation to the relevant formulae.

4.3 Interest Rates

Interest rates are a measure of the cost of borrowing money, or the return on investment over a set period of time.

$$\text{Interest Paid} = \text{Total Repayments} - \text{Amount Borrowed}$$

4.3.1 Nominal and Effective Rates

The *nominal annual interest rate* (r) is the yearly rate of interest which does not consider compounding. Financial calculators often denote this as I%.

The *effective annual interest rate* (r_{eff}) considers the effect of compounding throughout the year. It reflects the percentage increase in value, or cost of debt, over one year.

The formula for the effective annual interest rate is given as follows:

$$r_{\text{eff}} = \left(1 + \frac{r}{n_c}\right)^{n_c} - 1$$

4.4 Compound Interest

Compound interest is when the interest is calculated using any accumulated interest from previous periods in addition to the original principal.

The general formula for compound interest is:

$$V_f = V_0 \left(1 + \frac{r}{n_c}\right)^{n_c t}$$

where t is the time in years.

4.5 Continuous Compounding

When compounding occurs infinitely often (as $n_c \rightarrow \infty$), the formula becomes:

$$V_f = V_0 e^{rt}$$

Chapter 5

Sets

A *set* is a collection of distinct objects, called *elements* (or *members*). They are typically denoted by a capital letter (such as A, B, C , etc.), and their elements can be written inside curly brackets. For example:

$$A = \{1, 2, 3, 4, e, 9.\bar{9}, 8086\}$$

Here, $1 \in A$ denotes that “1 is an element of A ”, and $5 \notin A$ means that “5 is not an element of A ”.

5.1 Types of Sets

Name	Description	Example
Finite set	Contains a finite number of elements.	$B = \{1, 2, 3\}$
Infinite set	Contains an infinite number of elements	$\mathbb{N} = \{0, 1, 2, 3, \dots\}$
Singleton set	Contains exactly one element.	$C = \{256\}$
Universal set	Contains all objects under consideration in a certain context. Denoted as U .	
Empty set	(Sometimes called the <i>null set</i>) Contains no elements, and is denoted as \emptyset or $\{\}$	

5.2 Subsets

A set A is considered a *subset* of B if every element in A is also an element of B . In which case:

$$A \subseteq B$$

which means: “ A is a subset of or equal to B ”.

If A is a subset of B , but $A \neq B$, then A is considered a *proper subset* of B , shown as:

$$A \subset B$$

The empty set (\emptyset) is a subset of every set, including singletons, since it has no elements.

5.3 Standard number sets

There are a range of standard number sets which the reader should be aware of:

Symbol	Set name	Definition
\mathbb{N}	Natural numbers	$\{0, 1, 2, 3, \dots\}$
\mathbb{Z}	Integers	$\{\dots, -2, -1, 0, 1, 2, \dots\}$
\mathbb{Q}	Rational numbers	$\frac{p}{q}$, where $p, q \in \mathbb{Z}, q \neq 0$
\mathbb{R}	Real numbers	All \mathbb{Q} , as well as irrationals such as $\pi, e, \sqrt{2}$
\mathbb{C}	Complex numbers	All numbers in the form $a + bi$ where $a, b \in \mathbb{R}, i^2 = -1$

As shown by Figure 5.1, the sets are nested in the following way:

$$\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R} \subset \mathbb{C}$$

Note: The set of natural numbers (\mathbb{N}) is sometimes defined to exclude 0. In this book, we state that $0 \in \mathbb{N}$ for consistency. Complex numbers are introduced fully in **Chapter [[17. Complex Numbers]]**.




Figure 5.1: Number set hierarchy

5.4 Set operations

5.4.1 Union

5.4.2 Intersection

5.4.3 Mutual exclusivity

$$A \cap B = \emptyset$$

Chapter 6

Combinatorics

6.1 Basic Counting Principles

6.1.1 Multiplication Principle

The *Fundamental Counting Principle* (or the *Multiplication Principle*) states that if an event can occur in m ways, and another *independent* event is able to occur in n ways, then the total number of unique ways both events can occur is:

$$m \cdot n$$

For example, if there are m Formula 1 drivers, and n Formula 1 cars, each driver can be paired with any car. Therefore there are a total of mn different driver–car combinations that can be made.

This principle can be generalised to any number of events. If there are k independent events, where the i^{th} event can occur in n_i ways, then the total number of unique ways all of the events can occur is:

$$\prod_{i=1}^k n_i$$

6.1.2 Addition Principle

The *Addition Principle* states that given an event can occur in m ways, and another *mutually exclusive* event (recall mutual exclusivity from **Chapter** [[5. Sets]]) can occur in n ways, then the total number of ways either event can occur is:

$$m + n$$

Which can be generalised in the same way. If there are k mutually exclusive events, where the i^{th} event can occur in n_i ways, then the total number of possible outcomes is:

$$\sum_{i=1}^k n_i$$

6.2 Factorials

The *factorial* of a positive integer n is written as $n!$, and defined as the product of all positive integers from 1 to n :

$$n! = \prod_{i=1}^n i, \quad n \geq 1$$

By convention, $0! = 1$.

6.3 Permutations

A *permutation* is a unique ordering of a set of objects. For example ABC , and CBA are both permutations of three letters A , B , and C .

Note: If you take AAB , swapping the position of the two A s is still considered the same permutation, since the result is still AAB .

The number of permutations of n distinct objects taken in groups of r at a time, is written as nPr , is given by:

$$nPr = \frac{n!}{(n-r)!} \quad (\text{Assuming no identical objects})$$

From here we can see that if all r objects are used ($r = n$):

$$nPr = \frac{n!}{(n-n)!} = \frac{n!}{(0)!} = \frac{n!}{1} = n!$$

So the special case where all objects are used ($r = n$):

$$nPr = n! \quad \text{when } r = n$$

6.3.1 Permutations with Identical Objects

The formula for the number of permutations given earlier assumes all objects are distinct. In situations where some objects are identical (e.g. the example given earlier with AAB), there are fewer distinct permutations, as some are repeated.

The general formula is:

$$\frac{n!}{\prod_{i=1}^r n_i!}$$

where n is the total number of objects, n_1, n_2, \dots, n_r are the counts for each group of identical objects.

Worked Example 6A

Find the number of permutations of the letters in the word: MISSISSIPPI.

We use the general formula for permutations with identical objects:

$$\frac{n!}{\prod_{i=1}^r n_i!}$$

MISSISSIPPI has $n = 11$ letters, with 4 'I's, 4 'S's, 2 'P's, and 1 'M'. So the number of permutations is:

$$\begin{aligned} & \frac{11!}{4! \cdot 4! \cdot 2! \cdot 1!} \\ &= \frac{11 \cdot 10 \cdot 9 \cdot 8 \cdot 7 \cdot 6 \cdot 5 \cdot 4!}{4! \cdot 4! \cdot 2} \\ &= \frac{11 \cdot 10 \cdot 9 \cdot 8 \cdot 7 \cdot 6 \cdot 5}{4! \cdot 2} \\ &= \frac{11 \cdot 10 \cdot 9 \cdot 8 \cdot 7 \cdot 6 \cdot 5}{24 \cdot 2} \end{aligned}$$

Simplifying the denominator: $24 \cdot 2 = 48 = 6 \cdot 8$, which can be cancelled from the numerator:

$$\begin{aligned} &= 11 \cdot 10 \cdot 9 \cdot 7 \cdot 5 \\ &= 990 \cdot 7 \cdot 5 \end{aligned}$$

Worked Example 6A

= 34 650 permutations

6.4 Combinations

A *combination* is a set of objects, meaning that order does not matter. For example $\{A, B, C\}$ and $\{C, B, A\}$ are the same combination of three letters, irrespective of the fact they are different permutations.

Since each group of r objects can be arranged in $r!$ different orders, all representing the same combination, the number of combinations of n distinct objects in groups of r , written as nCr or $\binom{n}{r}$ (“ n choose r ”), is given by:

$$nCr = \binom{n}{r} = \frac{nPr}{r!} = \frac{n!}{r!(n-r)!}$$

We also define that $nC0 = 1$ (there is one way to choose nothing), and $nCn = 1$ (there is one way to choose all objects):

$$\binom{n}{0} = 1 \quad \text{and} \quad \binom{n}{n} = 1$$

6.5 Partitions of Sets

A *partition* of a set is a way to group its elements into non-empty subsets, where each element belongs to exactly one subset.

For example, the set $\{A, B, C\}$ has the following partitions:

- $\{\{A, B, C\}\}$
- $\{\{A\}\{B, C\}\}$
- $\{\{B\}\{A, C\}\}$
- $\{\{C\}\{A, B\}\}$
- $\{\{A\}\{B\}\{C\}\}$

Therefore there are 5 different ways to partition the set $\{A, B, C\}$ in total.

6.6 Partitions of Objects (Stars and Bars)

The *stars and bars* method is a way to count the number of ways n identical objects can be distributed into r distinct groups. The method involves representing the n objects as “stars” (\star), and the partitions between the groups as “bars” ($|$).

So taking another example of 5 objects and 3 groups, may look like:

$$\star \star \mid \star \mid \star \star$$

This represents the grouping of $(2, 1, 2)$.

Since $r - 1$ bars are required to create r groups, there are a total of $n + r - 1$ slots (positions). The number of ways to do this is:

$$\boxed{\binom{n+r-1}{r-1}}$$

6.7 Binomial Expansion

The *binomial theorem* gives a formula allowing us to expand powers of a binomial expression of the form $(a + b)^n$, without having to multiply out each term by hand. It states that (for any non-negative integer n):

$$(a + b)^n = \sum_{r=0}^n \binom{n}{r} a^{n-r} b^r$$

For example, the binomial expansion for $n = 3$ is:

$$(a + b)^3 = \binom{3}{0} a^3 + \binom{3}{1} a^2 b + \binom{3}{2} a b^2 + \binom{3}{3} b^3 = a^3 + 3a^2 b + 3a b^2 + b^3$$

6.7.1 Pascal's Triangle

The coefficients of $\binom{n}{r}$ which appear in the expansion can also be found through *Pascal's triangle*, where each number is the sum of the two numbers directly above it:

$$\begin{array}{ccccccc} 1 & & & & & & \\ 1 & 1 & & & & & \\ 1 & 2 & 1 & & & & \\ 1 & 3 & 3 & 1 & & & \\ 1 & 4 & 6 & 4 & 1 & & \\ 1 & 5 & 10 & 10 & 5 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{array}$$

6.8 Multinomial Theorem

The *multinomial theorem* is a generalisation of the binomial theorem to allow for the expansion of powers of a sum with more than two terms. This allows it to be applied to expressions such as $(a + b + c + \dots)^n$. It is formally defined as:

$$\left(\sum_{i=1}^m x_i \right)^n = \sum_{\substack{k_1+k_2+\dots+k_m=n \\ k_1, k_2, \dots, k_m \geq 0}} \frac{n!}{\prod_{j=1}^m k_j!} \prod_{r=1}^m x_r^{k_r}$$

Although this form is relatively cluttered, so typically, the *multinomial coefficient* is introduced:

$$\binom{n}{k_1, k_2, \dots, k_m} = \frac{n!}{\prod_{i=1}^m k_i!}, \text{ where } k_1 + k_2 + \dots + k_m = n$$

Notice that for $m = 2$, this becomes the binomial coefficient:

$$\binom{n}{k_1, k_2} = \frac{n!}{k_1! k_2!},$$

Since $k_1 + k_2 = n$, we can write $k_2 = n - k_1$, giving:

$$\binom{n}{k_1, k_2} = \frac{n!}{k_1! (n - k_1)!},$$

Therefore, for $m = 2$, the multinomial coefficient reduces to the binomial coefficient:

$$\binom{n}{k_1, k_2} = \binom{n}{k_1}, \quad (\text{since } k_2 = n - k_1)$$

Chapter 7

Sectors and Radians

Chapter 8

Geometry

Chapter 9

Trigonometry

Chapter 10

Statistics

Chapter 11

Probability

Chapter 12

Random Variables and Probability Distributions

Chapter 13

Hypothesis Testing

Chapter 14

Differentiation

Chapter 15

Integration

Chapter 16

Differential Equations

Chapter 17

Complex Numbers

We know that $\forall x \in \mathbb{R}, x^2 \geq 0$, but let $i^2 = -1$

Chapter 18

Vectors

Chapter 19

Matrices

19.1 Eigenvalues and Eigenvectors

19.2 Power Formula

$$A^n = PD^nP^{-1}$$

where A is a diagonalisable square matrix, $n \in \mathbb{N}$, P is the matrix of eigenvectors of A , and D is the diagonal matrix of the corresponding eigenvalues

Chapter 20

Graph Theory

20.1 Adjacency Matrices

$a_{ij} = 1$ if there exists an edge from vertex i to vertex j , otherwise $a_{ij} = 0$, for an unweighted graph.

If the graph is weighted, $a_{ij} = w_{ij}$, where w_{ij} represents the weight of the edge from i to j , or 0 if there exists no edge.

20.2 Transition Matrices

a_{ij} is the probability of moving from vertex i to vertex j in a given step. (**Note:** Some curricula define a_{ij} as the probability of moving from j to i)

Chapter 21

Modelling

21.1 Logistic Functions

The general form of a logistic function is:

$$\frac{L}{1 + Ce^{-kx}}, \quad \text{where } L, C, k \in \mathbb{R}^+,$$

and L is the function's maximum value, C is a constant resultant from the initial conditions, and k is the growth rate.

Appendix 1. Formula Reference

Exponents and Logarithms

Rules of Exponents

$$\begin{aligned}
 a^m \cdot a^n &= a^{m+n} \\
 a^m \div a^n &= a^{m-n} \\
 (a^m)^n &= a^{mn} \\
 a^0 &= 1, \text{ for all } a \text{ in } \mathbb{C}, a \neq 0 \\
 a^1 &= a \\
 a^{-m} &= \frac{1}{a^m} \\
 a^{\frac{1}{m}} &= \sqrt[m]{a} \\
 a^{\frac{n}{m}} &= (\sqrt[m]{a})^n \\
 (ab)^n &= a^n \cdot b^n \\
 \left(\frac{a}{b}\right)^n &= \frac{a^n}{b^n} \\
 a^x = b &\Leftrightarrow \log_a b = x
 \end{aligned}$$

*Laws of Logarithms (real case:
 $a, x, y > 0$, and $a \neq 1$)*

$$\begin{aligned}
 \log_a x + \log_a y &= \log_a (xy) \\
 \log_a x - \log_a y &= \log_a \left(\frac{x}{y}\right) \\
 n \log_a x &= \log_a (x^n)
 \end{aligned}$$

Sequences and Series

Sigma notation properties

Addition property

$$\sum_{n=1}^k (a_n + b_n) = \sum_{n=1}^k a_n + \sum_{n=1}^k b_n$$

Constant Multiplication

$$\sum_{n=1}^k c \cdot u_n = c \cdot \sum_{n=1}^k u_n$$

Property

Arithmetic Sequences and Series

The n^{th} term of an arithmetic sequence

$$u_n = u_1 + (n - 1)d$$

The sum of a finite arithmetic series

$$S_n = \frac{n}{2}(u_1 + u_n)$$

Sequences and Series

	OR $S_n = \frac{n}{2} (2u_1 + (n-1)d)$
<i>Geometric Sequences and Series</i>	
The n^{th} term of a geometric sequence	$u_n = u_1 r^{n-1}$
The sum of a finite arithmetic series (where $r \neq 1$)	$S_n = \frac{u_1(r^n - 1)}{r - 1}$
	OR $S_n = \frac{u_1(1 - r^n)}{1 - r}$
The sum of an infinite geometric series	$S_\infty = \frac{u_1}{1 - r}$, where $ r < 1$

Functions

Equation of a straight line	$y = mx + c$ OR $y = m(x - x_1) + y_1$
Line of symmetry	If $f(x) = ax^2 + bx + c$, the axis of symmetry of is $x = -\frac{b}{2a}$
Discriminant	$\Delta = b^2 - 4ac$
Quadratic formula	The solutions to $ax^2 + bx + c = 0$ are $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$, $a \neq 0$

Finance

Interest paid	Interest Paid = Total Repayments – Amount Borrowed
Effective annual interest rate	$r_{\text{eff}} = \left(1 + \frac{r}{n_c}\right)^{n_c} - 1$
Compound interest (discrete)	$V_f = V_0 \left(1 + \frac{r}{n_c}\right)^{n_c t}$
Compound interest (continuous)	$V_f = V_0 e^{rt}$

Sets

De Morgan's Laws

Union	$(A \cup B)^c = A^c \cap B^c$
Intersection	$(A \cap B)^c = A^c \cup B^c$

Combinatorics

Fundamental Counting Principle

 n factorial

Binomial theorem

Sectors and Radians

Radians and degrees equivalence π radians = 180° Arc length $l = r\theta$ Sector area $A = \frac{1}{2}r^2\theta$

Geometry

Distance between two points a and b in n -dimensional space $\sqrt{\sum_{i=1}^n (b_i - a_i)^2}$ Midpoint of a line segment with endpoints a and b in n -dimensional space $(\frac{1}{2}(a_i + b_i))_{1 \leq i \leq n}$ *Spheres of radius r* Surface area $A = 4\pi r^2$ Volume $V = \frac{4}{3}\pi r^3$ *Pyramids of height h* Volume $V = \frac{1}{3}Ah$ *Parallelogram*Area (Given \mathbf{v} and \mathbf{w} are adjacent sides of a parallelogram) $A = |\mathbf{v} \times \mathbf{w}|$ *Cone*Surface area of the curved surface $A = \pi rl$ Volume (right cone) $V = \frac{1}{3}\pi r^2 h$

Trigonometry

Rules for triangles with sides a, b, c , and angles *A, B, C , where X is opposite to x*

Sine Rule

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

Cosine Rule

$$c^2 = a^2 + b^2 - 2ab \cos C$$

Triangle Area

$$\frac{1}{2}ab \sin C$$

Trigonometric Identities

Pythagorean Identity

$$\cos^2 \theta + \sin^2 \theta = 1$$

Tangent Identity

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

Trigonometry

Double-Angle Identity	$\cos(2\theta) = \cos^2 \theta - \sin^2 \theta$
Reciprocal Identities	$\sin \theta = \frac{1}{\csc \theta}$ $\cos \theta = \frac{1}{\sec \theta}$ $\tan \theta = \frac{1}{\cot \theta}$

Statistics

Percentage error	$\varepsilon = \left \frac{v_a - v_e}{v_e} \right \cdot 100$
<i>Descriptive statistics</i>	
Interquartile Range	$\text{IQR} = Q_3 - Q_1$
Arithmetic Mean	$\bar{x} = \frac{\sum_{i=1}^n f_i x_i}{\sum_{i=1}^n f_i}$
<i>Sampling</i>	
Unbiased estimator of the population variance	$\hat{\sigma}^2 = \frac{n}{n-1} s_n^2$

Probability

Probability of an event E	$\mathbb{P}(E) = \frac{ E }{ U }$
Combined events	$\mathbb{P}(E \cup F) = \mathbb{P}(E) + \mathbb{P}(F) - \mathbb{P}(E \cap F)$
Mutually exclusive events	$\mathbb{P}(E \cap F) = 0$
Probability of E given F	$\mathbb{P}(E F) = \frac{\mathbb{P}(E \cap F)}{\mathbb{P}(F)}$

Random Variables and Probability Distributions

Expected value of a single discrete random variable X	$\mathbb{E}[X] = \sum_i x_i \mathbb{P}(X = x_i)$
<i>Linear Transformation of a single random variable X</i>	
Expected Value	$\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$
Variance	$\text{Var}[aX + b] = a^2 \text{Var}[X]$
<i>Linear combinations of random variables X_1, X_2, \dots, X_n</i>	

Random Variables and Probability Distributions

Expected Value <i>Linear combinations of independent random variables</i> X_1, X_2, \dots, X_n	$\mathbb{E}[\sum_{i=1}^n a_i X_i] = \sum_{i=1}^n a_i \mathbb{E}[X_i]$
Variance <i>Uniform Distribution</i> $X \sim \mathcal{U}(a, b)$	$\text{Var}[\sum_{i=1}^n a_i X_i] = \sum_{i=1}^n a_i^2 \text{Var}[X_i]$ $a = \text{lower bound}$ $b = \text{upper bound}$
Mean (μ)	$\mathbb{E}[X] = \frac{a+b}{2}$
Variance (σ^2) <i>Normal (Gaussian) Distribution</i> $X \sim \mathcal{N}(\mu, \sigma^2)$	$\text{Var}[X] = \frac{(b-a)^2}{12}$ $\mu = \text{mean}$ $\sigma^2 = \text{variance}$
Central Limit Theorem (CLT)	For large n , $\bar{X} \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right)$ (approximately)
<i>Binomial Distribution</i> $X \sim \mathcal{B}(n, p)$	$n = \text{number of trials}$ $p = \text{probability of success}$
Mean (μ)	$\mathbb{E}[X] = np$
Variance (σ^2) <i>Poisson Distribution</i> $X \sim \mathcal{P}(\lambda)$	$\text{Var}[X] = np(1-p)$ $\lambda = \text{mean and variance}$

Hypothesis Testing

z (mean, σ known)	$z = \frac{\bar{x} - \mu_0}{\sigma/\sqrt{n}}$
t (mean, σ unknown)	$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}, \quad df = n - 1$
z (proportion)	$z = \frac{\hat{p} - p_0}{\sqrt{\frac{p_0(1-p_0)}{n}}}$
χ^2 (GoF/independence) <i>Two sample</i>	$\chi^2 = \sum \frac{(O - E)^2}{E}$

Hypothesis**Testing**

$$t \text{ (two-sample, pooled variance)} \quad t = \frac{\bar{x}_1 - \bar{x}_2}{s_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}, \quad s_p^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}$$

$$z \text{ (two-sample proportions)} \quad z = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}(1 - \hat{p})(\frac{1}{n_1} + \frac{1}{n_2})}}, \quad \hat{p} = \frac{x_1 + x_2}{n_1 + n_2}$$

Differentiation

$$\text{Derivative of } f(x) \text{ using first principles} \quad f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$\text{Derivative of } x^n \quad f(x) = x^n \Rightarrow f'(x) = nx^{n-1}$$

*Standard**Derivatives*

$$f(x) = \sin x \Rightarrow f'(x) = \cos x$$

$$f(x) = \cos x \Rightarrow f'(x) = -\sin x$$

$$f(x) = \tan x \Rightarrow f'(x) = \frac{1}{\cos^2 x} = \sec^2 x$$

$$f(x) = \sin x \Rightarrow f'(x) = \cos x$$

$$f(x) = \cos x \Rightarrow f'(x) = -\sin x$$

*Standard**Differentiation**Rules*

$$\text{Chain rule} \quad \frac{d}{dx} f(g(x)) = f'(g(x)) \cdot g'(x)$$

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$$

$$\text{Product rule} \quad \frac{d}{dx} [u(x)v(x)] = u(x)v'(x) + u'(x)v(x)$$

$$\text{Quotient rule} \quad \frac{d}{dx} \left[\frac{u(x)}{v(x)} \right] = \frac{u'(x)v(x) - u(x)v'(x)}{v(x)^2}$$

$$\text{L'Hôpital's Rule} \quad \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}, \text{ for } \frac{0}{0} \text{ or } \frac{\infty}{\infty} \text{ forms, with } g'(x) \neq 0$$

Integration

$$\text{Integral of } x^n \quad \int x^n dx = \frac{x^{n+1}}{n+1} + C, \quad n \neq -1$$

Integration

Area of the region between the x -axis and $f(x)$, for the range (a, b)

$$A = \int_a^b y \, dx$$

Trapezoidal Rule

$$\int_a^b y \, dx \approx \frac{h}{2} (y_0 + y_n + 2(y_1 + \cdots + y_{n-1}))$$

Standard Integrals

$$\begin{aligned} \int \frac{1}{x} \, dx &= \ln|x| + C \\ \int \sin x \, dx &= -\cos x + C \\ \int \cos x \, dx &= \sin x + C \\ \int e^x \, dx &= e^x + C \\ \int \tan x \, dx &= -\ln|\cos x| + C \\ \int \cot x \, dx &= \ln|\sin x| + C \\ \int \sec^2 x \, dx &= \tan x + C \\ \int \sec x \, dx &= \ln|\sec x + \tan x| + C \\ \int \frac{1}{x^2 - a^2} \, dx &= \frac{1}{2a} \ln \left| \frac{x-a}{x+a} \right| + C \\ \int \frac{1}{x^2 + 1} \, dx &= \arctan(x) + C \end{aligned}$$

Differential Equations

Euler's method (step size h)

$$y_{n+1} = y_n + hf(x_n, y_n)$$

Coupled linear differential equations (diagonalisable case)

$$\begin{aligned} x_{n+1} &= x_n + h \\ \mathbf{x}(t) &= \begin{pmatrix} x(t) \\ y(t) \end{pmatrix} = C_1 e^{\lambda_1 t} \vec{p}_1 + C_2 e^{\lambda_2 t} \vec{p}_2 \end{aligned}$$

Complex Numbers

Rectangular form $z = a + bi$

Polar (modulus-argument) form $z = r(\cos \theta + i \sin(\theta))$

Exponential (Euler) form $z = re^{i\theta}$

Compliment conjugate $\bar{z} = a - bi$

De Moivre's theorem $(\cos \theta + i \sin \theta)^n = \cos(n\theta) + i \sin(n\theta)$

Vectors

Magnitude	$ \mathbf{v} = \sqrt{v_1^2 + v_2^2 + v_3^2}$
Scalar product	$ \mathbf{v} = \sqrt{\sum_{i=1}^n v_i^2}, \quad v_i = i^{\text{th}} \text{ component}$ $\mathbf{v} \cdot \mathbf{w} = \sum_{i=1}^n v_i w_i$ $\mathbf{v} \cdot \mathbf{w} = \mathbf{v} \mathbf{w} \cos \theta, \theta = \text{angle between } \mathbf{v}, \mathbf{w}$
Vector product	Given $\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}, \mathbf{w} = \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix}$ $\mathbf{v} \times \mathbf{w} = \begin{pmatrix} v_2 w_3 - v_3 w_2 \\ v_3 w_1 - v_1 w_3 \\ v_1 w_2 - v_2 w_1 \end{pmatrix}$ $ \mathbf{v} \times \mathbf{w} = \mathbf{v} \mathbf{w} \sin \theta$
Vector equation of a line	$\mathbf{r} = \mathbf{a} + \lambda \mathbf{d}, \lambda \in \mathbb{R}$
Parametric form of a line	$x = x_0 + \lambda l, \quad y = y_0 + \lambda m, \quad z = z_0 + \lambda n$

Matrices

Matrix addition	$(A + B)_{ij} = a_{ij} + b_{ij}$
Matrix multiplication	$(AB)_{ij} = \sum_{k=1}^n a_{ik} b_{kj}$ OR $AB = [\sum_{k=1}^n a_{ik} b_{kj}]_{1 \leq i \leq m, 1 \leq j \leq p}$
Determinant of a 2×2 matrix	$\det \begin{pmatrix} a & b \\ c & d \end{pmatrix} = ad - bc$
Determinant of a 3×3 matrix	$\det \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix} =$ $a(ei - fh) - b(di - fg) + c(dh - eg)$
Invertibility condition	A is invertible if $\det A \neq 0$
Power formula of a diagonalisable matrix A	$A^n = P D^n P^{-1}, n \in \mathbb{N}, P = \text{eigenvectors}, D = \text{eigenvalues}$
<i>2D Transformation Matrices</i>	(Assuming x -axis is horizontal, and y -axis is vertical)
Stretch with scale factor h horizontally, and v vertically	$\begin{pmatrix} h & 0 \\ 0 & v \end{pmatrix}$, centred at the origin
Anticlockwise rotation of angle θ	$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$, about the origin

Matrices

Reflection in the line through the origin with angle θ from the x -axis	$\begin{pmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & -\cos 2\theta \end{pmatrix}$
Shear with factor h horizontally, and v vertically	$\begin{pmatrix} 1 & h \\ v & 1 \end{pmatrix}$, about the origin

Graph Theory

Adjacency matrix	$a_{ij} = 1$ if edge $i \rightarrow j$, otherwise 0. If weighted, $a_{ij} = w_{ij}$
Transition matrix	a_{ij} = probability of moving $i \rightarrow j$ (convention can vary, however)
State after n transitions (row-stochastic convention)	$s_n = s_0 T^n$

Modelling

Logistic function	$\frac{L}{1 + Ce^{-kx}}$, where $L, C, k \in \mathbb{R}^+$
<i>Volume of revolution in the range (a, b)</i>	
about the x -axis	$V = \pi \int_a^b [f(x)]^2 dx$
about the y -axis	$V = \pi \int_a^b [f(y)]^2 dy$

Appendix 2. Exercise Solutions

Appendix 3. References and Notes

A significant portion of the material in this handbook was inspired by the notes provided by my IB Mathematics teacher. These resources are not formally published but were incredibly useful in shaping the explanations and content of this book.

References

1. Economopoulos, P., Halsey, T., Doering, S., Ortman, M., Sirinoglu Singh, N., Gray, P., Harris, D., & Wathall, J. C. (2019). *Oxford IB Diploma Programme: IB Mathematics: Applications and Interpretation, Higher Level*. Oxford University Press.
 2. Anderson, I. (2002). *A First Course in Combinatorial Mathematics* (2nd ed.). Oxford University Press.
 3. Wilson, R. J. (1996). *Introduction to Graph Theory* (2nd ed.). Longman.
-

Appendix 4. Worked Examples Template Page (TEMP)

Worked Example I

Find u_{10} for an arithmetic sequence where $u_1 = 3$ and $d = 5$.

We use the n^{th} term formula:

$$u_n = u_1 + (n - 1)d$$

$$u_{10} = 3 + (10 - 1) \cdot 5 = 3 + 45 = 48$$
