CS 70 Discrete Mathematics and Probability Theory Fall 2018 Alistair Sinclair and Yun Song Discussion 08B

1 Let's Talk Probability

- (a) When is $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B)$ true? What is the general rule that always holds?
- (b) When is $\mathbb{P}(A \cap B) = \mathbb{P}(A) * \mathbb{P}(B)$ true? What is the general rule that always holds?
- (c) If A and B are disjoint, are they independent?

Solution:

- (a) In general, we know $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) \mathbb{P}(A \cap B)$. This is the Inclusion-Exclusion Principle. Therefore if A and B are disjoint, such that $\mathbb{P}(A \cap B) = 0$, then $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B)$ holds.
- (b) $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$ holds if and only if *A* and *B* are independent (by definition). The general rule that always holds is $\mathbb{P}(A \cap B) = \mathbb{P}(B|A)\mathbb{P}(A) = \mathbb{P}(A|B)\mathbb{P}(B)$.
- (c) No, if two events are disjoint, we cannot conclude they are independent. Consider a roll of a fair six-sided die. Let A be the event that we roll a 1, and let B be the event that we roll a 2. Certainly A and B are disjoint, as $\mathbb{P}(A \cap B) = 0$. But these events are not independent: $\mathbb{P}(B \mid A) = 0$, but $\mathbb{P}(B) = 1/6$.

Since disjoint events have $\mathbb{P}(A \cap B) = 0$, we can see that the only time when A and B are independent is when either $\mathbb{P}(A) = 0$ or $\mathbb{P}(B) = 0$.

2 Aces

Consider a standard 52-card deck of cards:

- (a) Find the probability of getting an ace or a red card, when drawing a single card.
- (b) Find the probability of getting an ace or a spade, but not both, when drawing a single card.
- (c) Find the probability of getting the ace of diamonds when drawing a 5 card hand.
- (d) Find the probability of getting exactly 2 aces when drawing a 5 card hand.
- (e) Find the probability of getting at least 1 ace when drawing a 5 card hand.
- (f) Find the probability of getting at least 1 ace or at least 1 heart when drawing a 5 card hand.

Solution:

- (a) Inclusion-Exclusion Principle: $\frac{4}{52} + \frac{26}{52} \frac{2}{52} = \frac{28}{52} = \frac{7}{13}$.
- (b) Inclusion-Exclusion, but we exclude the intersection: $\frac{4}{52} + \frac{13}{52} 2 \cdot \frac{1}{52} = \frac{15}{52}$.
- (c) Ace of diamonds is fixed, but the other 4 cards in the hand can be any other card: $\frac{\binom{51}{4}}{\binom{52}{5}}$.
- (d) Account for the number of ways to draw 2 aces and the number of ways to draw 3 non-aces: $\frac{\binom{4}{2}\cdot\binom{48}{3}}{\binom{52}{5}}.$
- (e) Complement to getting no aces: $1 \frac{\binom{48}{5}}{\binom{52}{5}}$.
- (f) Complement to getting no aces and no hearts: $1 \frac{\binom{36}{5}}{\binom{52}{5}}$. This is because 52 13 3 = 36, where 13 is the number of hearts and 3 is the number of non-heart aces.
- 3 Balls and Bins

Throw n balls into n bins.

- (a) What is the probability that the first bin is empty?
- (b) What is the probability that the first k bins are empty?
- (c) Use the union bound to give an upper bound on the probability that at least k bins are empty.
- (d) What is the probability that the second bin is empty given that the first one is empty?
- (e) Are the events that "the first bin is empty" and "the first two bins are empty" independent?
- (f) Are the events that "the first bin is empty" and "the second bin is empty" independent?

Solution:

- (a) The probability that ball *i* does not land in the first bin is $\frac{n-1}{n}$. The probability that all of the balls do not land in the first bin is $\left(\frac{n-1}{n}\right)^n$.
- (b) The probability that ball *i* does not land in the first *k* bins is $\frac{n-k}{n}$. The probability that all of the balls do not land in the first *k* bins is $\left(\frac{n-k}{n}\right)^n$.

(c) We use the union bound. Let A be the event that at least k bins are empty. Notice that there are $m = \binom{n}{k}$ sets of k bins out of the total n bins. Then

$$\mathbb{P}(A) = \mathbb{P}\left(\bigcup_{i=1}^{m} A_i\right) \le \sum_{i=1}^{m} \mathbb{P}(A_i)$$

where A_i is the event that the *i*th set of *k* bins is empty. We know the probability of the first *k* bins being empty from part (b), and this is true for any set of *k* bins, so

$$\mathbb{P}(A_i) = \left(\frac{n-k}{n}\right)^n.$$

Then,

$$\mathbb{P}(A) \le m \cdot \left(\frac{n-k}{n}\right)^n = \binom{n}{k} \left(\frac{n-k}{n}\right)^n.$$

(d) Using Bayes' Rule:

$$\mathbb{P}[2\text{nd bin empty} \mid 1\text{st bin empty}] = \frac{\mathbb{P}[2\text{nd bin empty} \cap 1\text{st bin empty}]}{\mathbb{P}[1\text{st bin empty}]}$$

$$= \frac{(n-2)^n/n^n}{(n-1)^n/n^n}$$

$$= \left(\frac{n-2}{n-1}\right)^n$$
(1)

Alternate solution:

We know bin 1 is empty, so each ball that we throw can land in one of the remaining n-1 bins. We want the probability that bin 2 is empty, which means that each ball cannot land in bin 2 either, leaving n-2 bins. Thus for each ball, the probability that bin 2 is empty given that bin 1 is empty is (n-2)/(n-1). For n total balls, this probability is $[(n-2)/(n-1)]^n$.

- (e) They are dependent. Knowing the latter means the former happens with probability 1.
- (f) In part (c) we calculated the probability that the second bin is empty given that the first bin is empty: $[(n-2)/(n-1)]^n$. The probability that the second bin is empty (without any prior information) is $[(n-1)/n]^n$. Since these probabilities are not equal, the events are dependent.