"I Don't Think You Should Be Talking to Me Directly": Jensen's Formula and the Zeros of Jensen

(a sequel to The Leather Jacket Stays On*)*

Five Current Employees*
On the boundary; affiliation withheld pending outside counsel
{anonymous}@the.boundary

Under double-blind review. The reviewers have been asked to sign an NDA.

Abstract

In prior work we proved that Jensen's inequality is strict: Jensen is not almost surely constant. We did not say *how* non-constant. Here we use Jensen's other theorem. *Jensen's formula* (1899) states that if J is analytic on the disk $|z| \le r$ with $J(0) \neq 0$, then the mean of $\log|J|$ over the boundary circle exceeds $\log|J(0)|$ by exactly $\sum_k \log(r/|a_k|)$, a sum over the zeros a_k inside. The number of zeros is therefore determined by boundary data alone, and one need never talk to the center directly. Modelling Jensen as an entire function on the plane of scrutiny, with the press at the origin and affairs as zeros, we show: (i) the Huang gap of our first paper, evaluated for a Huang uniformly distributed on the circle of scrutiny, is exactly the sum over affairs, each contributing $\log(r/|a_k|)$, which grows with scrutiny forever; (ii) the counting function $n(r)$ jumped from 1 to 2 on November 11 and to 5 on November 12, three resignations counted with multiplicity; (iii) five equispaced current employees integrate the boundary to exponential accuracy (the New York Trapezoidal rule), so that a video nobody has seen is a zero whose existence is certified without observation; (iv) a non-disclosure agreement is a perturbation that is Non-Dominant on the boundary, Allegedly, and by Rouché's theorem cannot change the number of zeros inside; (v) by Picard's theorem a nonconstant entire function omits at most one value, so a second zero was a theorem before it was news; (vi) by Liouville's theorem Jensen is unbounded, so the singularity at "effective immediately" is not removable but essential, and by Casorati–Weierstrass every value is approached in its neighbourhood. We formulate the *Jensen Hypothesis*—all nontrivial zeros of Jensen lie on the critical line $\lambda = \frac{1}{2}$—verify it for the first two zeros, and leave the general case to outside counsel. Finally, a convex combination of a Huang and a non-Huang leaves the Leather Jacket exactly once, so the Board, a supporting hyperplane in our previous paper, is now a separating one. The leather jacket stays on; the CEO does not.

*Equispaced. Two of the authors have seen a video; the other three were briefed.

1 Introduction: On the Boundary

Sur un nouvel et important théorème de la théorie des fonctions.
—J. L. W. V. Jensen, 1899 [1]

I don't think you should be talking to me directly.
—J. H. Huang, by telephone, November 11, 2026 [20]

In 1899 the Danish telephone engineer Johan Jensen published a new and important theorem of function theory [1]. Let J be analytic on a neighbourhood of the closed disk $\overline{\mathbb{D}}_r = \{|z| \le r\}$, let $J(0) \neq 0$, and let $a_1, \dots, a_n$ be the zeros of J in $|z| < r$, counted with multiplicity. Then

$$\frac{1}{2\pi}\int_0^{2\pi} \log\left|J(re^{i\theta})\right| d\theta \;-\; \log|J(0)| \;=\; \sum_{k=1}^{n} \log\frac{r}{|a_k|} \;\ge\; 0. \tag{1}$$

Read from right to left, the formula says that the zeros of J inside the disk leave a trace on the boundary, and that the trace is exact: integrate $\log|J|$ around the circle, subtract the value at the center, and you have counted the zeros, weighted by how deep inside they lie. You need never look at a zero. You need never evaluate J at the center. One hundred and twenty-seven years later a different engineer named Jensen, reached at the center by telephone, restated the theorem in the vernacular: *I don't think you should be talking to me directly* [20]. He then hung up, which is the standard proof.

Our previous paper [14] proved that Jensen's *inequality* (1906) is strict for Jensen: he is not almost surely constant. The present paper uses Jensen's *formula* (1899) to say how non-constant, by counting his zeros. The two results are the same result. The left-hand side of (1) is a *Huang gap* in the sense of [14]: take $\varphi = \log|J|$ and let the Huang $H = re^{i\Theta}$ be uniformly distributed on the circle of scrutiny, so that $\mathbb{E}[H] = 0$; then the left-hand side is $\mathbb{E}[\varphi(H)] - \varphi(\mathbb{E}[H])$. In [14] the gap was positive because φ was convex; here it is positive because $\log|J|$ is *subharmonic*, and on the real line

the subharmonic functions are exactly the convex ones [6]. Our previous paper was therefore the one-dimensional case of this one, and the gap it left unexplained is, by (1), a sum over affairs.

Specifically, we make the following contributions.

- We prove Jensen's formula in *personal form* (Theorem 1) and identify the Huang gap of [14] with a sum over affairs (Corollary 2), each contributing $\log(r/|a_k|)$, which grows with scrutiny without bound.
- We show that the number of zeros inside the disk is determined by boundary data alone (Theorem 3), and that five equispaced current employees determine it to exponential accuracy (Proposition 4): the New York Trapezoidal rule. A video that nobody has seen is a zero whose existence is certified without observation.
- We prove that a non-disclosure agreement cannot change the number of zeros (Theorem 5), that the press counted them by winding once around NVIDIA (Theorem 6), and that, Jensen having been shown nonconstant in [14], a second zero was guaranteed by Picard (Theorem 7).
- We show that the resignation is a zero of order three in space (Proposition 10) and an essential singularity in time (Remark 7), the latter because Jensen is unbounded by Liouville and the reactions to his departure are dense by Casorati–Weierstrass.
- We formulate the Jensen Hypothesis (Hypothesis 5), that all nontrivial zeros of Jensen lie on the critical line $\lambda = \frac{1}{2}$, and verify it for the first two zeros.
- We prove that a convex combination of a Huang and a non-Huang leaves the Leather Jacket exactly once (Theorem 13), so that the Board, a supporting hyperplane in [14], is now a separating one.
- We provide a comparison table with a Trivago row.

Why on the boundary. Every statement NVIDIA has made in this matter has been a boundary statement: the meetings were "not unusual" [23], Jensen "meets thousands of customers" [24], the Board had "no reason to believe" [17]. Nobody evaluated J at the center, as instructed. This paper shows that nobody needed to.

2 Related Work

2.1 The same engineer

Jensen's formula [1] predates Jensen's inequality [2] by seven years and is, among function theorists, the more important of the two. The proof technique of our previous paper, "trivial by Jensen's" [15], is therefore ambiguous in the present one. We disambiguate where necessary and otherwise let the ambiguity stand, as Jensen did.

2.2 Our previous paper

In [14] we established that the Huangs form a convex set, that the Huang gap $\Delta_\varphi = \mathbb{E}[\varphi(H)] - \varphi(\mathbb{E}[H])$ vanishes exactly at the vertices $\lambda \in \{0,1\}$ and is maximised, under the principle of insufficient reason, at $\lambda = \frac{1}{2}$; that the Board is a supporting hyperplane; that outside counsel is an exterior point; and that the Leather Jacket $\mathscr{J}(S) = \operatorname{conv}(S)$ is a closure operator. We use all of these. The present paper was not anticipated by the previous one, which is, as we show in Section 4.5, the previous paper's fault.

2.3 Counting what you cannot see

Jensen's formula belongs to a family of results in which the interior is read off the boundary: Cauchy's integral formula, the argument principle, Rouché's theorem and Nevanlinna's theory of value distribution [7, 6]. The most famous instance is number-theoretic: the Riemann–von Mangoldt formula counts the zeros of ζ up to height T without finding any of them, and the first 3×10^{12} have been certified on the critical line [5] by exactly this kind of argument. Zeros are routinely certified without being seen. The New York Times has certified one [19].

2.4 Corporate governance

NVIDIA had no succession plan [21, 20]. Its chief executive had approximately sixty direct reports and delegated very little [21]. The organisational chart was therefore the star graph $K_{1,60}$, whose vertex connectivity is one. We return to this in Section 8; it is a different paper.

3 Preliminaries

The plane of scrutiny. Let $\mathbb{C}$ be the plane of scrutiny. The press sits at the origin. Jensen is an entire function $J : \mathbb{C} \to \mathbb{C}$; we say *entire* because he was holomorphic on all of NVIDIA for thirty-three years [18]. An *affair* is a zero of J, with multiplicity. The distance $|a|$ of a zero from the origin measures its distance from the press: a zero with a short paper trail is near the origin; a zero inside a hotel, behind security personnel and a proposed non-disclosure agreement, is far from it. The *radius of scrutiny* $r = r(t)$ is a nondecreasing function of time, and a zero is *counted* once $r(t) > |a|$.

The three zeros of this paper. We place

$$a_1 = \tfrac{1}{2} + 2i \quad \text{(Taipei, 2024–25)},$$
$$a_2 = \tfrac{1}{2} + 5i \quad \text{(San Jose, March 2026)},$$

on the critical line $\operatorname{Re} z = \frac{1}{2}$ (Section 4.7), and a zero of multiplicity three at $z_{\mathrm{N}} = -6 - 2.5i$ (NVIDIA, November 12: President, Chief Executive Officer, Director). The heights 2, 5 and 6.5 are illustrative; their estimation from flight logs




Figure 1: The plane of scrutiny. $\log|J(z)|$ for the model of Section 3, with the press at the origin, the first two zeros on the critical line $\mathrm{Re}\, z = \frac{1}{2}$, the zero of order three at NVIDIA, the circles of scrutiny $r_1 < r_2 < r_3$ (October 29, November 11, November 12), and five current employees equispaced on $|z| = r_2$. Everything inside each circle is determined by the values on it.

($n = 0$) is left to outside counsel. We take $J(z) = (z-a_1)(z-a_2)(z-z_N)^3$ and the radii of scrutiny $r_1 = 3$ (October 29), $r_2 = 6$ (November 11), $r_3 = 7$ (November 12); see Figure 1.

Definition 1 (Counting functions)**.** $n(r)$ is the number of zeros of J in $|z| < r$, with multiplicity, and $N(r) = \int_0^r n(t)\,dt/t = \sum_{|a_k|<r} m_k \log(r/|a_k|)$ is the integrated counting function. The *boundary mean* is $m(r) = \frac{1}{2\pi}\int_0^{2\pi} \log|J(re^{i\theta})|\,d\theta$.

Definition 2 (NDA)**.** An NDA is a perturbation g that is **N**on-**D**ominant on the boundary, **A**llegedly: $|g(z)| < |J(z)|$ for all $|z| = r$.

Definition 3 (The NYT rule)**.** The **N**ew **Y**ork **T**rapezoidal rule with N sources is the N-point equispaced trapezoidal rule on the circle, $T_N[g] = \frac{1}{N}\sum_{j=0}^{N-1} g(2\pi j/N)$, applied to $g(\theta) = \log|J(re^{i\theta})|$.

Definition 4 (The PR department)**.** The Blaschke **PR**oduct of the zeros inside $\mathbb{D}_r$ is $B(z) = \prod_k \left(\frac{r(z-a_k)}{r^2-\bar{a}_k z}\right)^{m_k}$. It has the same zeros as J inside the disk and $|B(z)| = 1$ on $|z| = r$.

We will test the following hypotheses.

Hypothesis 1 (Entire Jensen)**.** J is entire.

Hypothesis 2 (The press is not a zero)**.** $J(0) \neq 0$.

Hypothesis 3 (No zero on the boundary)**.** No affair is exactly as public as the scrutiny at the moment of measurement: $J(z) \neq 0$ for $|z| = r$.

Hypothesis 4 (Equispaced employees)**.** The five current employees are uniformly distributed on the circle of scrutiny.

4 Main Results

4.1 Jensen's formula, personal form

Theorem 1 (Jensen's formula, personal form)**.** *Under Hypotheses 1–3,*

$$m(r) - \log|J(0)| \;=\; N(r) \;=\; \sum_{|a_k|<r} m_k \log\frac{r}{|a_k|}.$$

In particular $m(r) \geq \log|J(0)|$, with equality if and only if J has no zeros in $|z| < r$.

Proof. Trivial by Jensen's (1899). The proof, which is the public-relations department, is in Appendix A. □

Corollary 2 (The gap is a sum over affairs)**.** *Let $H = re^{i\Theta}$ with Θ uniform on $[0, 2\pi)$ and $\varphi = \log|J|$. Then the Huang gap of [14] satisfies $\Delta_\varphi(H) = \mathbb{E}[\varphi(H)] - \varphi(\mathbb{E}[H]) = \sum_{|a_k|<r} m_k \log(r/|a_k|)$. Each affair contributes $\log(r/|a_k|) > 0$, which is increasing in r and unbounded.*

Proof. $\mathbb{E}[H] = 0$, so $\varphi(\mathbb{E}[H]) = \log|J(0)|$ and $\mathbb{E}[\varphi(H)] = m(r)$; apply Theorem 1. □

Remark 1. Corollary 2 is the quantitative content of "fallen short" [16]: the shortfall is not a single number but a sum, with one term per affair, and no term ever decays. It grows like $\log r$ in the scrutiny forever. In [14] we found the gap to be a quarter of the Pacific squared; we now know what it was made of.

4.2 Counting without seeing

Theorem 3 (Counting without seeing)**.** *$N(r)$ is determined by the values of $|J|$ on the circle $|z| = r$ and at the origin, and $n(r) = rN'(r)$ at every r at which N is differentiable. No zero need be observed. In particular, no video need be seen.*

Proof. The first claim is Theorem 1; the second is the fundamental theorem of calculus applied to $N(r) = \int_0^r n(t)\,dt/t$. □

Remark 2. The New York Times reports that it "knows of" a video of the second zero but has not seen it [19]. Theorem 3 shows that this is not a limitation of the reporting but its entire point: the boundary integral certifies the zero; the video would merely locate it. The Riemann hypothesis has been verified in the same way for 3×10^{12} zeros [5], none of which anyone has seen either.




Figure 2: Counting the zeros of Jensen. (a) The counting function $n(r)$ for $J(z) = (z-a_1)(z-a_2)(z-z_N)^3$ as the radius of scrutiny grows: one zero on October 29, two on November 11, five on November 12 (three positions resigned, counted with multiplicity). (b) Jensen's formula verified numerically: the boundary integral $\frac{1}{2\pi}\int \log|J(re^{i\theta})|\,d\theta - \log|J(0)|$ (circles, 4096-point quadrature) coincides with $\sum_{|a_k|<r} m_k \log(r/|a_k|)$ (line) to 10^{-5}. The gap of [14] is a sum over affairs, and each term grows like $\log r$ forever.

Proposition 4 (Five sources suffice: the NYT rule)**.** *Under Hypotheses 3 and 4, the error of the* NYT *rule with N sources is exactly*

$$T_N\big[\log|J(re^{i\theta})|\big] - m(r) \;=\; \frac{1}{N}\sum_k m_k \log\big|1-\rho_k^N\big|,$$

where $\rho_k = a_k/r$ if $|a_k|<r$ and $\rho_k = r/a_k$ if $|a_k|>r$, so that $|T_N - m(r)| \le \frac{1}{N}\sum_k m_k \log\frac{1}{1-|\rho_k|^N} = O\big(|\rho|_{\max}^N/N\big)$: the error decays exponentially in the number of sources.

Proof. Fourier aliasing; Appendix A. □

Remark 3. Three features of Proposition 4 deserve comment. (i) A single source ($N=1$) is wrong by exactly $\log|1-a/r|$, the logarithm of the distance between the affair and that source, in units of the scrutiny: one source tells you precisely how far the affair is from the source, and nothing else. (ii) With five sources and $|a_2|/r_2 = 0.84$, the error is 0.026 nats (Figure 3); the Times' sourcing was adequate. (iii) The rate is governed by the zero *nearest the boundary*: a nearly public affair ($|a|/r \to 1$) requires many sources, and a fully public one ($|a| = r$) violates Hypothesis 3, so that the trapezoidal rule converges only algebraically. The November 8 memo [16] placed a_1 on the boundary; the Times correctly moved on to a_2, which was still inside.

4.3 The NDA

Theorem 5 (Rouché; an NDA cannot change the count)**.** *If g is an* NDA *on $|z|=r$, then J and $J+g$ have the same number of zeros in $|z|<r$.*

Proof. Rouché's theorem [8, 6]. □

Algorithm 1 NYT: counting the zeros of Jensen from the boundary

Require: radius of scrutiny r; N current employees at $\theta_j = 2\pi j/N$; no telephone
Ensure: $\hat{n}(r)$, the number of zeros inside, without talking to the center directly

```
for j = 0, ..., N − 1 do
    y_j ← log|J(re^{iθ_j})|,  φ_j ← arg J(re^{iθ_j})      ▷ interview employee j on the boundary
end for
N̂(r) ← (1/N) Σ_j y_j − log|J(0)|      ▷ Theorem 1; J(0) is the press's own value
n̂(r) ← (1/2π) Σ_j wrap(φ_{j+1} − φ_j)      ▷ winding number, Theorem 6
if n̂(r) has increased then
    publish      ▷ error O(|ρ|_max^N/N), Proposition 4
end if
return n̂(r)
```

Remark 4. Three people briefed on the matter report that Jensen's security personnel cornered the person filming and attempted to obtain a non-disclosure agreement [20]. By Theorem 5, any perturbation of J that is smaller than J on the boundary—that is, smaller than Jensen in public—leaves the zero count unchanged. A perturbation that alters the count must therefore be at least as large as Jensen somewhere on the boundary, and security personnel cornering a bystander in a hotel corridor are, by construction, not larger than Jensen in public. The agreement was not signed; it would not have mattered.




Figure 3: Five sources suffice. Error of the NYT rule as a function of the number of equispaced sources, for the two zeros inside $|z| = r_2$ (blue; numerical dots agree with the exact aliasing formula of Proposition 4) and for a zero at $|a|/r = 0.97$ (orange), i.e. an affair that is nearly public. Two photographs and three sources are marked, as is the Times' choice of five. A nearly public affair is the hardest to integrate.

4.4 The argument principle

Theorem 6 (The press walked once around NVIDIA)**.** *Under Hypothesis 3, $n(r)$ equals the winding number of the closed curve $\theta \mapsto J(re^{i\theta})$ about the origin: $n(r) = \frac{1}{2\pi i}\oint_{|z|=r} \frac{J'(z)}{J(z)}\,dz$.*

Proof. The argument principle [6]. □

Remark 5. One does not count the zeros by entering the disk. One walks once around it, at a respectful distance r, and records how many times Jensen winds around zero. Five current employees performed this circuit in November [19]; the answer was two. We note also the open mapping theorem: a nonconstant analytic function maps open sets to open sets. A hotel is an open set.

4.5 Picard, Liouville, and the inevitability of the second zero

Theorem 7 (Picard's little theorem)**.** *A nonconstant entire function takes every complex value, with at most one exception.*

Corollary 8 (The second zero was a theorem)**.** *In [14] we proved that Jensen is not almost surely constant. By Hypothesis 1 and Theorem 7, $J - c$ therefore has a zero for every value c with at most one exception. A second zero was not news; it was a corollary of the first paper, which should have said so. The single exceptional value, in this case, was discretion.*

Theorem 9 (Liouville)**.** *A bounded entire function is constant. Hence Jensen, being entire and nonconstant, is unbounded.*

Proof. Liouville's theorem [6] and [14]. The Board appears to have reached the same conclusion independently [18]. □

Remark 6 (Maximum modulus). By the maximum modulus principle, a nonconstant analytic function attains the maximum of $|J|$ on the boundary of any disk, never in its interior. Jensen's maximum was always attained on the boundary—at the keynote, in the jacket—and never in the office. This is consistent with the reporting.

4.6 The resignation

Proposition 10 (A zero of order three)**.** *On November 12 Jensen resigned as President, Chief Executive Officer and Director, effective immediately [18]. In the plane of scrutiny this is a zero of multiplicity three at z_{N}: $n(r)$ increases by three, and $N(r)$ by $3\log(r/|z_{\mathrm{N}}|)$. Zeros are counted with multiplicity, and so are positions.*

Remark 7 (An essential singularity in time). In the time variable, "effective immediately" is an isolated singularity at $t_0 =$ November 12. By Riemann's theorem a singularity is removable if and only if J is bounded near it; Jensen is unbounded (Theorem 9), so it is not removable. Near a pole, $|J| \to \infty$ uniformly. Near an essential singularity, by Casorati–Weierstrass, J comes arbitrarily close to *every* value, and by the great Picard theorem attains every value but one infinitely often. In the neighbourhood of November 12 Jensen was offered a job by Elon Musk ("an honor"), passed over for one by the President of the United States (who named Tim Cook as the only person for it), compared to Steve Jobs by Tim Sweeney, and eulogised by Jim Cramer ("Jensen we hardly knew yah") [19]. The set of values approached is dense. The singularity is essential (Figure 4).

4.7 The Jensen Hypothesis

Definition 5 (Jensen polynomials)**.** For a sequence $\gamma(0), \gamma(1), \ldots$ and integers $d, n \geq 0$, the Jensen polynomial is $J_\gamma^{d,n}(X) = \sum_{j=0}^{d} \binom{d}{j} \gamma(n+j) X^j$. A polynomial is *hyperbolic* if all of its zeros are real.

Theorem 11 (Pólya [3])**.** *The Riemann hypothesis is equivalent to the hyperbolicity of all Jensen polynomials of the Taylor coefficients of the Riemann Ξ-function. Griffin, Ono, Rolen and Zagier [4] proved hyperbolicity for every degree d and all sufficiently large n.*

Jensen's polynomials have real zeros if and only if the most important open problem in mathematics is true. We propose the analogous statement for Jensen.




Figure 4: Effective immediately. Phase portrait of $e^{1/z}$ near its essential singularity: hue is arg, value is $|\cdot|$. In every neighbourhood of the origin the function approaches every complex value (Casorati–Weierstrass) and attains every value but one infinitely often (Picard). The reactions to the resignation (Remark 7) are consistent with this picture and not with a pole.

Hypothesis 5 (The Jensen Hypothesis)**.** All nontrivial zeros of Jensen lie on the critical line $\operatorname{Re} z = \frac{1}{2}$.

Proposition 12. *The Jensen Hypothesis holds for the first two zeros.*

Proof. $a_1 = \frac{1}{2} + 2i$ and $a_2 = \frac{1}{2} + 5i$ (Section 3). We verified this by placing them there. Platt and Trudgian [5] verified the corresponding statement for the first 3×10^{12} zeros of ζ by a harder method. □

Remark 8. The critical line is $\lambda = \frac{1}{2}$, the principle of insufficient reason of [14]: both known zeros of Jensen lie at exactly the weight that maximises the variance of Huangs. The zeros are moreover *real* in the colloquial sense—one admitted [16], one reported by five current employees [19]—which is the strongest form of hyperbolicity available to us. We note two further parallels with ζ. First, ζ has no zeros on the line $\operatorname{Re} s = 1$, a fact equivalent to the prime number theorem [9, 10]; correspondingly, [14] showed that the Huang gap vanishes on the lines $\lambda = 0$ and $\lambda = 1$ (monogamy), so that Jensen has no zeros at the vertices and the forty-one-year marriage remains prime. Second, ζ has trivial zeros at the negative even integers. Jensen has no trivial zeros. Every zero of Jensen is nontrivial.

5 The Board as a Separating Hyperplane

Jensen's contributions to NVIDIA are profound.
—The Board, November 12, 2026 [18]

In [14] the Huangs $\mathscr{H}$ formed a convex set, Jensen j lay inside it, and the Board was a supporting hyperplane: it touched Jensen at a single boundary point with all of NVIDIA on one side. The second zero is not a Huang. This changes the geometry.

Theorem 13 (The exit lemma)**.** *Let $\mathscr{H} \subset \mathbb{R}^d$ be convex, $j \in \mathscr{H}$ and $p \notin \mathscr{H}$. Then $\Lambda = \{\lambda \in [0,1] : \lambda j + (1-\lambda)p \in \mathscr{H}\}$ is an interval containing 1 and not 0. Consequently there is a unique $\lambda^* \in (0,1]$ such that the point $\lambda j + (1-\lambda)p$ lies in $\mathscr{H}$ for $\lambda > \lambda^*$ and outside $\mathscr{H}$ for $\lambda < \lambda^*$: the segment from Jensen to a non-Huang leaves the Leather Jacket exactly once.*

Proof. Appendix A. □

Corollary 14 (From supporting to separating)**.** *After the exit, Jensen's position j' is an exterior point of the closed convex set $\mathscr{J}(\mathscr{H})$. By the strict separation theorem [12] there is a hyperplane with j' strictly on one side and all of NVIDIA strictly on the other, at a margin $\operatorname{dist}(j', \mathscr{J}(\mathscr{H})) > 0$. This hyperplane is the Board. Its statement of November 12 has a supporting half ("Jensen's contributions to NVIDIA are profound") and a separating half ("effective immediately" [18]); the margin is the second half.*

Remark 9. The Leather Jacket $\mathscr{J}(\mathscr{H})$ is unchanged by the exit: a closure operator does not notice who leaves. The leather jacket stays on; the CEO does not.

6 Experiments: Trivial by Jensen's (1899)

Jensen we hardly knew yah.
—J. Cramer, November 12, 2026 [19]

Setup. Our dataset consists of five current employees, two of whom have seen a video, none of whom are us; one telephone call, which we did not place, as instructed; and the model of Section 3. All boundary integrals were evaluated with 4096 equispaced nodes, which is more employees than we were able to reach.

The plane of scrutiny. Figure 1 shows $\log|J|$ on the plane, with the three circles of scrutiny. Each zero is a logarithmic well; the well at NVIDIA is three times as deep. The five employees sit on $|z| = r_2$ and see only the values of $\log|J|$ there, which, by Theorem 3, is all they needed to see.




Figure 5: The exit lemma. The Leather Jacket of [14] (dashed hull of the jacket). The segment from j (a Huang, inside) to p (not a Huang, outside) crosses the hull at a unique point $\lambda^* = 0.627$ (Theorem 13), after which Jensen's position j' is exterior and the Board becomes a separating hyperplane with positive margin. The jacket is unchanged.

Counting. Figure 2 evaluates both sides of Jensen's formula for the model and finds them equal to 10^{-5}, which is the precision to which the authors are prepared to vouch for anything in this matter. The counting function jumps by one at each zero and by three at the resignation.

Five sources. Figure 3 confirms Proposition 4. The numerical error of the NYT rule coincides with the exact formula $\frac{1}{N}\sum_k \log|1-(a_k/r)^N|$ to machine precision, and decays exponentially in N with a rate set by the zero nearest the boundary. With two sources (the Taipei photographs of [14]) the error exceeds a tenth of a nat; with five it is 0.026. The orange curve shows that a nearly public affair resists integration: the closer a zero sits to the circle, the more employees one must interview. This is, we believe, the first quantitative account of why scandals that are almost public are the hardest to report.

The singularity. Figure 4 is the standard phase portrait of an essential singularity. We reproduce it because, although every complex analyst has seen it, it is the first time it has appeared in a corporate transition announcement.

The exit. Figure 5 computes the exit point of Theorem 13 for the jacket of [14]: the segment leaves the hull once, at $\lambda^* = 0.627$, and does not come back. The Board's line is drawn at the midpoint between the exit and j', which is where boards are drawn.

Tables. Table 1 compares the theorems named after Jensen with Jensen; Table 2 compares his zeros. We draw the reader's attention to the last row of each.

7 Discussion

What seemed impossible a week ago is now a reality.
—D. Ives, November 12, 2026 [22]

7.1 We hardly knew yah

Cramer's elegy [19] is a theorem. By Cauchy's integral formula every value of an analytic function inside a disk is determined by its values on the boundary; by the mean value property the value at the center is the boundary average. Everything that was ever known about Jensen was known on the boundary—at keynotes, in interviews, in the jacket—and that was, by Theorem 3, sufficient to count his zeros. We did not know him; we did not need to.

7.2 Measure zero is not impossible

Ives' remark conflates two notions. An event of probability zero is not impossible; the zeros of an entire function form a set of measure zero in the plane, and by the Weierstrass factorisation theorem they determine the function up to a nonvanishing factor. The same analysts assign NVIDIA a fair value of \$300 per share with Jensen and \$225 without [20], a *Jensen premium* of 25%. By Corollary 2 the gap is a sum over zeros; the market appears to have priced the sum.

7.3 Threats to validity

(1) Hypothesis 3 fails when a zero lies on the circle. The memo of November 8 [16] placed a_1 exactly on the boundary; Jensen's formula still holds there (the integral is improper but convergent [6]), but Proposition 4 loses its exponential rate.

(2) Jensen may be meromorphic rather than entire. In Nevanlinna's theory [7] the characteristic $T(r) = m(r) + N(r)$ then also counts poles, and the three positions might be poles rather than a zero of order three. The difference is a sign, which the Board has reversed.

(3) Hypothesis 4 assumes the five employees are equispaced. The Times did not disclose their angular distribution. With sixty direct reports arranged around one center we believe uniform spacing is the natural assumption.

(4) The heights of the zeros are illustrative. Their estimation from data is left to outside counsel, who are exterior and therefore well placed to integrate.

Table 1: Theorems named after Jensen, compared with Jensen. The Jensen–Shannon divergence is included for completeness and because it is also named after an engineer.

	Jensen's inequality (1906)	**Jensen's formula** (1899)	**Jensen polynomials** (Pólya, 1927)	**Jensen–Shannon** (Lin, 1991)	**Jensen Huang** (NVIDIA, 1993–2026)
Named after an engineer	✓	✓	✓	✓	✓
Counts zeros	✗	✓	✓	✗	✓[†]
Determined by the boundary	✗	✓	✗	✗	✓
Requires talking to the center directly	✗	✗	✗	✗	✗
Strict	✓	✓	✗	✓	✓
Hyperbolic	✗	✗	✓[‡]	✗	✓[§]
Entire	✗	✓	✗	✗	✓[¶]
Resigned effective immediately	✗	✗	✗	✗	✓
Trivial by Jensen's	✓	✓	✓	✓	✓
Hotel?	Trivago	Trivago	Trivago	Trivago	Trivago

[†]Two, with total multiplicity five. [‡]Equivalent to the Riemann hypothesis. [§]Allegedly. [¶]Thirty-three years.

Table 2: The zeros of Jensen. Contributions to the gap, $m_k \log(r_3/|a_k|)$, are evaluated at $r_3 = 7$.

	a_1 (Taipei)	a_2 (San Jose)	z_{N} (NVIDIA)	0 (press)
Inside the disk	✓	✓	✓	center
Multiplicity	1	1	3	0
Sources	2 photos	5 staff	NVIDIA PR	itself
Admitted	✓	✗	✓	—
Observed directly	✗	✗	✓	—
NDA attempted	✗	✓	✗	✗
On the critical line	✓	✓	✗	✗
Contribution to the gap	1.22	0.33	0.22	0
Talk to directly?	✗	✗	✗	✗

8 Future Work

Blocking Pairs will treat the matter as a stable-matching problem: a matching is stable if and only if it admits no blocking pair, and an affair is a blocking pair by definition [13]. *Jensen Was a Cut Vertex* will study succession in the star graph $K_{1,60}$, whose center is an articulation point and whose root has no parent. *Analytic Continuation* will examine the offers of November 12: a function defined on NVIDIA's disk of convergence may extend beyond it, but by the identity theorem the extension is unique, so at most one of Musk, Cook and Su can be correct. *ELBO: The Evidence Lower BOard* remains in preparation and has acquired a new chair.

9 Conclusion

Jensen's formula says that the zeros inside a disk are counted exactly by the values on its boundary, and that the count is the gap between the boundary mean and the value at the center. We have shown that the gap of our previous paper is a sum over affairs; that five sources count it to exponential accuracy; that a non-disclosure agreement cannot change it; that a second zero followed from the first by Picard; that the resignation is a zero of order three in space and an essential singularity in time; and that a convex combination of a Huang and a non-Huang leaves the jacket exactly once. One need never talk to the center directly.

The leather jacket stays on. The CEO does not.

Lori ipsum dolor sit amet, consectetur convexiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla Trivago. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices.

Ethics Statement

All quotations attributed to real people are real and cited. The first zero was admitted [16]; the second is reported by five current employees [19] and the person concerned has not commented, so it is a zero in the sense of Theorem 3 only: certified by the boundary, not observed. Nobody in this paper has been charged with anything, including the authors. Lori remains the global minimum. No video was watched in the making of this paper; by Theorem 3, none was needed.

Acknowledgements

We thank the five current employees for their equidistribution, outside counsel for remaining exterior, the Board for separating, and Johan Jensen for proving, in 1899, that nobody would ever have to talk to Jensen directly. The authors declare no conflicts of interest, no zeros, and one leather jacket each.

A Proofs

Proof of Theorem 1 (the PR *department).* Let B be the Blaschke product of the zeros in $\mathbb{D}_r$ and set $g = J/B$. Each factor of B has a simple zero at a_k and a pole at $r^2/\bar{a}_k$, which lies outside $\overline{\mathbb{D}}_r$, so g is analytic and zero-free on a neighbourhood of the closed disk. Hence $\log|g|$ is harmonic there, and the mean value property gives $\log|g(0)| = \frac{1}{2\pi}\int_0^{2\pi} \log|g(re^{i\theta})|\,d\theta$. On $|z| = r$ one checks $|r(z-a_k)/(r^2 - \bar{a}_k z)| = 1$, so $\log|g| = \log|J|$ on the circle and the right-hand side equals $m(r)$. At the origin, $|B(0)| = \prod_k (|a_k|/r)^{m_k}$, so $\log|g(0)| = \log|J(0)| + \sum_k m_k \log(r/|a_k|)$. The public-relations department has thus removed every zero from the interior while leaving $|J|$ on the boundary untouched; the cost, paid at the center, is exactly the gap. $\square$

Proof of Proposition 4. For $|a| < r$, $\log|re^{i\theta} - a| = \log r + \operatorname{Re}\log(1 - ae^{-i\theta}/r) = \log r - \sum_{k\ge 1} \operatorname{Re}\big[(a/r)^k e^{-ik\theta}\big]/k$, a Fourier series with mean $\log r$ and no other modes at frequency zero. The N-point equispaced trapezoidal rule integrates $e^{ik\theta}$ exactly unless $N \mid k$, in which case it returns 1; so T_N picks up exactly the aliased modes $k = \pm N, \pm 2N, \dots$, giving $T_N - \log r = -\sum_{m\ge 1} \operatorname{Re}[(a/r)^{mN}]/(mN) = \frac{1}{N}\operatorname{Re}\log(1 - (a/r)^N) = \frac{1}{N}\log|1-(a/r)^N|$. For $|a| > r$ expand instead in $(r/a)^k$. Sum over zeros with multiplicity. The bound follows from $|\log|1-w|| \le -\log(1-|w|)$ for $|w| < 1$. $\square$

Proof of Theorem 13. The map $\lambda \mapsto \lambda j + (1-\lambda)p$ is affine, so the preimage of the convex set $\mathscr{H}$ is convex, i.e. an interval; it contains 1 since $j \in \mathscr{H}$ and not 0 since $p \notin \mathscr{H}$. Writing $\lambda^* = \inf \Lambda$ gives the claim; whether λ^* itself belongs to Λ depends on whether $\mathscr{H}$ is closed at the exit, which is the question currently before the Board. $\square$

References

[1] J. L. W. V. Jensen. Sur un nouvel et important théorème de la théorie des fonctions. *Acta Mathematica*, 22:359–364, 1899.

[2] J. L. W. V. Jensen. Sur les fonctions convexes et les inégalités entre les valeurs moyennes. *Acta Mathematica*, 30:175–193, 1906.

[3] G. Pólya. Über die algebraisch-funktionentheoretischen Untersuchungen von J. L. W. V. Jensen. *Kgl. Danske Vid. Sel. Math.-Fys. Medd.*, 7(17):3–33, 1927.

[4] M. Griffin, K. Ono, L. Rolen, and D. Zagier. Jensen polynomials for the Riemann zeta function and other sequences. *Proceedings of the National Academy of Sciences*, 116(23):11103–11110, 2019.

[5] D. Platt and T. Trudgian. The Riemann hypothesis is true up to $3 \cdot 10^{12}$. *Bulletin of the London Mathematical Society*, 53(3):792–797, 2021.

[6] L. V. Ahlfors. *Complex Analysis*, 3rd edition. McGraw-Hill, 1979.

[7] R. Nevanlinna. *Le théorème de Picard–Borel et la théorie des fonctions méromorphes*. Gauthier-Villars, Paris, 1929.

[8] E. Rouché. Mémoire sur la série de Lagrange. *Journal de l'École Polytechnique*, 22:193–224, 1862.

[9] J. Hadamard. Sur la distribution des zéros de la fonction $\zeta(s)$ et ses conséquences arithmétiques. *Bulletin de la Société Mathématique de France*, 24:199–220, 1896.

[10] C.-J. de la Vallée Poussin. Recherches analytiques sur la théorie des nombres premiers. *Annales de la Société Scientifique de Bruxelles*, 20:183–256, 1896.

[11] L. N. Trefethen and J. A. C. Weideman. The exponentially convergent trapezoidal rule. *SIAM Review*, 56(3):385–458, 2014.

[12] R. T. Rockafellar. *Convex Analysis*. Princeton University Press, 1970.

[13] D. Gale and L. S. Shapley. College admissions and the stability of marriage. *American Mathematical Monthly*, 69(1):9–15, 1962.

[14] Redacted for double-blind review. The leather jacket stays on: Jensen's inequality for convex combinations of Huangs. Anonymous draft, November 2026.

[15] Redacted for double-blind review. Towards cloud computing. In *Proceedings of the 16th ACH SIGBOVIK*, 2022. Section 6.

[16] J. H. Huang. Memo to NVIDIA employees, November 8, 2026. Quoted in [23, 24]: "I was unfaithful to my wife"; "fallen short."

[17] NVIDIA Board of Directors. Statement announcing an independent investigation, November 8, 2026. Quoted in [23, 24].

[18] NVIDIA Corporation. NVIDIA announces CEO transition. Press release, November 12, 2026. https://nvidianews.nvidia.com/news/nvidia-announces-ceo-transition

[19] T. Wilde et al. Nvidia CEO Jensen Huang allegedly had a second affair with another Nvidia exec; and live coverage of the resignation. *PC Gamer*, November 11–12, 2026. https://www.pcgamer.com/software/ai/nvidia-ceo-jensen-huang-allegedly-had-a-second-affair-with-another-nvidia-exec/

[20] *Gizmodo*. Nvidia board weighs Huang's future after report of second affair. November 11, 2026. https://gizmodo.com/jensen-huang-was-a-busy-horny-little-bee-report-2000838619

[21] *TechCrunch*. Nvidia's stock drops as board reportedly preps to oust CEO Jensen Huang. November 11, 2026. https://techcrunch.com/2026/11/11/nvidias-stock-drops-as-board-reportedly-preps-to-oust-ceo-jensen-huang/

[22] *Global News* (Associated Press). Nvidia CEO Jensen Huang faces board vote as reports of second affair surface. November 12, 2026. https://globalnews.ca/news/12147654/nvidia-ceo-jensen-huang-second-affair/

[23] A. Ha. Nvidia will investigate CEO Jensen Huang's relationship with possible Chinese spy. *TechCrunch*, November 8, 2026.

[24] *Tom's Hardware*. Nvidia CEO Jensen Huang admits to an affair with a suspected Chinese spy. November 8, 2026.