

The Importance of the Microbiota and Individual Community Members in Colorectal Cancer

Marc A Sze¹ and Patrick D Schloss^{1†}

† To whom correspondence should be addressed: pschloss@umich.edu

¹ Department of Microbiology and Immunology, University of Michigan, Ann Arbor, MI

Co-author e-mails:

- marcsze@med.umich.edu

1 **Abstract**

2 **Background.**

3 **Results.**

4 **Conclusions.**

5 **Keywords**

6 microbiota; colorectal cancer; polyps; treatment; risk factor.

7 Background

8 **Results**

9 ***Section 1***

10 ***Section 2***

11 ***Section 3***

12 ***Section 4***

¹³ **Discussion**

¹⁴ **Conclusion**

Methods

Obtaining Data Sets: Studies used for this meta-analysis were identified through the review articles written by Keku, et al. and Vogtmann, et al. [1,2]. All studies were included that used tissue or feces as their sample source for 16S rRNA gene sequencing analysis. Studies using either 454 or Illumina sequencing technology were included. Only data sets that had the raw sequences available for analysis were included. Some studies did not have publically available raw sequences or did not have meta data in which the authors were able to share. After this filtering step the following studies remained: Ahn [3], Baxter [4], Brim [5], Burns [6], Chen [7], Dejea [8], Flemer [9], Geng [10], Hale [11], Kostic [12], Lu [13], Sanapareddy [14], Wang [15], Weir [16], and Zeller [17]. The Zackular [18] study was not included because the 90 individuals analyzed within the study are contained within the larger Baxter study. The Kostic study was not used since after sequence processing all the case samples did not have more than 100 sequences remaining. This left a total of 13 studies in which complete analysis could be completed.

Data Set Breakdown: In total there were 7 studies with only fecal samples (Ahn, Baxter, Brim, Hale, Wang, Weir, and Zeller), 5 studies with only tissue samples (Burns, Dejea, Geng, Lu, Sanapareddy), and 2 studies with both fecal and tissue samples (Chen and Flemer). The total number of individuals initially run through the sequence processing for the fecal samples was 1899 and for the tissue samples was 462.

Sequence Processing: For the majority of studies raw sequences were downloaded from the SRA (<ftp://ftp-trace.ncbi.nih.gov/sra/sra-instant/reads/ByStudy/sra/SRP/>) and metadata was obtained from the following website: <http://www.ncbi.nlm.nih.gov/Traces/study/> by searching the respective accession number of the study. Of the studies that did not have sequences and meta data on the SRA one study had the data stored on DBGap [3] and four studies the data was obtained directly from the authors [9,11,14,16]. Each

study was processed using the mothur (v1.39.3) software program [19]. Where possible quality filtering utilized the default methods used in mothur for either 454 or Illumina based sequencing. If it was not possible to use these defaults the author stated quality cut-offs were used instead. Chimeras were identified and removed using the VSEARCH [20] program and *de novo* OTU clustering at 97% similarity using the OptiClust algorithm [21] was utilized.

Statistical Analysis. All statistical analysis after sequence processing utilized the R software package (v3.4.2). For the alpha diversity analysis values were power transformed using the rcompanion (v1.10.1) package and then Z-score normalized using the car (v2.1.5) package. Testing for alpha diversity differences utilized both T-tests and ANOVA. Linear mixed-effect models were created using the lme4 (v1.1.14) package to correct for both study and variable region effect in the diversity measures when analyzing colorectal cancer groups. Relative Risk was analyzed using both the epiR (v0.9.87) and metafor (v2.0.0) packages. Relative risk significance testing utilized the chi-squared test. Beta-diversity differences utilized a Bray-Curtis distance matrix and PERMANOVA executed with the vegan (v2.4.4) package. Random Forest models were built using both the caret (v6.0.77) and randomForest (v4.6.12) packages. Random Forest testing of the obtained AUC versus a random model AUC utilized T-tests. Power analysis and estimations were made using the pwr (v1.2.1) and statmod (v1.4.30) packages. All figures were created using both ggplot2 (v2.2.1) and gridExtra (v2.3) packages.

Reproducible Methods. The code and analysis can be found here https://github.com/SchlossLab/Sze_CRCMetaAnalysis_XXXX_2016. Unless mentioned otherwise the accession number for the raw sequences for the studies used in this analysis can be found directly in the respective batch file, on the GitHub repository or in the original manuscript.

64 **Declarations**

65 **Ethics approval and consent to participate**

66 Need to fill in.

67 **Consent for publication**

68 Not applicable.

69 **Availability of data and material**

70 Need to fill in.

71 **Competing Interests**

72 All authors declare that they do not have any relevant competing interests to report.

73 **Funding**

74 Need to fill in.

75 **Authors' contributions**

76 Need to fill in.

⁷⁷ **Acknowledgements**

⁷⁸ Need to fill in.

References

1. Keku TO, Dulal S, Deveau A, Jovov B, Han X. The gastrointestinal microbiota and colorectal cancer. *American Journal of Physiology - Gastrointestinal and Liver Physiology* [Internet]. 2015 [cited 2017 Oct 30];308:G351–63. Available from: <http://ajpgi.physiology.org/lookup/doi/10.1152/ajpgi.00360.2012>
2. Vogtmann E, Goedert JJ. Epidemiologic studies of the human microbiome and cancer. *British Journal of Cancer* [Internet]. 2016 [cited 2017 Oct 30];114:237–42. Available from: <http://www.nature.com/doi/10.1038/bjc.2015.465>
3. Ahn J, Sinha R, Pei Z, Dominianni C, Wu J, Shi J, et al. Human gut microbiome and risk for colorectal cancer. *Journal of the National Cancer Institute*. 2013;105:1907–11.
4. Baxter NT, Ruffin MT, Rogers MAM, Schloss PD. Microbiota-based model improves the sensitivity of fecal immunochemical test for detecting colonic lesions. *Genome Medicine*. 2016;8:37.
5. Brim H, Yooseph S, Zoetendal EG, Lee E, Torralbo M, Laiyemo AO, et al. Microbiome analysis of stool samples from African Americans with colon polyps. *PloS One*. 2013;8:e81352.
6. Burns MB, Lynch J, Starr TK, Knights D, Blekhman R. Virulence genes are a signature of the microbiome in the colorectal tumor microenvironment. *Genome Medicine*. 2015;7:55.
7. Chen W, Liu F, Ling Z, Tong X, Xiang C. Human intestinal lumen and mucosa-associated microbiota in patients with colorectal cancer. *PloS One*. 2012;7:e39743.
8. Dejea CM, Wick EC, Hechenbleikner EM, White JR, Mark Welch JL, Rossetti BJ, et al. Microbiota organization is a distinct feature of proximal colorectal cancers. *Proceedings of*

the National Academy of Sciences of the United States of America. 2014;111:18321–6.

9. Flemer B, Lynch DB, Brown JMR, Jeffery IB, Ryan FJ, Claesson MJ, et al. Tumour-associated and non-tumour-associated microbiota in colorectal cancer. *Gut*. 2017;66:633–43.

10. Geng J, Fan H, Tang X, Zhai H, Zhang Z. Diversified pattern of the human colorectal cancer microbiome. *Gut Pathogens*. 2013;5:2.

11. Hale VL, Chen J, Johnson S, Harrington SC, Yab TC, Smyrk TC, et al. Shifts in the Fecal Microbiota Associated with Adenomatous Polyps. *Cancer Epidemiology, Biomarkers & Prevention: A Publication of the American Association for Cancer Research, Cosponsored by the American Society of Preventive Oncology*. 2017;26:85–94.

12. Kostic AD, Gevers D, Pedamallu CS, Michaud M, Duke F, Earl AM, et al. Genomic analysis identifies association of *Fusobacterium* with colorectal carcinoma. *Genome Research*. 2012;22:292–8.

13. Lu Y, Chen J, Zheng J, Hu G, Wang J, Huang C, et al. Mucosal adherent bacterial dysbiosis in patients with colorectal adenomas. *Scientific Reports*. 2016;6:26337.

14. Sanapareddy N, Legge RM, Jovov B, McCoy A, Burcal L, Araujo-Perez F, et al. Increased rectal microbial richness is associated with the presence of colorectal adenomas in humans. *The ISME journal*. 2012;6:1858–68.

15. Wang T, Cai G, Qiu Y, Fei N, Zhang M, Pang X, et al. Structural segregation of gut microbiota between colorectal cancer patients and healthy volunteers. *The ISME journal*. 2012;6:320–9.

16. Weir TL, Manter DK, Sheflin AM, Barnett BA, Heuberger AL, Ryan EP. Stool microbiome and metabolome differences between colorectal cancer patients and healthy adults. *PLoS*

124 One. 2013;8:e70803.

125 17. Zeller G, Tap J, Voigt AY, Sunagawa S, Kultima JR, Costea PI, et al. Potential of
126 fecal microbiota for early-stage detection of colorectal cancer. *Molecular Systems Biology*.
127 2014;10:766.

128 18. Zackular JP, Rogers MAM, Ruffin MT, Schloss PD. The human gut microbiome as
129 a screening tool for colorectal cancer. *Cancer Prevention Research (Philadelphia, Pa.)*.
130 2014;7:1112–21.

131 19. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, et al.
132 Introducing mothur: Open-Source, Platform-Independent, Community-Supported Software
133 for Describing and Comparing Microbial Communities. *Appl. Environ. Microbiol.* [Internet].
134 2009 [cited 12AD Jan 1];75:7537–41. Available from: [http://aem.asm.org/cgi/content/](http://aem.asm.org/cgi/content/abstract/75/23/7537)
135 [abstract/75/23/7537](http://aem.asm.org/cgi/content/abstract/75/23/7537)

136 20. Rognes T, Flouri T, Nichols B, Quince C, Mahé F. VSEARCH: A versatile open source
137 tool for metagenomics. *PeerJ*. 2016;4:e2584.

138 21. Westcott SL, Schloss PD. OptiClust, an Improved Method for Assigning
139 Amplicon-Based Sequence Data to Operational Taxonomic Units. *mSphere*. 2017;2.

142 **Figure 1:**

143 **Figure 2:**

144 **Figure 3:**

145 **Figure 4:**

146 **Figure S1:**

147 **Figure S2:**

148 **Figure S3:**