

The Microbiota and Individual Community Members in Colorectal Cancer: Is There a Common Theme?

Marc A Sze and Patrick D Schloss

Supplemental

Table S1: Bray-Curtis Distance Matrix Analyzed by PERMANOVA for Adenoma

F statistic	R2	P-value	Study	Adenoma n	Control n
0.63	0.059	0.7562	Brim	6	6
1.85	0.021	0.0096	Zeller	37	50
1.04	0.003	0.3788	Baxter	198	172
1.01	0.001	0.3658	Hale	214	473

Table S2: Bray-Curtis Distance Matrix Analyzed by PERMANOVA for Carcinoma

F statistic	R2	P-value	Study	Carcinoma n	Control n
3.47	0.034	0.0001	Wang	46	56
1.07	0.107	0.3431	Weir	7	4
2.18	0.010	0.0033	Ahn	62	148
2.38	0.019	0.0006	Zeller	41	87
2.15	0.004	0.0013	Baxter	120	370
0.97	0.001	0.7321	Hale	17	687
1.27	0.016	0.0460	Flemer	43	37

Table S3:

Table S2: Summary of Common OTUs for All Diagnosis Groups

Table S4: Summary of Adenoma based on Surgery Received

Table S5: Summary of mtry ranges used for the treatment and diagnosis models