

# BASiCS workflow: a step-by-step analysis of expression variability using single cell RNA sequencing data

Nils Eling<sup>\*1,2</sup>, Alan O'Callaghan<sup>3</sup>, John C. Marioni<sup>1,2</sup>, and Catalina A. Vallejos<sup>†3,4</sup>

<sup>1</sup>European Molecular Biology Laboratory, European Bioinformatics Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge CB10 1SD, UK

<sup>2</sup>Cancer Research UK Cambridge Institute, University of Cambridge, Li Ka Shing Centre, Cambridge, CB2 0RE, UK

<sup>3</sup>MRC Human Genetics Unit, Institute of Genetics & Molecular Medicine, University of Edinburgh, Western General Hospital, Crewe Road, Edinburgh, EH4 2XU, UK

<sup>4</sup>The Alan Turing Institute, British Library, 96 Euston Road, London, NW1 2DB, UK

**Abstract** Cell-to-cell gene expression variability is an inherent feature of complex biological systems. Single-cell RNA sequencing can be used to quantify this heterogeneity, but it is prone to strong technical noise. Here, we describe a step-by-step computational workflow which uses the BASiCS Bioconductor package to robustly quantify expression variability within and between known cell populations (such as experimental conditions or cell types). BASiCS provides an integrated framework for data normalisation, technical noise quantification and downstream analyses, whilst propagating statistical uncertainty across these steps. Within a single seemingly homogeneous cell population, BASiCS can be used to identify highly variable genes that drive the heterogeneity within the population as well as lowly variable genes that might exhibit housekeeping-like behavior. BASiCS also provides a probabilistic rule to identify changes in expression variability between cell populations, while avoiding confounding effects related to differences in technical noise or in overall abundance. Using two publicly available datasets, we guide users through a complete pipeline which includes preliminary steps for quality control and data exploration using the *scater* and *scran* Bioconductor packages. Data for the first case study was generated using the Fluidigm® C1 system, in which extrinsic spike-in RNA molecules were added in order to quantify technical noise. The second dataset was generated using a droplet-based system, for which spike-in RNA is not available. The latter analysis provides an example, in which differential variability testing reveals insights regarding a possible early cell fate commitment process.

## Keywords

Single-cell RNA sequencing, expression variability, transcriptional noise, differential expression testing

---

\*eling@ebi.ac.uk

†catalina.vallejos@igmm.ed.ac.uk

## Introduction

Single-cell RNA-sequencing (scRNA-seq) enables the study of genome-wide transcriptional heterogeneity in cell populations that remains otherwise undetected in bulk experiments [1, 2, 3]. Existing applications range from characterising cell types in immunity [4, 5, 6] and development [7, 8, 9] to dissecting the mechanisms for cell fate commitment [10, 11]. Transcriptional heterogeneity within a population of cells can relate to different reasons. On the broadest level, this heterogeneity can reflect the presence of distinct expression profiles associated to cell subtypes or states which could be characterised through clustering [12]. More subtle gene expression variability within a seemingly homogeneous cell population can be due to deterministic or stochastic events, the later being referred to as ‘noise’ [13].

## References

- [1] Oliver Stegle, Sarah a. Teichmann, and John C. Marioni. Computational and analytical challenges in single-cell transcriptomics. *Nature Reviews Genetics*, 16(3):133–145, jan 2015. ISSN 1471-0056. doi: 10.1038/nrg3833. URL <http://www.nature.com/doi/10.1038/nrg3833>.
- [2] Sanjay M. Prakadan, Alex K. Shalek, and David A. Weitz. Scaling by shrinking: empowering single-cell ‘omics’ with microfluidic devices. *Nature Reviews Genetics*, 18(6):345–361, 2017. ISSN 1471-0056. doi: 10.1038/nrg.2017.15. URL <http://www.nature.com/doi/10.1038/nrg.2017.15>.
- [3] Simona Patange, Michelle Girvan, and Daniel R. Larson. Single-cell systems biology: Probing the basic unit of information flow. *Current Opinion in Systems Biology*, 8:7–15, 2018. ISSN 24523100. doi: 10.1016/j.coisb.2017.11.011. URL <https://doi.org/10.1016/j.coisb.2017.11.011>.
- [4] Tapio Lönnberg, Valentine Svensson, Kylie R. James, Daniel Fernandez-Ruiz, Ismail Sebina, Ruddy Montandon, Megan S. F. Soon, Lily G. Fogg, Arya Sheela Nair, Urijah N. Liligeto, Michael J. T. Stubbington, Lam-Ha Ly, Fredrik Otzen Bagger, Max Zwiessele, Neil D. Lawrence, Fernando Souza-Fonseca-Guimaraes, Patrick T. Bunn, Christian R. Engwerda, William R. Heath, Oliver Billker, Oliver Stegle, Ashraf Haque, and Sarah A. Teichmann. Single-cell RNA-seq and computational analysis using temporal mixture modeling resolves Th1/Tfh fate bifurcation in malaria. *Science Immunology*, 2(9):eaal2192, 2017. ISSN 2470-9468. doi: 10.1126/sciimmunol.aal2192. URL <http://immunology.sciencemag.org/lookup/doi/10.1126/sciimmunol.aal2192>.
- [5] Alexandra Chloé Villani, Rahul Satija, Gary Reynolds, Siranush Sarkizova, Karthik Shekhar, James Fletcher, Morgane Griesbeck, Andrew Butler, Shiwei Zheng, Suzan Lazo, Laura Jardine, David Dixon, Emily Stephenson, Emil Nilsson, Ida Grundberg, David McDonald, Andrew Filby, Weibo Li, Philip L. De Jager, Orit Rozenblatt-Rosen, Andrew A. Lane, Muzlifah Haniffa, Aviv Regev, and Nir Hacohen. Single-cell RNA-seq reveals new types of human blood dendritic cells, monocytes, and progenitors. *Science*, 356(6335), 2017. ISSN 10959203. doi: 10.1126/science.aah4573.
- [6] Grace X.Y. Zheng, Jessica M. Terry, Phillip Belgrader, Paul Ryvkin, Zachary W. Bent, Ryan Wilson, Solongo B. Ziraldo, Tobias D. Wheeler, Geoff P. McDermott, Junjie Zhu, Mark T. Gregory, Joe Shuga, Luz Montesclaros, Jason G. Underwood, Donald A. Masquelier, Stefanie Y. Nishimura, Michael Schnall-Levin, Paul W. Wyatt, Christopher M. Hindson, Rajiv Bharadwaj, Alexander Wong, Kevin D. Ness, Lan W. Beppu, H. Joachim Deeg, Christopher McFarland, Keith R. Loeb, William J. Valente, Nolan G. Ericson, Emily A. Stevens, Jerald P. Radich, Tarjei S. Mikkelsen, Benjamin J. Hindson, and Jason H. Bielas. Massively parallel digital transcriptional profiling of single cells. *Nature Communications*, 8: 1–12, 2017. ISSN 20411723. doi: 10.1038/ncomms14049. URL <http://dx.doi.org/10.1038/ncomms14049>.
- [7] Ximena Ibarra-Soria, Wajid Jawaid, Blanca Pijuan-Sala, Vasileios Ladopoulos, Antonio Scialdone, David J. Jörg, Richard C.V. Tyser, Fernando J. Calero-Nieto, Carla Mulas, Jennifer Nichols, Ludovic Vallier, Shankar Srinivas, Benjamin D. Simons, Berthold Göttgens, and John C. Marioni. Defining murine organogenesis at single-cell resolution reveals a role for the leukotriene pathway in regulating blood progenitor formation. *Nature Cell Biology*, 20(2):127–134, 2018. ISSN 14764679. doi: 10.1038/s41566-017-0013-z.
- [8] Daniel E. Wagner, Caleb Weinreb, Zach M. Collins, James A. Briggs, Sean G. Megason, and Allon M. Klein. Single-cell mapping of gene expression landscapes and lineage in the zebrafish embryo. *Science*, 4362:1–12, 2018. ISSN 10959203. doi: 10.1126/science.aar4362.
- [9] Blanca Pijuan-Sala, Jonathan A. Griffiths, Carolina Guibentif, Tom W. Hiscock, Wajid Jawaid, Fernando J. Calero-Nieto, Carla Mulas, Ximena Ibarra-Soria, Richard C.V. Tyser, Debbie Lee Lian Ho, Wolf Reik, Shankar Srinivas, Benjamin D. Simons, Jennifer Nichols, John C. Marioni, and Berthold Göttgens. A single-cell molecular map of mouse gastrulation and early organogenesis. *Nature*, 566(7745):490–495, 2019. ISSN 14764687. doi: 10.1038/s41586-019-0933-9.
- [10] Mubeen Goolam, Antonio Scialdone, Sarah J. L. Graham, Iain C. MacAulay, Agnieszka Jedrusik, Anna Hupalowska, Thierry Voet, John C. Marioni, and Magdalena Zernicka-Goetz. Heterogeneity in Oct4 and Sox2 Targets Biases Cell Fate in 4-Cell Mouse Embryos. *Cell*, 165(1):61–74, 2016. ISSN 10974172. doi: 10.1016/j.cell.2016.01.047. URL <http://dx.doi.org/10.1016/j.cell.2016.01.047>.
- [11] Yusuke Ohnishi, Wolfgang Huber, Akiko Tsumura, Minjung Kang, Panagiotis Xenopoulos, Kazuki Kurimoto, Andrzej K. Oleś, Marcos J. Araújo-Bravo, Mitunori Saitou, Anna-Katerina Hadjantonakis, and Takashi Hiiragi. Cell-to-cell expression variability followed by signal reinforcement progressively segregates early mouse lineages. *Nature Cell Biology*, 16(1):27–37, 2014. ISSN 1476-4679. doi: 10.1038/ncb2881. URL <http://dx.doi.org/10.1038/ncb2881>.
- [12] Vladimir Yu Kiselev, Tallulah S. Andrews, and Martin Hemberg. Challenges in unsupervised clustering of single-cell RNA-seq data. *Nature Reviews Genetics* 2018, 2019. ISSN 1471-0064. doi: 10.1038/s41576-018-0088-9. URL [https://www.nature.com/articles/s41576-018-0088-9?utm\\_source=feedburner&utm\\_medium=feed&utm\\_campaign=Feed%2F3A+nrg%2Frs%2Fcurrent%28Nature+Reviews+Genetics%29](https://www.nature.com/articles/s41576-018-0088-9?utm_source=feedburner&utm_medium=feed&utm_campaign=Feed%2F3A+nrg%2Frs%2Fcurrent%28Nature+Reviews+Genetics%29).

- [13] Michael B Elowitz, Arnold J Levine, Eric D Siggia, and Peter S Swain. Stochastic gene expression in a single cell. *Science*, 297(5584):1183–1186, 2002. ISSN 1095-9203. doi: 10.1126/science.1070919. URL <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve{&}db=PubMed{&}dopt=Citation{&}list{&}uids=12183631>.