

JMP Exam Notes

ONE MEAN · POWER · TWO PROPORTIONS

➤ = then click · ▼ = the little red triangle next to that report's title · → = where the answer shows up

SYMBOL KEY — WHAT EACH LETTER MEANS

GREEK = THE TRUE VALUE YOU CAN'T SEE — μ "mew" — the true mean of the whole population · σ "sigma" — the true standard deviation, only when a problem gives it · α "alpha" — the significance level, the cutoff you compare p to (.05, .01) · β "beta" — the chance of missing a real difference · β_0 , β_1 "beta-zero, beta-one" — the true intercept and slope · Σ big sigma — "add up over every data point" · χ^2 "chi-square" — the test JMP prints for a 2x2 table.

LATIN = WHAT YOUR SAMPLE GIVES YOU — $\bar{x}$ "x-bar" — your sample average · s your sample standard deviation · n how many observations · $\hat{p}$ "p-hat" — a sample proportion · b_0 , b_1 the intercept and slope JMP prints · $\hat{y}$ "y-hat" — a predicted y · e a residual: actual – predicted · **SE** standard error = $\sigma/\sqrt{n}$ or $s/\sqrt{n}$ · r correlation, –1 to +1 · **R²** share of y explained · **df** degrees of freedom · **k** how many predictors.

A hypothesis is always about the Greek letter, never $\bar{x}$. Careful: **B** — alone is the Type II error chance, but **B₀**, **B₁** — are regression slopes.

STUDENT CALCULATORS & TEACHING MODULES

GETTING THERE, THEN WHICH FILE — **Help** > **Sample Index** → under ① Content Type click **Tool** → under ② Tool Type pick **Calculators** → double-click the file. (Some builds list these under the **Student** menu.) One mean, test statistic + p-value → **TestMean.jsl** · one proportion → **TestProportion.jsl** · two means → **TestTwoSampleMeans.jsl** · two proportions → **TestTwoSampleProportions.jsl** · CI for a mean → **ConfidenceIntervalsForTheMean.jsl** · CI for a proportion → **ConfidenceIntervalsForProportion.jsl** · CI comparing two groups → **CITwoMeans.jsl**, **CITwoProportions.jsl** · any normal / t / binomial probability → **DistributionCalculator.jsl** · how big a sample → **SampleSizeForConfidenceIntervals.jsl**.

TEACHING MODULES — SHAPE, MEAN, SD, AND A GUIDED TEST — Same path but Tool Type **Teaching Modules**. **Sampling Distribution of Sample Means**: set *Population Shape* and *Population Mean* → *Population Summary* gives the shape, Mean and Std Dev, and the sample-means panel gives the mean of $\bar{x}$ and its Std Dev (that's $\sigma/\sqrt{n}$). **Hypothesis Test on Mean**: pick *z-test* or *t-test*, pick the alternative (not equal / less / greater), then tick *Reveal Decision*.

TESTING ONE MEAN

Is it a number or a group? If you could take an average of it, it's a number → use a **mean**. If it just sorts each person into a group (yes/no, accepted/rejected), it's a group → use a **proportion**.

Writing the hypotheses. H_0 always says μ equals the claimed number. "less / went down" → $H_1: \mu <$ · "more / went up" → $>$ · "different / changed" → $\neq$.

Shape, average, spread, count. Analyze > Distribution > Y, Columns = variable > OK → the histogram shows the shape; **Summary Statistics** gives Mean, Std Dev, N and Std Err Mean.

t test — when they don't give you σ . Analyze > Distribution > Y = variable > OK > ▼variable > Test Mean > type the claimed number > leave the standard deviation box empty > OK → the **Test Mean** panel, **t Test** row.

z test — when they do give you σ . Same path, but also fill in "Enter True Standard Deviation to do z-test rather than t test" → the panel now shows a **z Test** row instead.

Which p-value to read. H_1 has $\neq$ → **Prob>|t|** · H_1 has $>$ → **Prob>t** · H_1 has $<$ → **Prob<t**. Ignore the Signed-Rank line.

Only summary numbers given. Distribution needs raw data, so use **TestMean.jsl** above, or $t = (\bar{x} - \mu_0) / (s/\sqrt{n})$, $df = n - 1$. If they give s^2 , square-root it first.

WHAT HAS TO BE TRUE, AND INTERVALS

Conditions for the t test. Independence: it's a random sample, it's less than 10% of the whole group, and each person is measured once. **Bell shape:** the problem says bell-shaped, or $n \geq 30$ (big samples are fine either way), or the histogram looks even with no far-out points.

Checking the shape in JMP. ▼variable > Continuous Fit > Normal > ▼Fitted Normal > Goodness of Fit → **Shapiro-Wilk Prob<W**; if it's bigger than α , the bell shape is fine. Also ▼variable > Normal Quantile Plot — the dots should hug the straight line.

Confidence interval for the mean. **Summary Statistics** already lists Upper / Lower 95% Mean; other levels ▼variable > Confidence Interval > 0.99. Formula $\bar{x} \pm t(\alpha/2, n-1) \cdot s/\sqrt{n}$.

IF H_0 WERE TRUE, AND POWER

Describe the spread of sample averages when H_0 is true. Give all three. **Shape:** bell-shaped, because the population is bell-shaped or $n \geq 30$. **Center:** μ_0 . **Spread:** $\sigma/\sqrt{n}$.

Which sample averages make you reject? cutoff = $\mu_0 + (z)(SE)$, where $SE = \sigma/\sqrt{n}$ and z carries the sign of the tail: –1.645 for a "less than" test, +1.645 for a "greater than" test at $\alpha = .05$; for a two-sided test use $\mu_0 \pm 1.96 \cdot SE$. Reject when $\bar{x}$ is past that cutoff.

β — the chance of missing a real difference. Take that same cutoff, but measure it from the **true** mean instead: $z = (\text{cutoff} - \mu_{\text{true}}) / (\sigma/\sqrt{n})$. Then β is the chance $\bar{x}$ lands in the "don't reject" zone.

Power = 1 – β . These all mean power: "chance of detecting this difference", "chance of rejecting H_0 when the true mean is _____", "chance of correctly concluding". These mean β : "chance of failing to reject", "chance of a Type II error".

Power from JMP. DOE > Design Diagnostics > Sample Size and Power > One Sample Mean, fill in Alpha, Std Dev, Difference to Detect and Sample Size, leave **Power** empty > Continue. Leaving **Sample Size** empty instead tells you how big a sample you'd need. It assumes a two-sided test, so on one-sided questions use the formula above.

Drawing it. Two bell curves on one line, one vertical line at the cutoff through both: top curve at μ_0 , tail past the cutoff shaded = α ; bottom curve at the true mean, "don't reject" side shaded = β , the rest power.

WORKED EXAMPLE —

POWER Soap: $\mu_0 = 33$, $\sigma = 12.4$, $n = 35$, $\alpha = .05$ "less than", true mean 29
SE = $\sigma/\sqrt{n} = 12.4/\sqrt{35} = 12.4/5.9161 \approx 2.096$
 $\bar{x}c = \mu_0 + (z)(SE) = 33 + (-1.645)(2.096) = 33 - 3.448 \approx 29.55$
 $z = (\bar{x}c - \mu_{\text{true}})/SE = (29.55 - 29)/2.096 = 0.55/2.096 \approx 0.263$
Power = $P(Z < 0.263) \approx .6039 \rightarrow 60.39\%$ $\beta = 1 - .6039 \approx .3961$
t TEST Life Expectancy: $SE = 16.41/\sqrt{200} \approx 1.1604$, $t = 0.98/1.1604 \approx 0.8446$

TWO PROPORTIONS — SEPARATE GROUPS

A table of counts. File > New > Data Table with Group (Nominal), Outcome (Nominal), Count (Continuous), one row per box, like (Help, Accepted, 222). Then Analyze > Fit Y by X > Y = Outcome > X = Group > Freq = Count > OK → **Contingency Table** and **Tests**. Skip **Freq** and JMP thinks you have 4 people.

Best guess for the difference. ▼Contingency Table > Row %. Each $\hat{p}$ is the count divided by that group's own row total, not the overall total. Example: $222/515 = 55/190 = .4311 - .2895 = .1416$.

Hypotheses. $H_0: p_1 - p_2 = 0$ — if both groups have the same rate, the difference is zero. "improves" → > 0 · "reduces" → < 0 · "different" → $\neq 0$.

What has to be true. Two separate random groups, or people randomly assigned to them; one answer per person; and all four counts in the table at least 5.

Test and interval. ▼Contingency Analysis > Two Sample Test for Proportions → the difference, its standard error, **Prob>|z|** / **Prob>z** / **Prob<z**, and the confidence interval. **Pearson ChiSquare** in the Tests panel is the same two-sided test.

Formulas. combined $\hat{p} = (x_1+x_2)/(n_1+n_2)$, then $z = (\hat{p}_1-\hat{p}_2) / \sqrt{\hat{p}(1-\hat{p})(1/n_1+1/n_2)}$. The interval does **not** combine them: $(\hat{p}_1-\hat{p}_2) \pm z\alpha/2\sqrt{\hat{p}_1\hat{q}_1/n_1 + \hat{p}_2\hat{q}_2/n_2}$.

TWO PROPORTIONS — SAME PEOPLE TWICE

Spotting it and reading the table. Same people measured twice, or matched pairs. **a** and **d** are the people who didn't change; **b** and **c** are the ones who switched, one each way. Smoking: $a = 20$, $b = 5$, $c = 16$, $d = 9$, $n = 50$. Don't use the test above — it ignores the pairing.

Only the switchers count. $n' = b + c$. Everyone who stayed the same ($a + d$) gets dropped. T = how many switched in the direction H_1 expects. Smoking: 29 dropped, $n' = 21$, $T = 16$. H_0 says switching is equally likely both ways.

The p-value. If H_0 is true, T acts like flipping a coin n' times: $T \sim \text{Binomial}(n', .05)$. Cols > New Column > Formula > Discrete Probability > Binomial Distribution(.5, n' , k) gives the chance of k or fewer. Smoking: $1 - \text{BinDist}(.5, 21, 15) = .0133$, under .05, so the video worked.

Or let JMP do it. Analyze > Fit Y by X > Y = After, X = Before, Freq = Count > OK > ▼ > Agreement Statistic → the **test of symmetry** is McNemar's test, but it only gives the two-sided answer.

Interval. $(b-c)/n \pm z\alpha/2\sqrt{[(a+d)(b+c) + 4bc] / n^3}$. Smoking: (–.389, –.051), all below zero. Puzzle: (–.196, .077), which includes zero — no real difference.

FILL-IN-THE-BLANK ANSWERS

DEFINE THE PARAMETER — μ = the true mean _____ (with units) for all _____.

INTERPRET THE TEST STATISTIC — The sample average is _____ standard errors [above / below] _____, the number H_0 claims.

INTERPRET THE P-VALUE — If the true mean really were _____, there is a _____ % chance of getting a sample average at least this far from _____ just by chance.

CONCLUSION — WORKS FOR EVERY TEST — Since $p =$ _____ [$<$ / $>$] $\alpha =$ _____, we [reject / fail to reject] H_0 . There [is / is not] enough evidence to say that _____.

Last blank → **ONE MEAN:** — the mean [variable] is [less than / greater than / different from] ____ · **TWO PROPORTIONS:** — the proportion of ____ is [higher / lower / different] for ____ than for ____ · **SLOPE:** — [x] helps predict [y] · **OVERALL F:** — the model helps predict [y] · **CURVED MODEL:** — the squared term makes the model better. Never write "accept H_0 ." If your instructor words it "sufficient evidence to conclude," that is the same sentence.

INTERPRET ANY CONFIDENCE INTERVAL — We are _____ % confident that _____ is between _____ and _____.

Middle blank → **M:** — the true mean [variable] for all [group] · **P₁–P₂:** — the difference in the proportion of ____ between ____ and ____ · **SLOPE:** — the average change in [y] for each extra one [x-unit] · **MEAN AT X:** — the average [y] for all [units] with $x =$ ____.

PREDICTION: — a single [unit] with $x =$ ____ will have [y]. If the interval includes 0 → no real difference / doesn't help predict.

TYPE I ERROR IN CONTEXT — We decide that _____ when really _____ . The chance of this is $\alpha =$ _____.

TYPE II ERROR IN CONTEXT — We decide there isn't enough evidence that _____ when really _____ is true. The chance of this is $\beta =$ _____.

DESCRIBE THE SPREAD OF SAMPLE AVERAGES — **Shape:** bell-shaped, because _____ . **Center:** $\mu =$ _____ . **Spread:** $\sigma/\sqrt{n} =$ _____ / $\sqrt{\quad} =$ _____.

Reason blank: "the problem says the population is bell-shaped" or " $n \geq 30$, so it's fine either way".

CHECK THE T-TEST CONDITIONS — **Independence:** the _____ are a random sample and fewer than 10% of all _____ . **Bell shape:** _____.

POWER AND B — If the true mean is really _____, there is a _____ % chance we [correctly reject / fail to reject] H_0 . "Correctly reject" = power = $1 - \beta$. "Fail to reject" = β = the chance of a Type II error.

WHICH ERROR WAS POSSIBLE — We [rejected / failed to reject] H_0 , so if we made a mistake it has to be a Type [I / II] error: _____.

LARGE ENOUGH SAMPLE, TWO PROPORTIONS — The counts _____, _____, _____ and _____ are all at least 5, so both groups are big enough.

JMP Exam Notes

CORRELATION AND THE BEST-FIT LINE

Which variable goes where. "Predict A using B" → A is the response and goes in **Y, Response**; B does the predicting and goes in **X, Factor**.

Scatterplot. Analyze > Fit Y by X > Y, Response = response > X, Factor = factor > OK.

Add the line. **▼**Bivariate Fit > Fit Line → equation printed right above **Summary of Fit**. For the curved bands around it: **▼**Linear Fit > Confid Curves Fit or Confid Curves Indiv (picture only).

Correlation r. Analyze > Multivariate Methods > Multivariate > both variables in Y, Columns > OK → **Correlations**, the number where the two variables cross (the 1.0000s are each variable with itself). Or $r = \pm \sqrt{R^2}$, using the sign of the slope. r always lands between -1 and +1: the sign is the direction, the size is the strength — $|r| \geq .8$ strong, .5-.8 medium, under .5 weak.

Equation of the best-fit line. Analyze > Fit Model > Y = response > select x > Add > Run → **Parameter Estimates, Estimate** column: the Intercept row is b_0 , the x row is b_1 .

Predict. Put the x value into $\hat{y} = b_0 + b_1x$. Watch the units — if x is in thousands, 34,000 means $x = 34$. Don't predict past the range of the data you have.

CHECKING IT'S OK TO USE THE LINE

Save the leftovers. **▼**Response y > Save Columns > Residuals, then again for **Predicted Values** and **Studentized Residuals** → new columns at the right of the data table. A residual is just actual – predicted.

Are the leftovers bell-shaped? Analyze > Distribution > Y = Residual > OK > **▼**Residual > Normal Quantile Plot, then Continuous Fit > Normal > **▼**Fitted Normal > Goodness of Fit → **Shapiro-Wilk**.

Leftovers plot. **▼**Response y > Row Diagnostics > Plot Residual by Predicted. A shapeless band around zero is what you want · an arch or U means a line is the wrong shape · a funnel means the spread isn't even · a studentized value past ± 3 is a far-out point.

DOES X ACTUALLY HELP PREDICT Y?

Test the slope. Fit Model > Run > Parameter Estimates, on the x row → **t Ratio** is the test number, **Prob>|t|** is the two-sided p-value, **Std Error** is the slope's wobble. H_0 : the slope is 0, $df = n - 2$.

One-sided? Cut **Prob>|t|** in half, but only if the slope's sign matches what H_1 says. If it doesn't, the p-value is over .5 and you fail to reject.

Testing the slope against a number other than 0. JMP only tests 0. Either check whether the confidence interval contains that number, or use $t = (b_1 - c) / \text{Std Error}$, $df = n - 2$.

How much of y is explained. **Summary of Fit** → **RSquare**, times 100 for a percent.

THE THREE INTERVALS

Interval for the slope. Right-click inside **Parameter Estimates** > Columns > Lower 95% + Upper 95% → read them on the **x** row, not the Intercept row. If it doesn't include 0, x helps predict y .

99% or 98% instead. **▼**Response y > Set Alpha Level > 0.01 (or **Other** → 0.02), then read those columns again. If yours is stuck at 95%: $b_1 \pm t(\alpha/2, n-2) \cdot \text{Std Error}$, with t from $t_{\text{Quantile}(1-\alpha/2, n-2)}$.

Interval for the average y — "for ALL cases with $x = \text{___}$ ". Add a row with that x and **leave Y empty** (that keeps it out of the line), then Fit Model > Run > **▼**Response y > Save Columns > Mean Confidence Interval → **Lower / Upper 95% Mean** on that row.

Interval for ONE new case. Same setup, then Save Columns > Indiv Confidence Interval → **Lower / Upper 95% Indiv**, always wider. A singular word — "a truck", "a school" — means use Indiv.

Far-out points. Click the point to light up its row, then Rows > Exclude/Unexclude and Run again. "How many more votes than predicted" = actual – predicted, which is that row's residual.

SEVERAL PREDICTORS AT ONCE

Look at them first. Analyze > Multivariate Methods > Multivariate > response + all predictors in Y, Columns > OK → **Correlations** and a grid of mini scatterplots. Two predictors with $|r|$ above about .8 measure nearly the same thing, which causes trouble.

Fit it. Analyze > Fit Model > Y = response > highlight all the predictors > Add > Run → **Parameter Estimates** gives $\hat{y} = b_0 + b_1x_1 + \dots + b_kx_k$.

Does the whole model help? **Analysis of Variance** → **F Ratio** and **Prob > F**. H_0 says every slope is 0. Rejecting means at least one predictor helps — not that they all do.

Does one predictor help, with the others in? **Parameter Estimates**, that predictor's row → **t Ratio, Prob>|t|**, $df = n - k - 1$. A predictor can look strong alone and still be useless here, because another one already covers it.

Interval for one coefficient. Right-click **Parameter Estimates** > Columns > Lower 95% + Upper 95%; change the level first with **Set Alpha Level**.

Checking conditions. Save **Residuals**, use **▼**Response y > Row Diagnostics > Plot Residual by Predicted, and run Distribution on the residuals for the plot and Shapiro-Wilk. For overlap between predictors: right-click **Parameter Estimates** > Columns > VIF — over 5 a warning, over 10 a problem.

PICKING THE BEST MODEL

Use adjusted R^2 . **Summary of Fit** → **RSquare Adj**, highest one wins. Plain R^2 always goes up when you add a predictor, so it can't compare models with different numbers of predictors; $R^2_{\text{adj}} = 1 - (1 - R^2)(n-1)/(n-k-1)$ takes a penalty for extras.

Every model at once. Fit Model > add all > Personality = Stepwise > Run > **▼** > All Possible Models → every combination with R^2 , R^2_{adj} , Cp, AICc, BIC.

Dropping predictors one at a time. Fit Model > add all > Stepwise > Run, set Direction = **Backward**, Stopping Rule = **P-value Threshold**, Prob to Leave = **0.1**, click **Go** → **Current Estimates**, a check under **Entered** means it stayed. Then **Make Model** → **Run**. The idea: throw out the biggest p-value above 0.1, refit, repeat.

Predicting from the model you picked. Put the values into that model's equation, or add a row with those x 's and an empty Y , save **Predicted Values**, or use **▼**Response y > Factor Profiling > Profiler.

CORRELATION · ONE PREDICTOR · SEVERAL PREDICTORS · CURVED

CURVED (QUADRATIC) MODELS

Quick way. Analyze > Fit Y by X > Y, X > OK > **▼**Bivariate Fit > Fit Polynomial > 2, quadratic → a **Polynomial Fit Degree=2** block with the plain equation printed above it.

Full way (use this one for intervals). Analyze > Fit Model > Y > select x > Macros > Polynomial to Degree with Degree = 2 > Run → **Parameter Estimates** gains an **x^2** row.

JMP shifts x first. By default it prints $(x-\bar{x}) \cdot (x-\bar{x})$ instead of x^2 , so the numbers aren't the plain b_0, b_1, b_2 . Uncheck **▼**Response y > Center Polynomials and Run again to get the plain equation.

Does the curve help? Test the squared term only: **Parameter Estimates**, the **x^2** row (or **Effect Tests**) → **Prob>|t|**. H_0 says the squared term is 0, meaning a straight line is enough. Don't use the overall F here — it can look good just from the straight-line part.

The change in y isn't the same every time. change = $b_1(x_2 - x_1) + b_2(x_2^2 - x_1^2)$. Going 0 to 1 gives $b_1 + b_2$; going 2 to 3 gives $b_1 + 5b_2$. They're different on purpose — that's the whole point of asking both.

Intervals from a curved model. Add the row with x and an empty Y , fit with **Polynomial to Degree 2**, then **Save Columns** → **Mean Confidence Interval** for the average of all such cases, **Indiv Confidence Interval** for one specific new case.

NUMBERS YOU MAY NEED TO LOOK UP

df: $n - 1$ for one mean · $n - 2$ for a slope · $n - k - 1$ for one predictor out of k · k and $n - k - 1$ for the overall F .

z cutoffs: one-sided $\alpha = .10 \rightarrow 1.282, .05 \rightarrow 1.645, .025 \rightarrow 1.960, .01 \rightarrow 2.326$ · two-sided 90% → 1.645, 95% → 1.960, 98% → 2.326, 99% → 2.576.

THE FORMULAS BEHIND THE LINE

Three building blocks. JMP never shows these three — they're only for working a small data set out yourself. $S_{xx} = \sum (x - \bar{x})^2$ spread of the x 's, feeds the slope and its standard error · $S_{yy} = \sum (y - \bar{y})^2$ spread of the y 's, feeds r and R^2 · $S_{xy} = \sum (x - \bar{x})(y - \bar{y})$ how much x and y move together, feeds the slope and r .

Everything else comes from those three. slope $b_1 = S_{xy}/S_{xx}$ · intercept $b_0 = \bar{y} - b_1\bar{x}$ (needs b_1 first) · the line $\hat{y} = b_0 + b_1x$ · $r = S_{xy}/\sqrt{(S_{xx} \cdot S_{yy})}$ · $R^2 = r^2$ · one point's leftover $e = y - \hat{y}$. To check your work, your b_0 and b_1 should match **Parameter Estimates**.

FILL-IN-THE-BLANK ANSWERS — REGRESSION

COMMENT ON THE SCATTERPLOT —There is a [positive / negative], [weak / medium / strong], [straight / curved] relationship between _____ and _____.
Add if needed: "There is an odd point at ____." If it curves, say a squared term should be tried.

INTERPRET R — $r = \text{_____}$ means a [weak / medium / strong] [positive / negative] straight-line relationship between _____ and _____.
WRITE THE EQUATION —Predicted _____ = _____ + _____ (_____).

INTERPRET THE SLOPE —For every extra one _____, the predicted (average) _____ [goes up / goes down] by _____.
INTERPRET THE INTERCEPT —When _____ is 0, the predicted _____ is _____.
If $x = 0$ is outside the data: add "but $x = 0$ isn't in the range of the data, so this doesn't really mean anything."

INTERPRET R^2 — _____ % of the changes in _____ are explained by the [straight / curved] relationship with _____.
DOES X HELP PREDICT Y? —Since $p = \text{_____}$ [$< / >$] $\alpha = \text{_____}$, we [reject / fail to reject] H_0 : the slope is 0. There [is / is not] enough evidence to say that _____ helps predict _____.
INTERVAL FOR THE SLOPE —We are _____ % confident that for every extra one _____, the average _____ changes by between _____ and _____.
INTERVAL FOR THE AVERAGE Y —We are _____ % confident that the average _____ for all _____ with _____ = _____ is between _____ and _____.
INTERVAL FOR ONE NEW CASE —We are _____ % confident that a single _____ with _____ = _____ will have a _____ between _____ and _____.
WHAT THE LEFTOVERS PLOT SHOWS —The plot shows _____, so the [straight-line / even spread] condition [is / is not] met. Shapiro-Wilk gives $p = \text{_____}$ [$> / <$] α , so the bell shape [is / is not] reasonable.
First blank: "a shapeless band around zero" · "an arched pattern" · "a funnel shape".

INTERPRET A COEFFICIENT (SEVERAL PREDICTORS) —For every extra one _____, the predicted _____ [goes up / goes down] by _____, while the other predictors stay the same.

DOES THE WHOLE MODEL HELP? —Since $p = \text{_____}$ [$< / >$] $\alpha = \text{_____}$, we [reject / fail to reject] H_0 : every slope is 0. There [is / is not] enough evidence to say that the model helps predict _____.
ONE PREDICTOR, OTHERS ALREADY IN —With the other predictors already in the model, there [is / is not] enough evidence to say that _____ helps predict _____ ($t = \text{_____}$, $p = \text{_____}$).

PICKING A MODEL —The model with _____ has the highest adjusted R^2 (_____), so it's the one to use.

DOES THE CURVE HELP? —The squared term has $p = \text{_____}$ [$< / >$] $\alpha = \text{_____}$, so there [is / is not] enough evidence to say that the curved model fits better than a straight line.