List of Figures

1 Phase portraits of H_t on $[-7.5, 7.5] \times [-5, 5]$ at (a) $t = 0$, (b) $t = 2/3$, (c) $t = 4/3$, and (d) $t = 2$: the color shows $\arg H_t$ (key at right), the thin lines are the level sets $\arg H_t \in \frac{\pi}{6}\mathbb{Z}$, and the shading steps in $|H_t|$ by factors of $e^{2\pi/12} \approx 1.69$; the white line is $|H_t| = 1$. 2

2 The deformation H_t from the identity ($t = 0$) to F ($t = 2$), at (a) $t = 0$, (b) $t = 2/3$, (c) $t = 4/3$, and (d) $t = 2$. 3

3 The deformation H_t on the Riemann sphere, with ∞ at the north pole: the image of the set $|\operatorname{Im} z| < \pi/2$, the line $\operatorname{Im} z = 0.8$, and the point $z_0 = 0$ under F ($t = 2$) over its boundary at $t = 0$ (gray). 4

4 Phase portrait of H_t on $[-7, 7] \times [-5, 5]$: the color shows $\arg H_t$ (key at right), the thin lines are the level sets $\arg H_t \in \frac{\pi}{6}\mathbb{Z}$, and the shading steps in $|H_t|$ by factors of $e^{2\pi/12} \approx 1.69$; the white line is $|H_t| = 1$. 5

1 Body

Text. See Figure 4.




Figure 1: Phase portraits of H_t on $[-7.5, 7.5] \times [-5, 5]$ at (a) $t = 0$, (b) $t = 2/3$, (c) $t = 4/3$, and (d) $t = 2$: the color shows $\arg H_t$ (key at right), the thin lines are the level sets $\arg H_t \in \frac{\pi}{6}\mathbb{Z}$, and the shading steps in $|H_t|$ by factors of $e^{2\pi/12} \approx 1.69$; the white line is $|H_t| = 1$. Zeros are dots and poles are open circles. Here H_t deforms the identity ($t = 0$) into F ($t = 2$) along the homotopy H_t, $0 \le t \le 2$, that runs the steps in turn ($H_t = H^{(k)}_{t-k+1}$ on $k-1 \le t \le k$): for f_1, the zoom homotopy $H^{(1)}_s(z) = (e^{sz} - 1)/s + s$ for $0 < s \le 1$, $H^{(1)}_0(z) = z$; for f_2, the elliptic Möbius transformations $H^{(2)}_s$ that fix $p = i$ and $q = -i$, with $H^{(2)\prime}_s(p) = e^{-\pi i s/2}$. The map is

$$F(z) = \frac{e^z - 1}{e^z + 1}.$$




Figure 2: The deformation H_t from the identity ($t = 0$) to F ($t = 2$), at (a) $t = 0$, (b) $t = 2/3$, (c) $t = 4/3$, and (d) $t = 2$, where

$$F(z) = \frac{e^z - 1}{e^z + 1}.$$

Solid: the paths $t \mapsto H_t(z)$ of the marked points up to the frame's t; dotted: the rest of each path; dashed: the boundary at $t = 0$.



Figure 3: The deformation H_t on the Riemann sphere, with ∞ at the north pole: the image of the set $|\operatorname{Im} z| < \pi/2$, the line $\operatorname{Im} z = 0.8$, and the point $z_0 = 0$ under F ($t = 2$) over its boundary at $t = 0$ (gray). The arrow is the path $t \mapsto H_t(z)$ of one point, from z (dot) to $F(z)$. The map is

$$F(z) = \frac{e^z - 1}{e^z + 1}.$$




Figure 4: Phase portrait of H_t on $[-7,7] \times [-5,5]$: the color shows $\arg H_t$ (key at right), the thin lines are the level sets $\arg H_t \in \frac{\pi}{6}\mathbb{Z}$, and the shading steps in $|H_t|$ by factors of $e^{2\pi/12} \approx 1.69$; the white line is $|H_t| = 1$. Zeros (dots) at -1.843. Poles (open circles) at $1.843 + \pi i$ and $1.843 - \pi i$. Shown at $t = 1.20$, where H_t deforms the identity ($t = 0$) into F ($t = 2$) along the homotopy H_t, $0 \le t \le 2$, that runs the steps in turn ($H_t = H^{(k)}_{t-k+1}$ on $k-1 \le t \le k$): for f_1, the zoom homotopy $H^{(1)}_s(z) = (e^{sz}-1)/s + s$ for $0 < s \le 1$, $H^{(1)}_0(z) = z$; for f_2, the elliptic Möbius transformations $H^{(2)}_s$ that fix $p = i$ and $q = -i$, with $H^{(2)\prime}_s(p) = e^{-\pi i s/2}$. The map is

$$F(z) = \frac{e^z - 1}{e^z + 1}.$$