

Figure 2 Tumor cells in the presence of 5 gm of a sublethal dose

S Kale, R Raja, D Thorat, G Soundararajan, TV Patil
and GC Kundu

University of Texas Southwestern Medical Center

Figure 3. Tumor cells are detected by ELISA enzymatic reaction of a sub-lethal dose of P3. Tumor cells are detected by ELISA enzymatic reaction of phosphodiesterase inhibitor 1 (P3) for 24 h and then washed three times. The proteins are expressed as a percentage of the percentage of the cells constituting the TUMOR cell fraction. (A) Tumor cells were detected by ELISA enzymatic reaction of A1-P3 with a total of 60 ng/ml of phosphodiesterase inhibitor 1 (P3) for 24 h and then washed three times. The proteins are expressed as a percentage of the percentage of the cells constituting the TUMOR cell fraction. (B) Tumor cells were detected by ELISA enzymatic reaction of A1-P3 with a total of 60 ng/ml of phosphodiesterase inhibitor 1 (P3) for 24 h and then washed three times. The proteins are expressed as a percentage of the percentage of the cells constituting the TUMOR cell fraction. (C) Tumor cells were detected by ELISA enzymatic reaction of A1-P3 with a total of 60 ng/ml of phosphodiesterase inhibitor 1 (P3) for 24 h and then washed three times. The proteins are expressed as a percentage of the percentage of the cells constituting the TUMOR cell fraction. The TUMOR cell fraction was measured using a liquid chromatography-mass spectrometry method. (A) Tumor cells were detected by ELISA enzymatic reaction of A1-P3 with a total of 60 ng/ml of phosphodiesterase inhibitor 1 (P3) for 24 h and then washed three times. The proteins are expressed as a percentage of the percentage of the cells constituting the TUMOR cell fraction. (B) Tumor cells were detected