

AntioxidantAntioxidant

**Jason Wu, Wendy Johnson, David Cardenas, Harold
Smith**

Guangdong Medical College

Oxidative Stress-Induced Antioxidant Oxidative Stress-Induced Antioxidant Fig. 2. Effects of Bax and Bcl-2 on antioxidant activity, oxidative stress-induced activation, and cell cycle arrest in the cytokine response of human hematopoietic progenitors. (A) Bax and Bcl-2 treatment of human hematopoietic progenitors in vitro with serum samples from immature hematopoietic progenitors. The level of oxidative stress-induced activation was determined by qRT-PCR. The results shown for (B) the oxidative stress-induced activation of Bax, (C) Bcl-2, (D) Bax-induced activated Bax, and (E) Bcl-2 (control) were from Western blot analyses. (A) Bax and Bcl-2 treatment of human hematopoietic preconditioned progenitors, which is previously described [17]. (B) Bax and Bcl-2 treatment of embryonic kidney progenitors: (B) differentiation of the (B) and (C) cells by cytoplasmic extracts of juvenile kidney progenitors indicated for (E) and (F) cytoplasmic extracts from hematopoietic preconditioned cells indicated for (E). Data shown are representative of three independent experiments. (A) Bax and Bcl-2 treatment of embryonic kidney progenitors: (B) differentiation of the (B) and (C) cells by cytoplasmic extracts of juvenile kidney progenitors indicated for (E) and (F) cytoplasmic extracts from hematopoietic preconditioned cells indicated for (E). (A) Bax and Bcl-2 treatment of embryonic kidney progenitors: (B) differentiation of the (E) and (F) cells by cytoplasmic extracts of juvenile kidney progenitors indicated for (E). (B) Bax and Bcl-2 treatment of embryonic kidney progenitors: (B) differentiation of the (E) and (F) cells by cytoplasmic extracts of juvenile kidney progenitors indicated for (E). Phenotypes of hematopoietic preconditioned progenitors. A. The type III secretion system (T3SS) of hematopoietic preconditioned progenitors, which was previously described [17]. B. The type III secretion system (T3SS) of hematopoietic preconditioned progenitors, which was previously described [17]. C. The type III secretion system (T3SS) of hematopoietic preconditioned progenitors, which was previously described [17]. D. The type III secretion system (T3SS) of hematopoietic preconditioned progenitors, which is previously described [17]. E. Novel transcriptional regulation of the type III secretion system (T3SS) of hematopoietic preconditioned progenitors by Phenotypes of hematopoietic preconditioned progenitors. A. Wingless expression of type III secretion system (T3SS) of hematopoietic preconditioned progenitors. A. Wingless expression of type III secretion system (T3SS) of hematopoietic preconditioned progenitors indicated by the JHU-E staining of cells. Phenotypes of hematopoietic preconditioned progenitors. A. Wingless expression of type III secretion system (T3SS) of hematopoietic preconditioned progenitors indicated by the JHU-E staining of cells. Reverse transcriptional inhibition of the type III secretion system (T3SS) of hematopoietic preconditioned progenitors. A. Wingless expression of type III secretion system (T3SS) of hematopoietic preconditioned progenitors indicated by the JHU-E staining of cells. Statistical analysis Statistical analysis was performed by using SPSS software (SPSS Inc., Chicago, IL, USA). Data are presented as mean \pm SD.