

KRFirthSMHaidaraAMRabinovichandKSBackstromRo

**Christopher Spears, Christopher Adams, Dr. Stephen
Hess**

Logistical College of Chinese People's Armed Police Force

redox regulation by the retinoid receptor in mammals. *J. Immunol.* 163:2245-2256. L. K. Yang, S. T. S. Lee, Y. Y. E. Kim, S. E. Lee, S. S. Lee, N. H. Lee, Y. M. Kim, and J. L. H. Kim, "A novel receptor for retinoid receptors. *J. Biochem. Biophys. Res. Commun.* 557:2282-2287. H. H. Lee, S. S. Lee, and D. S. Kim, "Retinoid receptors as targets for the retinoid receptor-mediated pathogenesis of mammalian malarial fever. *Curr. Opin. Pathol.* 10:257-266. M. M. Rabinovich, S. M. Haidara, S. M. Haidara, A. M. Rabinovich, and K. S. Backstrom, "Retinoid receptors as novel targets for the pathogenesis of human malarial fever. *J. Immunol.* 163:2187-2196. R. A. Saha, D. Saha, and M. A. Saha, "Retinoid receptor and pathogenesis in human malarial fever. *Nature* 454:962-906. H. M. Rabinovich, J. L. Haidara, A. Saha, and K. S. Backstrom, "Retinoid receptors and mechanism of pathogenesis in human malarial fever. *J. Immunol. Physiol.* 165:2213-2219. H. M. Rabinovich, J. L. Haidara, and K. S. Backstrom, "Retinoid receptor and pathogenesis in human malarial fever. *J. Immunol.* 165:2213-2219. K. S. Backstrom, "Retinoid receptors and pathogenesis in human malaria. *Ann. N. Y. Acad. Sci.* 2:275-291. M. A. Saha, D. Saha, and K. S. Backstrom, "Retinoid receptors and pathogenesis in human malaria. *Ann. N. Y. Acad. Sci.* 2:275-291. M. A. Saha, D. Saha, and K. S. Backstrom, "Retinoid receptors and pathogenesis in human malaria. *Ann. N. Y. Acad. Sci.* 2:275-291. Introduction Malaria is one of the most common causes of death in the world's population. Malaria is the leading cause of morbidity and mortality. A high prevalence of malaria is estimated to account for 60% of the world's population. The aim of this study was to investigate the role of the retinoid receptor in the pathogenesis of acute malaria. The objective was to investigate the mechanism by which retinoid receptors mediate MDR-mediated pathogenesis of acute malaria. Methods Caco-2 cells were cultured in DMEM and maintained in the absence of retinoid receptors. The Caco-2 cells were seeded in 24-well plates containing various concentrations of retinoid. The retinoic acid receptor (RAR) (1 : 10) was added to each well of the medium. The medium was replaced every 3 weeks, and the plates were incubated for 2 months (12 weeks). The retinoic acid receptor (receptor 1 : 5) was added every 3 weeks to each well of the medium. Each well was incubated for 1 year. The cells were then tested for the presence of retinoid receptors. The proteins were detected by immunoblotting with anti-6-methyl-RAR (1 : 10), and the proteins were detected by Western blot analysis using anti-rabbit IgG (1 : 10), anti-mouse IgA (1 : 10), and anti-mouse IgG (1 : 10). The protein concentration was adjusted.