

# **4542Effectof1PAOIon**

**Tony Alvarez, Daniel Mclaughlin, Megan Vazquez, Jill  
Brewer, Kayla Mills, Misty White, Nicole Murphy,  
Benjamin Clark, Allen Gonzalez**

**Paris Descartes, CPSC**

MCAO-induced MCAO cell killing cells MCAO-induced MCAO cell killing was induced by 1-1 PAOI at concentrations of 20 ng/ml and 2,000 ng/ml. The mRNA and protein levels of the two subunits were determined by RT-PCR analysis. 4.5.4.3 Effect of 1-PAOI on MCAO-induced mRNA polymerase activity MDA-MB-231 and MDA-MB-468 cells were transfected with 1-PAOI for 30 min at 37°C in 10 mM Tris-HCl, and the nucleotide and protein levels were measured by RT-PCR. The effect of proteinase activity on MCAO-induced mRNA polymerase activity was determined by Western blotting after treatment with 1-PAOI. 4.5.5.1 Effect of 1-PAOI on MCAO-induced mRNA polymerase activity The mRNA and protein levels of the two subunits were determined by RT-PCR analysis. The signal transduction pathways and mRNA stability were determined by Western blotting after treatment with 1-PAOI. 4.5.6 Effect of 1-PAOI on the MCAO-induced mRNA polymerase activity MDA-MB-231 and MDA-MB-468 cells were transfected with 1-PAOI at 37°C in 10 mM Tris-HCl, and the nucleotide and protein levels were measured by RT-PCR. The mRNA polymerase activity was determined by Western blotting after treatment with 1-PAOI. 4.5.7 Effect of 1-PAOI on mRNA and protein levels of MCAO-induced extracellular matrix Intracellular matrix (ECM) was isolated from MDA-MB-231 and MDA-MB-468 cells, by immunoprecipitation and western blotting. MCAO-induced extracellular matrix (ECM) was isolated from MDA-MB-231 and MDA-MB-468 cells, using a standard double-sandwich chromatographic analysis kit [19,20]. The cell extracts were prepared from the extracellular matrix (ECM) of MDA-MB-231 and MDA-MB-468 cells, and the mRNA and protein levels were determined by RT-PCR. 4.5.8 Effect of 1-PAOI on extracellular matrix The mRNA and protein levels of extracellular matrix (ECM) of MDA-MB-231 and MDA-MB-468 cells, were determined by Western blotting after treatment with 1-PAOI. 4.5.9 MCAO-induced extracellular matrix (ECM) The mRNA and protein levels of extracellular matrix (ECM) of MDA-MB-231 and MDA-MB-468 cells, were determined by Western blotting after treatment with 1-PAOI. 4.5.10 MCAO-induced extracellular matrix (ECM) The mRNA and protein levels of extracellular matrix (ECM) of MDA-MB-231 and MDA-MB-468 cells, were determined by Western blotting after treatment with 1-PAOI. 4.5.11 MCAO-induced extracellular matrix (ECM) The mRNA and protein levels of extracellular matrix (ECM) of MDA-MB-231 and MDA-MB-468 cells, were determined by Western blotting after treatment with 1-PAOI. 4.5.12 MCAO-induced extracellular matrix (ECM) The mRNA and protein levels of extracellular matrix (ECM) of MDA-MB-231 and MDA-MB-468 cells, were determined by Western blotting after treatment with 1-PAOI. 4.5.13 MCAO-induced extracellular matrix (ECM) The mRNA and protein levels of extracellular matrix (ECM) of MDA-MB-231 and MDA-MB-468 cells, were determined by Western blotting after treatment with 1-PAOI. 4.5.14 MCAO-induced extracellular matrix (ECM) The mRNA and protein levels of extracellular matrix (ECM) of MDA-MB-231 and MDA-MB-468 cells were determined by Western blotting after