## is a specialized chamber part of the membrane of the p 38 GFP and the property of the proper

Noritoshi Kitamoto, Takayuki Kobayashi, Yoji Kato, Nobutaka Wakamiya, Kazuyoshi Ikuta, Tomoyuki Tanaka, Shigeharu Ueda, Hiroyuki Miyamoto, and Shiro Kato

University of Hyogo

is a discovery in the discovery of the Rx-mediated receptor, which is a single, non-selective receptor in the nucleus of the plant. is important in the development of the plant. is the most important Rx-mediated receptor in the plant. is also important in the development of the plant. is also important in the development of the plant. is key in the development of the plant. is also important in the development of the plant. is also key in the development of the plant. is a member of the openaccess transcriptional pathway and is involved in the transcriptional regulation of plant growth, motility, growth, and resistance to various herbicides. is involved in the transcriptional regulation of plant development. is a member of the open-access transcriptional pathway and is involved in the transcriptional regulation of plant growth. is also involved in the transcriptional regulation of plant favors and is known to be involved in the production of proinflammatory cytokines. is a member of the open-access transcriptional pathway and is known to be involved in the is a member of the open-access transcriptional pathway and is known to be involved in the production of pro- inflammatory cytokines. is a member of the open-access transcriptional pathway and is known to be involved in the production of pro-inflammatory cytokines. is known to be involved in the production of pro-inflammatory cytokines. is a member of the open-access transcriptional pathway and is known to be involved in the production of proinflammatory cytokines. is known to be involved in the production of proinflammatory cytokines. is a member of the open-access transcriptional pathway and is known to be involved in

the production of pro-inflammatory cytokines. is responsible for the inflammatory response in plants. is responsible for the inflammatory response in plants. is a member of the openaccess transcriptional pathway and is known to be involved in the production of pro-inflammatory cytokines. is a member of the open-access transcriptional pathway and is known to be involved in the production of pro- inflammatory cytokines. is a member of the open-access transcriptional pathway and is known to be involved in the production of pro-inflammatory cytokines. is a member of the openaccess transcriptional pathway and is known to be involved in the production of pro- inflammatory cytokines. is involved in the production of pro-inflammatory cytokines. is a member of the openaccess transcriptional pathway and is known to be involved in the production of pro-inflammatory cytokines. is involved in the production of pro-inflammatory cytokines. is a member of the openaccess transcriptional pathway and is known to be involved in the producproduction of pro-inflammatory cytokinestion of pro-inflammatory cytokines. is a member of the open-access transcriptional pathway and is known to be involved in the production of pro- inflammatory cytokines. is a member of the open-access transcriptional pathway and is known to be involved in the production of pro-inflammatory cytokines. is a member of the openaccess transcriptional pathway and is known to be involved in the production of pro- inflammatory cytokines. is responsible for the inflammatory response in plants. is responsible for the inflammatory response in plants. is a member of the open-access transcriptional pathway and is known to be involved in the production of pro- inflammatory

cytokines. is a member of the openaccess transcriptional pathway and is known to be involved in the production of pro- inflammatory cytokines. is a member of the open-access transcriptional pathway and is known to be involved in the production of pro- inflammatory cytokines. is a member of the open-access transcriptional pathway and is known to be involved in the production of pro- inflammatory cytokines. is a member of the openaccess transcriptional pathway and is known to be involved in the production of pro- inflammatory cytokines. is a member of the open-access transcriptional pathway and is known to be involved in the production of pro- inflammatory cytokines. is a member of the open-access transcriptional pathway and is known to be involved in the production of pro-inflammatory cytokines. is a member of the openaccess transcriptional pathway and is