

The Core Oligosaccharide and Thioredoxin of *Vibrio cholerae* Are Necessary for Binding and Propagation of Its Typing Phage VP3[▽]

Jingyun Zhang,[†] Wei Li,^{†‡} Qian Zhang, Hongxia Wang, Xiao Xu, Baowei Diao, Lijuan Zhang, and Biao Kan^{*}

National Institute for Communicable Disease Control and Prevention and State Key Laboratory for Infectious Disease Prevention and Control, Chinese Center for Disease Control and Prevention, P.O. Box 5, Changping, Beijing 102206, Peoples Republic of China

Received 1 October 2008/Accepted 22 January 2009

VP3 is a T7-like phage and was used as one of the typing phages in a phage-biotyping scheme that has been used for the typing of *Vibrio cholerae* O1 biotype El Tor. Here, we studied the receptor and other host genes of *V. cholerae* necessary for the lytic propagation of VP3. Six mutants resistant to VP3 infection were obtained from the random transposon insertion mutant bank of the sensitive strain N16961. The genes VC0229 and VC0231, which belong to the *wav* gene cluster encoding the core oligosaccharide (OS) region of lipopolysaccharide, were found to be interrupted by the transposon in five mutants, and the sixth mutant had the transposon inserted between the genes *rhlB* and *trxA*, which encode the ATP-dependent RNA helicase RhlB and thioredoxin, respectively. Gene complementation, transcription analysis, and the loss of VP3 sensitivity by the gene deletion mutants confirmed the relationship between VP3 resistance and VC0229, VC0231, and *trxA* mutation. The product of VP3 gene 44 (gp44) was predicted to be a tail fiber protein. gp44 could bind to the sensitive wild-type strain and the *trxA* mutant, but not to VC0229 and VC0231 mutants. The results showed that OS is a VP3 receptor on the surface of N16961, thioredoxin of the host strain is involved in the propagation of the phage, and gp44 is the tail fiber protein of VP3. This revealed the first step in the infection mechanism of the T7-like phage VP3 in *V. cholerae*.

Vibrio cholerae is the pathogenic agent of cholera and has caused seven pandemics. Only the O1 and O139 serogroups of *V. cholerae* have been associated with cholera epidemics, and strains of the O1 serogroup are subdivided into two biotypes, classical and El Tor (12). *V. cholerae* is the host for many bacteriophages, such as the tailed phage CP-T1 (18), K139 (39), and many filamentous phages (3, 24, 49). Phages infect and destroy host strains with high specificity, so phage-typing schemes were developed to differentiate strains of the same serogroups or genera in many bacteria, such as *Salmonella enterica* serovar Typhi (28) and *Staphylococcus aureus* (22). In *V. cholerae*, phage-typing schemes were also developed for typing of the O1 biotype El Tor (7) and O139 (5). A phage-biotyping scheme has been used for nearly 40 years in China for the typing of O1 El Tor strains (14). Based on the lytic patterns of the five typing phages, El Tor strains can be clustered into 32 phage types.

The first step of phage infection is the specific binding of phages to receptors on the host cell surface, triggering the ejection of phage DNA into the cell and initiating the propagation process (1). Some outer membrane proteins serve as receptors for phage adsorption, such as LamB (19), OmpA (8),

and OmpC (52) of *Escherichia coli* and OmpK of *Vibrio parahaemolyticus* (23). Lipopolysaccharide (LPS) is a common phage receptor. Cytotoxin-converting Φ CTX from *Pseudomonas aeruginosa* (51) and FC3-10 from *Klebsiella* spp. (4) bind to the core region of LPS, while vibriophage CP-T1 (17) and Φ YeO3-12, specific for *Yersinia enterocolitica* serotype O:3 (35), bind to the O side chain. Other structures that can be used as receptors include flagella (32) and pili (49). Besides the bacterial receptors, there are some other host components necessary for the propagation of virulent phages, such as the host RNA polymerase involved in the transcription of phage genes and the components participating in the assembly of phage particles.

VP3 is one of the five typing phages in the phage-biotyping scheme (14). The complete genome sequence comprises 39,481 bp with an overall G+C content of 42.62%, and 52 open reading frames (ORFs) were predicted (our unpublished data). It is a T7-like phage by morphology and genome sequence comparison. Here, we have identified the receptor and other host genes of *V. cholerae* that are necessary for the lytic propagation of VP3. In addition, VP3 tail fiber protein was also predicted and proved to function as a receptor-binding protein.

MATERIALS AND METHODS

Bacterial strains, plasmids, and media. The strains and plasmids used in this study are listed in Table 1. Phage VP3 was propagated on host strain 2477c. The El Tor strain N16961, whose genome was sequenced (20), is sensitive to VP3. Streptomycin (Sm)-resistant N16961-Sm^r was selected on Luria broth (LB) agar with Sm (100 μ g/ml) and was also sensitive to VP3. All strains were grown in liquid or solid (15 g/liter agar) LB medium, which could be supplemented with kanamycin (Kan) (50 μ g/ml), Sm (100 μ g/ml), chloramphenicol (Cm) (15 μ g/ml),

^{*} Corresponding author. Mailing address: Department of Diarrheal Diseases, National Institute for Communicable Disease Control and Prevention, Chinese Center for Disease Control and Prevention, P.O. Box 5, Changping, Beijing 102206, Peoples Republic of China. Phone: 86-10-61739458. Fax: 86-10-61739156. E-mail: kanbiao@icdc.cn.

[†] J.Z. and W.L. contributed equally to this work.

[‡] Present address: Laboratory of Nucleic Acid Vaccines, Department of Medicine, University of Massachusetts Medical School, Worcester, MA.

[▽] Published ahead of print on 6 February 2009.

TABLE 1. Strains and plasmids used in this study

Strain or plasmid	Description	Source
<i>V. cholerae</i>		
2477c	O1 El Tor, Ogawa	Laboratory stock
N16961	O1 El Tor, Inaba	Laboratory stock
N16961-Sm ^r	Spontaneous mutant of N16961; Sm ^r	This study
N44G1	N16961-Sm ^r VC0229::mini-Tn5Km2 Sm ^r Kan ^r VP3 ^r	This study
N67B1	N16961-Sm ^r VC0231::mini-Tn5Km2 Sm ^r Kan ^r VP3 ^r	This study
N69H3	N16961-Sm ^r VC0229::mini-Tn5Km2 Sm ^r Kan ^r VP3 ^r	This study
N77C6	N16961-Sm ^r VC0229::mini-Tn5Km2 Sm ^r Kan ^r VP3 ^r	This study
N93C3	N16961-Sm ^r VC0231::mini-Tn5Km2 Sm ^r Kan ^r VP3 ^r	This study
N60C6	N16961-Sm ^r mutant with mini-Tn5Km2 inserted between <i>rhlB</i> and <i>trx4</i> ; Sm ^r Kan ^r VP3 ^r	This study
C29	N16961ΔVC0231	This study
N16961-d0306	N16961Δ <i>trx4</i>	This study
<i>E. coli</i>		
JM109	<i>endA1 recA1 gyrA96 thi-1 hsdR17</i> (r _K ⁻ m _K ⁻) <i>relA1 supE44</i> °(<i>lac-proAB</i>) [F' <i>traD36 proAB lacI</i> °Z ^M M15]	TaKaRa
SM10(<i>λpir</i>)	<i>thi thr leu tonA lacY supE recA</i> ::RP4-2-TC::Mu Km <i>λpir</i>	44
SM10-km2	SM10(<i>λpir</i>) carrying plasmid pUTkm2; Kan ^r Amp ^r	This study
BL21(DE3)	F ⁻ <i>ompT hsdSB</i> (r _B ⁻ m _B ⁻) <i>gal dcm</i> (DE3)	Tiagen
Plasmids		
pUTkm2	Suicide plasmid carrying transposon mini-Tn5Km2; Kan ^r Amp ^r TnpA ⁺	10
pET30-a(+)	Expression vector; <i>fl ori</i> T7 <i>lac</i> promoter Kan ^r	Novagen
pUC19	Cloning vector; <i>ori lacZ</i> Amp ^r	TaKaRa
p0231UL	pUC19 carrying the upstream and downstream fragments flanking VC0231	This study
pDS132	Suicide plasmid; <i>R6K ori mob RP4 cat sacB</i>	37
pDS-0231	pDS132 carrying the upstream and downstream fragments flanking VC0231	This study
pTcat	pMD18-T::cat	29
p0306U-cat-L	pTcat carrying the upstream and downstream fragments flanking <i>trx4</i>	This study
pCVD442	Suicide plasmid; <i>R6K ori mob RP4 bla sacB</i>	11
pCVD-0306	pCVD442 with <i>cat</i> inserted between the upstream and downstream fragments flanking <i>trx4</i>	This study
pBR322	Cloning vector; ColEI <i>ori</i> Amp ^r Tet ^r	TaKaRa
pBR0229c	pBR322 carrying gene VC0229	This study
pBR0231c	pBR322 carrying gene VC0231	This study
pBR0305c	pBR322 carrying gene <i>rhlB</i>	This study
pBR0306c	pBR322 carrying gene <i>trx4</i>	This study
pBR0305-6c	pBR322 carrying both gene <i>rhlB</i> and <i>trx4</i>	This study

or ampicillin (100 µg/ml). During the deletion mutant construction, solid LB medium with 20% sucrose and without NaCl was used to select plasmid excision from the chromosome.

Propagation of phage VP3 and DNA extraction. Twenty milliliters of 2477c culture (1×10^8 CFU/ml) and VP3 (1×10^8 PFU) were added to 80 ml of LB, incubated at 37°C, and shaken until the lysate was clear. The mixture was incubated at 56°C for 30 min and then centrifuged to remove cell debris. With a titer of 10^9 to 10^{10} PFU/ml, the supernatant was stored at 4°C as a stock solution. When phage DNA was extracted (40), the supernatant was treated with DNase I and RNase A, and then solid NaCl and polyethylene glycol 8000 were added, and it was left overnight at 4°C. Precipitated phage particles were recovered by centrifugation and were resuspended in 2 ml of SM buffer (40). The polyethylene glycol 8000 and cell debris were extracted from the phage suspension by adding an equal volume of chloroform, and it was centrifuged to recover the aqueous phase. Proteinase K and sodium dodecyl sulfate (SDS) were added to the solution, and it was incubated at 56°C for 1 h. An equal volume of phenol was added to the chilled phage suspension, mixed, and centrifuged at $3,000 \times g$ for 5 min at room temperature. The aqueous phase was extracted with a 1:1 mixture of equilibrated phenol and chloroform and an equal volume of chloroform. DNA was precipitated with ethanol and dissolved in deionized water.

Transposon mutagenesis and the selection of VP3-resistant mutants. Plasmid pUTkm2 (10) was transformed into *E. coli* SM10(*λpir*) (44) to obtain SM10-km2 (Kan^r). Conjugation was performed between N16961-Sm^r as the recipient strain and SM10-km2 as the donor strain. The donor and recipient strain (optical density at 600 nm [OD₆₀₀] ≈ 0.8) cultures were mixed 1:9 in a 1.5-ml microcen-

trifuge tube, washed twice with 1 ml of LB, resuspended in 100 µl of LB, and transferred onto 0.45-µm-pore-size membranes overlaid on LB plates. After being incubated at 37°C for 4 to 6 h, the membranes were washed with LB and the collected bacteria were diluted and plated onto the LB agar plates (Kan, 50 µg/ml, and Sm, 100 µg/ml) to select mutants.

Separated transconjugants were picked and inoculated into 150 µl of LB (Kan, 50 µg/ml, and Sm, 100 µg/ml) contained in 96-well plates (Corning Costar 3599) and incubated for 3 to 5 h until the OD was 0.5 to 0.8. This was monitored with a Tecan Genios Basic Microplate Reader at 395 nm. Then, 10 µl of the cultures was inoculated into 140 µl of LB (Kan, 50 µg/ml, and Sm, 100 µg/ml) with phage VP3 (1×10^8 PFU/ml) contained in new 96-well plates and incubated for 3 h. The wells with an OD₃₉₅ of 0.5 or more were selected as the candidates for VP3-resistant mutants, which were confirmed subsequently by double-layer plaque assay. Briefly, 4 ml of melted 0.7% LB agar was mixed with 100 µl of cell cultures and poured onto an LB agar plate, and 10 µl of VP3 10-fold serial dilutions was dropped onto the plate when the upper layer solidified. After overnight incubation, mutants with no plaque formation were confirmed as VP3 resistant. The Kan resistance gene (*kan*) was detected by PCR with the primers pkan-low/pkan-up, and the transposase gene *tnpA* was detected with the primers ptnpA-low/ptnpA-up (Table 2). Strains with *kan* and without *tnpA* had the transposon inserted into chromosomes.

Southern blot analysis and determination of transposon insertion sites. Southern blot analysis was performed by a standard procedure (40). Total DNA was extracted from mutants and digested with restriction enzymes, such as EcoRI, PstI, KpnI, and SacI, and separated by electrophoresis. The fragment

TABLE 2. Primers used in this study

Primer	Sequence (5'–3') ^a	Restriction site
Detection of <i>kan</i> and <i>tnpA</i>		
pkan-low	TGGCAAGATCCTGGTATCGGT	No
pkan-up	CAATCAGGTGCGACAATCTAT	No
ptnpA-low	AGGACAAACTGGCGCATAAC	No
ptnpA-up	TAAACAGCTGACCGCAACAA	No
Used in mutagenesis of VC0231 of N16961		
m0231U-1	CTCGGTACCTTCTCGGCTGTCTGAATC	KpnI
m0231U-2	CGGGGATCCGATGATCGTCGCCGCAA	BamHI
m0231U-3	TCCTCTAGATTCTCGGCTGTCTGAATC	XbaI
m0231L-1	CGGGGATCCAGCGACTTGATGCTGAAA	BamHI
m0231L-2	CAGGCATGCTGGTTTTCTGATGCTCAA	SphI
inner0231-1	TATGATGGCTCTGTTCCG	No
inner0231-2	ATTGACCTCCAGTGGCTT	No
Used in mutagenesis of <i>trxA</i> of N16961		
m0306U-1	TTCGAGCTCCTTGAATGTGATTGAGGTTG	SacI
m0306U-2	CGGGGATCCCTTTCACTCCAATGTGAT	BamHI
m0306L-1	AGAGTCGACTGAGTCAGACCGCCAAG	Sall
m0306L-2	CTTGCAATGCAGTGTTCACTTTGAGCAATG	SphI
inner0306-1	CGATGACGGTTTCGAGAA	No
inner0306-2	CTACCACACTGCCATCTT	No
Used in construction of complementary plasmids		
VC0229cU	GGGGATCCTTTTGGTATTTGTCACCT	BamHI
VC0229cL	GAGTCGACAGAATGGGAGCGAGGCGA	Sall
VC0231cU	GGGGATCCAGCTGTTGACCGAGTGTT	BamHI
VC0231cL	GAGTCGACTCCTATCATTACTGAGTG	Sall
VC0305cU	GGGGATCCTTGTCTGCACACTGCCAT	BamHI
VC0305cL	GAGTCGACTCGTTCTCGAAACCGTCA	Sall
VC0306cU	GGGGATCCGTCGGCGAACTTATGCTC	BamHI
VC0306cL	GAGTCGACATTAACGTGCGATAAACT	Sall

^a Restriction sites are underlined.

amplified from pUTKm2 with the primers pkan-low/pkan-up was labeled with digoxigenin using a DIG DNA Labeling and Detection Kit (Roche) and served as the DNA probe for hybridization. The blots were prehybridized for 2 h and hybridized for 12 h at 42°C in 50% formamide. If the restricted fragments containing the insert were of suitable size (2.5 kb to 4.4 kb in this study), they were ligated into pUC19 and sequenced to determine the transposon insertion sites.

Construction of deletion mutants and complementation experiments. An in-frame VC0231 deletion mutant of *V. cholerae* N16961 was constructed by homologous recombination using the suicide plasmid pDS132 (37). The flanking sequences on the 5' and 3' sides of the fragment that was to be deleted were PCR amplified from N16961 chromosomal DNA using the primers m0231U-1/m0231U-2 and m0231L-1/m0231L-2 (Table 2), respectively. Then, the fragments were cloned into pUC19, generating the plasmid p0231UL. The fragment containing the in-frame deletion pattern of VC0231 was obtained by PCR from a p0231UL template using primers m0231U-3/m0231L-2 (Table 2) and cloned into pDS132 to construct the plasmid pDS-0231, which was conjugally transferred into N16961 from the donor strain *E. coli* SM10(λ pir). Transconjugants were selected on thiosulfate-citrate-bile-sucrose (TCBS) agar with 15 μ g/ml Cm and streaked onto LB agar plates with 20% sucrose and without NaCl. Colonies from sucrose selection medium were screened by PCR using the primers inner0231-1/inner0231-2 (Table 2), which were specific for the deleted fragment. Sequencing confirmed that the negatively PCR-amplified colonies were mutants.

A *trxA* deletion mutant of N16961 was constructed using the suicide plasmid pCVD442 (11) in a similar way. The flanking sequences on the 5' and 3' sides of *trxA* were PCR amplified using the primers m0306U-1/m0306U-2 and m0306L-1/m0306L-2 (Table 2), respectively, and were subsequently cloned into pTcat (29), generating the plasmid p0306U-cat-L. The plasmid pTcat was constructed previously in our laboratory by inserting a *cat* gene, which confers Cm resistance, on pMD18-T (TaKaRa). The plasmid p0306U-cat-L was restricted with SacI and SphI, and the 2-kb band containing the in-frame deletion pattern of *trxA* was purified and cloned into pCVD442 to construct the plasmid pCVD-0306. The following procedures were similar to those for the construction of the VC0231

deletion mutant, except that the sucrose selection medium contained Cm (15 μ g/ml). Mutants were screened using the primers inner0306-1/inner0306-2 (Table 2) and were confirmed by sequencing.

When the complementary plasmid pBR0231c was constructed, the fragment containing VC0231 was PCR amplified from chromosomal DNA of N16961 with the primers VC0231cU/VC0231cL and inserted into pBR322. In the same way, the plasmids pBR0229c, pBR0305c, pBR0306c, and pBR0305-6c were constructed with the primers VC0229cU/VC0229cL, VC0305cU/VC0305cL, VC0306cU/VC0306cL, and VC0305cU/VC0306cL (Table 2), respectively. The fragment inserted into pBR0305-6c covered the genes *rhIB* and *trxA*. The complementary plasmids were transformed into different mutants, as mentioned in Results. The plasmid pBR322 was transformed into the mutants as a control. The sensitivity of the complemented mutants was examined by double-layer plaque assay.

Detection of gene transcription. The transcription of *rhIB* and *trxA* was detected by reverse transcription (RT)-PCR. Total RNA was extracted with a Qiagen RNeasy Mini Kit from N16961, N60C6, and N60C6 complemented with different plasmids and reverse transcribed with SuperScript III Reverse Transcriptase (Invitrogen). The resulting cDNAs were used as PCR templates amplified with gene-specific primers.

Expression, purification, and FITC labeling of the predicted VP3 tail fiber protein. A DNA fragment containing gene 44, which was predicted to encode VP3 tail fiber protein, was PCR amplified from VP3 DNA with the primers gp44NB1 (5'-TATACATATGCACCATCATCATCATTCAGGCACTCGTGCTCCT-3') and gp44NB2 (5'-CGCACTCGAGTTAATTTAAAGGGATAGT-3'). The former contained the coding sequence of a hexahistidine tag and the restriction site for NdeI, and the latter contained the restriction site for XhoI (both underlined). The PCR product was digested with NdeI and XhoI and inserted into NdeI/XhoI-digested pET30-a (+). The resulting plasmid was transformed into the host strain *E. coli* BL21(DE3). Expression of His₆-gp44 was induced with 0.4 mmol/liter isopropyl- β -D-thiogalactopyranoside (IPTG) at 28°C for 2 to 3 h. The total protein of the induced strain was separated by SDS-polyacrylamide gel electrophoresis (PAGE) and detected by Western blotting using mouse anti-His antibody at a concentration of 1:1,000 and horseradish




FIG. 1. Transposon insertion in the *wav* gene cluster and VP3 infection of the mutants. (A) Organization of the *wav* gene cluster of N16961 and the transposon insertion sites of the five VP3-resistant mutants. The large black arrows represent *wav* genes that were assumed to be essential. The white arrows represent genes that interfere with O-antigen attachment (34, 41). The grey arrows represent genes in the *wav* gene cluster. The transposon insertion positions are marked with small black arrows. The numbers after the mutant names stand for the nucleotide (nt) site just following the transposon in chromosome I of *V. cholerae* N16961. (B) Double-layer plaque assay of mutants (N67B1, N77C6, and C29) and complemented mutants (N67B1-vc0231, N77C6-vc0229, and C29-vc0231).

peroxidase-conjugated goat anti-mouse secondary antibody at a concentration of 1:500. His₆-gp44 was purified with a His₆ Bind Purification Kit (Novagen) and labeled with a fluorescein isothiocyanate (FITC) labeling kit (Guangzhou Chang Rui Biotech, China) according to the manufacturer's instructions. His₆-gp44 labeled with FITC is referred to as FITC-gp44 below.

Adsorption of FITC-gp44 to strains. Strains were cultured in LB at 37°C until the OD₆₀₀ was 0.3 to 0.4. Cells in 1 ml of the culture were collected by centrifugation at 4,000 × g for 4 min, washed once with phosphate-buffered saline (PBS), incubated for 1 h in FITC-gp44 solution, washed twice with PBS to remove the FITC-gp44 not adsorbed by the cells, and finally suspended in PBS. Two negative controls were made; one was *E. coli* BL21(DE3) treated with FITC-gp44 in parallel to the test strains, and the other was N16961 treated with PBS instead of the FITC-gp44 solution. To detect the dose dependence of agglutination of N16961 by FITC-gp44, cells of N16961 were incubated with twofold serial dilutions (1:2 to 1:32) of FITC-gp44 in PBS. All procedures were carried out at 4°C. After the FITC-gp44 solution was added, the samples were kept in the dark.

Each sample was mixed with an equal volume of 0.5% agarose preheated to 50°C, and 4 μl of the mixture was immediately dropped onto a slide, covered with a coverslip, and then observed with a fluorescence microscope (Olympus; BX51) and by confocal laser scanning microscopy (CLSM) (Olympus; FV500).

Nucleotide sequence accession number. The GenBank accession number for VP3 gene 44 is FJ217960.

RESULTS

Selection of VP3-resistant transposon insertion mutants and determination of the insertion sites. Nine thousand one hundred insertion mutants of strain N16961-Sm^r were generated by transposon mutagenesis with plasmid pUTKm2. By monitoring the growth rates of the mutants with the microplate reader when VP3 was coinoculated and testing VP3 resistance by double-layer plaque assay, we finally obtained six VP3-resistant mutants (N44G1, N60C6, N67B1, N69H3, N77C6, and N93C3). These mutants were positive for *kan* and negative for *tnpA* by PCR amplification, suggesting the transposon had been inserted into chromosomes. Southern blot analysis showed a single band, implying each mutant had one copy of the transposon inserted. To clone the transposon insertion fragments in the mutants, the restriction fragments containing the transposon insert, which were 2.5 kb to 4.4 kb in size in the Southern blots, were inserted into vector pUC19 and sequenced. The mutation sites of three mutants (N77C6, N67B1,

and N60C6) were determined by this method. In mutants N77C6 and N67B1, the transposon was inserted into the genes VC0229 and VC0231, respectively, both belonging to the *wav* gene cluster responsible for the synthesis of the core oligosaccharide (OS) region of LPS (34). In mutant N60C6, the transposon was inserted into the interval region between *rhlB* (locus tag VC0305), encoding the ATP-dependent RNA helicase RhlB, and *trx4* (locus tag VC0306), encoding thioredoxin (see Fig. 2A). The two genes are transcribed in opposite directions, and the insertion site lies in the 5' noncoding regions of both genes. Primers were then designed flanking the genes mentioned above and were used to detect the other three mutants (N44G1, N69H3, and N93C3) by PCR. We found that the transposon was inserted into VC0229 in mutants N44G1 and N69H3 and into VC0231 in mutant N93C3. The transposon was inserted at different positions within the ORFs of VC0229 and VC0231 in these mutants (Fig. 1A). By random transposon mutagenesis, four genes that are probably necessary for VP3 infection were screened out, i.e., *rhlB*, *trx4*, and two genes in the *wav* gene cluster. We then studied the effects of the genes on VP3 infection.

The *wav* gene cluster of *V. cholerae* N16961 is related to VP3 infection. To further determine the relationship between VP3 resistance and the functions of genes in the *wav* cluster, a deletion mutant of N16961 in VC0231 (C29) was constructed by homologous recombination. Strain C29 showed resistance to VP3 when examined by double-layer plaque assay. The mutants N67B1, N93C3, and C29 regained VP3 sensitivity when they were complemented with the plasmid pBR0231c, and so did N44G1, N69H3, and N77C6 when complemented with pBR0229c (Fig. 1B).

***V. cholerae* *trx4* encoding thioredoxin is relevant for VP3 infection.** In mutant N60C6, the transposon lies in the 5' noncoding regions of *rhlB* and *trx4*. To determine which gene (or both) is responsible for resistance to VP3 infection, three complementary plasmids were constructed and used to transform N60C6: pBR0305c, carrying *rhlB*; pBR0306c, carrying *trx4*; and pBR0305-6c, carrying both genes (Fig. 2B). N60C6 re-




FIG. 2. *trxA* and VP3 infection. (A) In mutant N60C6, a transposon was inserted into the region between *rhIB* (VC0305) and *trxA* (VC0306) in the 5' noncoding regions of both genes. nt., nucleotides. (B) Three complementary plasmids were constructed, each carrying a fragment containing *rhIB*, *trxA*, or both. The lines show the regions covered by the cloned fragments. (C) Response to VP3 infection by double-layer plaque assay. N60C6 became VP3 sensitive when it was transformed by pBR0306c (generating N60C6-vc0306), but not when the transforming plasmid was pBR0305c (generating N60C6-vc0305). (D) The transcription of *rhIB* and *trxA* was detected by RT-PCR. The lanes marked with a plus are positive controls.

gained VP3 sensitivity when it was transformed with pBR0306c or pBR0305-6c, but not pBR0305c (Fig. 2C). We then tested whether both genes were transcribed in N60C6. When tested by RT-PCR, *rhIB* was still transcribed while *trxA* was not. The transcription of *trxA* was detected when N60C6 was complemented with pBR0306c (Fig. 2D). These results indicate that *trxA* was inactivated in N60C6, which led to the VP3 resistance. The *trxA* deletion mutant (N16961-d0306) resisted VP3 infection, and it became VP3 sensitive when transformed with pBR0306c (Fig. 2C), further confirming that *trxA* is indispensable for VP3 propagation.

gp44 of VP3 can adsorb to VP3-sensitive strains. In order to find out whether the mutations influenced the adsorption of VP3 or the subsequent processes, the gene for VP3 tail fiber protein was predicted and expressed in vitro. The predicted gene 44 of VP3 encodes a protein (designated gp44) of 753 amino acid residues (aa), the N-terminal portion (aa 8 to 244) of which showed an identity of 48% with the N-terminal part of T7 gp17 (tail fiber protein; 553 aa) and showed high similarity to tail fiber proteins of other T7 family phages (Fig. 3). By searching in the Pfam database (13), VP3 gp44 was found to contain two Pfam-A (Phage_T7_tail and Collar) and one Pfam-B (Pfam-B_57397) domains. These domains are all common in phage tail fiber proteins, and domain Phage_T7_tail (corresponding to aa 15 to 244 of VP3 gp44) attaches the fibers

to phage particles in phage T7 (45). Therefore, gp44 was predicted to be a tail fiber protein of VP3 and probably capable of binding to receptors with its C terminus.

VP3 gp44 was expressed in *E. coli* BL21(DE3) with a six-His tag added to the N terminus. The purified His₆-gp44 formed two bands when separated on an SDS-PAGE gel; one was the predicted size, while the other was about double the size, and both contained the six-His tag as revealed by Western blotting. If the His₆-gp44 sample was mixed with an equal volume of 7 M urea before SDS-PAGE, the larger band grew weaker and the smaller band grew stronger, suggesting the larger band was the dimer of His₆-gp44.

His₆-gp44 was labeled with FITC and used to treat VP3-sensitive N16961. Upon observation with a fluorescence microscope, many green masses were found in the sample, and most bacteria were agglutinated. Green-fluorescent masses were not found for the two negative controls, N16961 treated with PBS and *E. coli* BL21(DE3) treated with FITC-gp44 (data not shown). A similar phenomenon was also found under CLSM. The agglutination of N16961 changed with the amount of FITC-gp44 added (Fig. 4). The less FITC-gp44 added, the weaker the fluorescence and the smaller the bacterial masses formed, which showed dose dependence of gp44 for agglutination of N16961. These results suggest that gp44 adheres to VP3-sensitive strains by binding to receptors on the cell surface, and the possible oligomerization of gp44 causes the cells




FIG. 3. Comparison of tail fiber protein sequences of phage T7, T3, ΦYeO3-12, gh-1, and VP3. T7, T3, ΦYeO3-12, and gh-1 are all T7-like phages. At the top are shown the identities and conservation of the full-length proteins. Darker blue dots mean higher identities. Lighter and higher yellow columns mean more conservation. The box shows multiple alignment of the N termini of the tail fiber proteins, where the similarity is particularly high.

to agglutinate. If so, instead of VP3 particles, gp44 can be used to determine whether a strain can adsorb VP3.

Mutants with inactivated genes in the *wav* gene cluster lost the ability to adsorb phage VP3. The adsorption of phage VP3 to the mutants was also examined by observing the binding of FITC-gp44 with a fluorescence microscope and by CLSM (Fig. 5). Mutants with inactivated genes in the *wav* gene cluster, including the five transposon mutants with the inactivated gene VC0229 or VC0231 and the deletion mutant C29, had no FITC-gp44 binding on the cell surface. The binding of FITC-gp44 was restored when the mutants were complemented with corresponding genes. Thus, mutations in the *wav* gene cluster probably altered the binding sites of VP3. Mutant N60C6 with inactivated *trxA* could adsorb FITC-gp44, suggesting that thioredoxin encoded by *trxA* is not involved in VP3 adsorption.

DISCUSSION

A phage-biotyping scheme is used in China for subtyping of the *V. cholerae* O1 El Tor strains. VP3 is one of the five typing phages included in the scheme, and based on its genome sequence and morphology (our unpublished data), it belongs to the T7 group. In order to gain more knowledge about the infection mechanism of VP3, this study investigated VP3 re-

ceptors on the El Tor strain and other host genes necessary for VP3 propagation.

Many members of the T7 group use LPS as a receptor. The receptor of *Y. enterocolitica* phage ΦYeO3-12 is the O chain of serotype O:3, which consists of the rare sugar 6-deoxy-L-altropyranose (35); the receptor of lytic *Pseudomonas putida* bacteriophage gh-1 is proposed to be LPS, as well (27); and the LPS core is the binding site for T7 (38). The *wav* gene cluster of N16961, responsible for the synthesis of the OS region of LPS, lies in the 17.2-kb region between ORFs VC0223 and VC0240, including a group of glycosyl transferases and O-antigen ligase (Fig. 1A). It has been reported that *waaA* (VC0233), *wavC* (VC0227), and *waaC* (VC0225) are highly conserved and putatively essential core OS genes in *V. cholerae* (34). Among all the *wav* genes, only mutations in *waaL* (VC0237), *wavA* (VC0223), and *wavL* (VC0239) affect O antigen attachment (34, 41). Here, VC0229 and VC0231 were found to be relevant for N16961 sensitivity to phage VP3 in the process of phage attachment. Since genes related to O-antigen synthesis in these VC0229 and VC0231 insertion mutants are integral, O antigen is not proposed to function as the VP3 receptor. The exact functions of VC0229 and VC0231 are still unknown. VC0231 encodes a protein predicted to contain the catalytic domain of serine/threonine protein kinases. Genes encoding similar pro-




FIG. 4. Dose dependence analysis of FITC-gp44 for agglutination of N16961. Cells of N16961 were treated with twofold serial dilutions of prepared FITC-gp44 (1:2 to 1:32) and observed by CLSM. The dilution factors of FITC-gp44 are shown below.




FIG. 5. Binding of FITC-gp44 on the surfaces of different strains observed by CLSM. *E. coli* BL21(DE3) treated with FITC-gp44 was used as a negative control. N60C6 is an N16961 mutant with a transposon inserted into the intergenic region between *rhlB* and *trxA*. N67B1 and N77C6 are N16961 mutants with a transposon inserted into VC0231 and VC0229, respectively. C29 is an N16961 mutant with an in-frame deletion in VC0231. N67B1-vc0231, N77C6-vc0229, and C29-vc0231 are mutants with complementary plasmids.

teins exist in the *V. parahaemolyticus* O- and K-antigen biosynthesis gene cluster (GI193787939) and the *Actinobacillus suis* serotype K1 capsular gene cluster (GI29469153) ORF 7 (2). We predicted that the product of VC0231 possibly affects the activities of some enzymes needed in the process of OS synthesis. The OS was concluded to be the VP3 receptor on strain N16961.

Our results also implicate the gene *trxA*, encoding thioredoxin, in VP3 infection. In phage T7, *E. coli* thioredoxin is the processivity factor of T7 DNA polymerase (the gene 5 product of T7), increasing the processivity of nucleotide polymerization (31, 46), and influences the binding of T7 helicase to a DNA polymerase-thioredoxin complex (15). The situation may be similar for DNA polymerase of phage T3 (9). *E. coli* mutants with inactivated *trxA* become resistant to T7 (21, 38) but adsorb T7 normally (6). Similarly, thioredoxin is probably involved in the life process of VP3 as a subunit of VP3 DNA polymerase but is not involved in the binding of VP3. This supposition was also supported by the binding of FITC-gp44 on the surface of N60C6, which suggested that the inactivation of *trxA* did not alter the adsorption sites of phage VP3.

The tail fiber protein of a tailed phage has the receptor-binding function and decides the host specificity of the phage (16, 47, 50). VP3 gp44 is a homologue of tail fiber protein gp17 of phage T7, especially the proximal part. Three domains are predicted in gp44, and the proximal domain, Phage_T7_tail, is common in T7 family members, such as bacteriophage T7, T3 (36), K1F (43), and *Pseudomonas* phage gh-1 (27), and it links tail fibers to phage particles (45). The other two domains are also common in phage tail proteins. Domain Collar exists in the short tail fiber protein gp12 and the long tail fiber protein gp37 responsible for the receptor recognition of bacteriophage T4 (33, 48), while PfamB_PB057397 exists in ORF35 (tail fiber protein) of *Vibrio* phage K139 and some K139-like phages, such as Ch457 and E8498 (25). Therefore, gp44 of phage VP3 was predicted to be the tail fiber protein and to have a recep-

tor-binding function; this was validated by the binding of gp44 to the surface of wild-type N16961. N16961 mutants with mutations in the *wav* gene cluster did not bind gp44, further proving the interaction of gp44 with the OS region of LPS.

We found that the expressed His₆-gp44 tended to form oligomers, and strong denaturing conditions promoted the dissociation of the oligomer. In addition, fluorescence microscopy and CLSM showed that when FITC-gp44 was added, cells of VP3-sensitive strains agglutinated while cells of VP3-resistant *wav* gene mutants were dispersed. When N16961 and N60C6 were blended with a His₆-gp44 solution dropped onto a slide, the cells were agglutinated, just like the reaction with *V. cholerae* antiserum, while *E. coli* BL21(DE3) and the *wav* gene mutants were not agglutinated (data not shown). These results implied that His₆-gp44 is oligomeric in solution, which is the common form for phage tail fiber proteins. For example, T4 short tail fibers are trimeric gp12 (30), and tail fibers of T3 (26) and T7 (45) are trimers of the gp17 monomer. The agglutination of VP3-sensitive strains could be caused by the oligomerization of His₆-gp44.

Conclusion. Some bacteria cannot be lysed by a phage because they do not adsorb the phage particles, owing either to a lack of receptors on the cell surface or to the fact that the receptors are not accessible due to other cell surface components (42). Some bacteria are able to adsorb phages but fail to serve as host cells for phage multiplication if they lack any components necessary for phage propagation (1). We confirmed that mutants with an altered *wav* gene cluster resisted the virulent infection of VP3 because they could not adsorb the phage particles through the interaction between VP3 gp44 and the OS of *V. cholerae*. Thioredoxin, encoded by *trxA*, is necessary for VP3 proliferation and may be a subunit of VP3 DNA polymerase. These results can help us to understand the mechanism of T7-like phage infection of *V. cholerae*.

ACKNOWLEDGMENTS

This research was supported by the National Basic Research Priorities Program (grant G1999054102) and partially by the High Tech Research and Development Program (grant 2006AA02Z425) of the Ministry of Science and Technology, Peoples Republic of China.

REFERENCES

- Adams, M. H. (ed.). 1959. Bacteriophages. Interscience Publishers, Inc., New York, NY.
- Boyce, J. D., J. Y. Chung, and B. Adler. 2000. Genetic organisation of the capsule biosynthetic locus of *Pasteurella multocida* M1404 (B:2). *Vet. Microbiol.* 72:121–134.
- Campos, J., E. Martinez, E. Suzarte, B. L. Rodriguez, K. Marrero, Y. Silva, T. Ledon, R. del Sol, and R. Fando. 2003. VGJΦ, a novel filamentous phage of *Vibrio cholerae*, integrates into the same chromosomal site as CTXΦ. *J. Bacteriol.* 185:5685–5696.
- Camprubi, S., S. Merino, V. J. Benedi, and J. M. Tomas. 1991. Isolation and characterization of bacteriophage FC3-10 from *Klebsiella* spp. *FEMS Microbiol. Lett.* 67:291–297.
- Chakrabarti, A. K., A. N. Ghosh, G. B. Nair, S. K. Niyogi, S. K. Bhattacharya, and B. L. Sarkar. 2000. Development and evaluation of a phage typing scheme for *Vibrio cholerae* O139. *J. Clin. Microbiol.* 38:44–49.
- Chamberlin, M. 1974. Isolation and characterization of prototrophic mutants of *Escherichia coli* unable to support the intracellular growth of T7. *J. Virol.* 14:509–516.
- Chattopadhyay, D. J., B. L. Sarkar, M. Q. Ansari, B. K. Chakrabarti, M. K. Roy, A. N. Ghosh, and S. C. Pal. 1993. New phage typing scheme for *Vibrio cholerae* O1 biotype El Tor strains. *J. Clin. Microbiol.* 31:1579–1585.
- Cole, S. T., U. Chen-Schmeisser, I. Hindennach, and U. Henning. 1983. Apparent bacteriophage-binding region of an *Escherichia coli* K-12 outer membrane protein. *J. Bacteriol.* 153:581–587.
- Davidson, J. F., R. Fox, D. D. Harris, S. Lyons-Abbott, and L. A. Loeb. 2003.

- Insertion of the T3 DNA polymerase thioredoxin binding domain enhances the processivity and fidelity of *Taq* DNA polymerase. *Nucleic Acids Res.* **31**:4702–4709.
10. de Lorenzo, V., M. Herrero, U. Jakubzik, and K. N. Timmis. 1990. Mini-Tn5 transposon derivatives for insertion mutagenesis, promoter probing, and chromosomal insertion of cloned DNA in gram-negative eubacteria. *J. Bacteriol.* **172**:6568–6572.
 11. Donnenberg, M. S., and J. B. Kaper. 1991. Construction of an *eae* deletion mutant of enteropathogenic *Escherichia coli* by using a positive-selection suicide vector. *Infect. Immun.* **59**:4310–4317.
 12. Feeley, J. C. 1965. Classification of *Vibrio cholerae* (*Vibrio comma*), including El Tor vibrios, by infrasubspecific characteristics. *J. Bacteriol.* **89**:665–670.
 13. Finn, R. D., J. Mistry, B. Schuster-Bockler, S. Griffiths-Jones, V. Hollich, T. Lassmann, S. Moxon, M. Marshall, A. Khanna, R. Durbin, S. R. Eddy, E. L. L. Sonnhammer, and A. Bateman. 2006. Pfam: clans, web tools and services. *Nucleic Acids Res.* **34**:D247–D251.
 14. Gao, S., S. Wu, and B. Liu. 1984. Characteristics of typing phages of *Vibrio cholerae* biotype El Tor. *Fu Huo Luan Zi Liao Hui Bian* **4**:237–245.
 15. Ghosh, S., S. M. Hamdan, T. E. Cook, and C. C. Richardson. 2008. Interactions of *Escherichia coli* thioredoxin, the processivity factor, with bacteriophage T7 DNA polymerase and helicase. *J. Biol. Chem.* **283**:32077–32084.
 16. Grundy, F. J., and M. M. Howe. 1984. Involvement of the invertible G segment in bacteriophage mu tail fiber biosynthesis. *Virology* **134**:296–317.
 17. Guidolin, A., and P. A. Manning. 1985. Bacteriophage CP-T1 of *Vibrio cholerae*. Identification of the cell surface receptor. *Eur. J. Biochem.* **153**: 89–94.
 18. Guidolin, A., G. Morelli, M. Kamke, and P. A. Manning. 1984. *Vibrio cholerae* bacteriophage CP-T1: characterization of bacteriophage DNA and restriction analysis. *J. Virol.* **51**:163–169.
 19. Hazelbauer, G. L. 1975. Role of the receptor for bacteriophage lambda in the functioning of the maltose chemoreceptor of *Escherichia coli*. *J. Bacteriol.* **124**:119–126.
 20. Heidelberg, J. F., J. A. Eisen, W. C. Nelson, R. A. Clayton, M. L. Gwinn, R. J. Dodson, D. H. Haft, E. K. Hickey, J. D. Peterson, L. Umayam, S. R. Gill, K. E. Nelson, T. D. Read, H. Tettelin, D. Richardson, M. D. Ermolaeva, J. Vamathevan, S. Bass, H. Qin, I. Dragoi, P. Sellers, L. McDonald, T. Utterback, R. D. Fleischmann, W. C. Nierman, O. White, S. L. Salzberg, H. O. Smith, R. R. Colwell, J. J. Mekalanos, J. C. Venter, and C. M. Fraser. 2000. DNA sequence of both chromosomes of the cholera pathogen *Vibrio cholerae*. *Nature* **406**:477–483.
 21. Holmgren, A., G. B. Kallis, and B. Nordstrom. 1981. A mutant thioredoxin from *Escherichia coli* tsnC 7007 that is nonfunctional as subunit of phage T7 DNA polymerase. *J. Biol. Chem.* **256**:3118–3124.
 22. Hood, A. M. 1953. Phage typing of *Staphylococcus aureus*. *J. Hyg. (London)* **51**:1–15.
 23. Inoue, T., S. Matsuzaki, and S. Tanaka. 1995. A 26-kDa outer membrane protein, OmpK, common to *Vibrio* species is the receptor for a broad-host-range vibriophage, KVP40. *FEMS Microbiol. Lett.* **125**:101–105.
 24. Jouravleva, E. A., G. A. McDonald, J. W. Marsh, R. K. Taylor, M. Boesman-Finkelstein, and R. A. Finkelstein. 1998. The *Vibrio cholerae* mannose-sensitive hemagglutinin is the receptor for a filamentous bacteriophage from *V. cholerae* O139. *Infect. Immun.* **66**:2535–2539.
 25. Kapfhammer, D., J. Blass, S. Evers, and J. Reidl. 2002. *Vibrio cholerae* phage K139: complete genome sequence and comparative genomics of related phages. *J. Bacteriol.* **184**:6592–6601.
 26. Kato, H., H. Fujisawa, and T. Minagawa. 1985. Purification and characterization of gene 17 product of bacteriophage T3. *Virology* **146**:22–26.
 27. Kovalyova, I. V., and A. M. Kropinski. 2003. The complete genomic sequence of lytic bacteriophage gh-1 infecting *Pseudomonas putida*—evidence for close relationship to the T7 group. *Virology* **311**:305–315.
 28. Lalko, J., and A. Gunnell. 1967. A new Vi-phage-type of *Salmonella typhi* and preparation of the typing phage. *Bull. W. H. O.* **36**:227–230.
 29. Liu, G., M. Yan, G. Qi, Y. Liu, S. Gao, and B. Kan. 2004. Chloramphenicol resistance marker exchange of lysogenic phage CTXΦ of *Vibrio cholerae* and its induction. *Zhonghua Wei Sheng Wu Xue He Mian Yi Xue Za Zhi* **24**:253–257.
 30. Makhov, A. M., B. L. Trus, J. F. Conway, M. N. Simon, T. G. Zurabishvili, V. V. Mesyanzhinov, and A. C. Steven. 1993. The short tail-fiber of bacteriophage T4: molecular structure and a mechanism for its conformational transition. *Virology* **194**:117–127.
 31. Mark, D. F., and C. C. Richardson. 1976. *Escherichia coli* thioredoxin: a subunit of bacteriophage T7 DNA polymerase. *Proc. Natl. Acad. Sci. USA* **73**:780–784.
 32. Merino, S., S. Camprubi, and J. M. Tomas. 1990. Isolation and characterization of bacteriophage PM3 from *Aeromonas hydrophila* the bacterial receptor for which is the monopolar flagellum. *FEMS Microbiol. Lett.* **57**:277–282.
 33. Miller, E. S., E. Kutter, G. Mosig, F. Arisaka, T. Kunisawa, and W. Ruger. 2003. Bacteriophage T4 genome. *Microbiol. Mol. Biol. Rev.* **67**:86–156.
 34. Nesper, J., A. Kraiss, S. Schild, J. Blass, K. E. Klose, J. Bockemuhl, and J. Reidl. 2002. Comparative and genetic analyses of the putative *Vibrio cholerae* lipopolysaccharide core oligosaccharide biosynthesis (*wav*) gene cluster. *Infect. Immun.* **70**:2419–2433.
 35. Pajunen, M., S. Kiljunen, and M. Skurnik. 2000. Bacteriophage ΦYcO3-12, specific for *Yersinia enterocolitica* serotype O:3, is related to coliphages T3 and T7. *J. Bacteriol.* **182**:5114–5120.
 36. Pajunen, M. I., M. R. Elizondo, M. Skurnik, J. Kieleczawa, and I. J. Molineux. 2002. Complete nucleotide sequence and likely recombinatorial origin of bacteriophage T3. *J. Mol. Biol.* **319**:1115–1132.
 37. Philippe, N., J. P. Alcaraz, E. Coursange, J. Geiselmann, and D. Schneider. 2004. Improvement of pCVD442, a suicide plasmid for gene allele exchange in bacteria. *Plasmid* **51**:246–255.
 38. Qimron, U., B. Marintcheva, S. Tabor, and C. C. Richardson. 2006. Genomewide screens for *Escherichia coli* genes affecting growth of T7 bacteriophage. *Proc. Natl. Acad. Sci. USA* **103**:19039–19044.
 39. Reidl, J., and J. J. Mekalanos. 1995. Characterization of *Vibrio cholerae* bacteriophage K139 and use of a novel mini-transposon to identify a phage-encoded virulence factor. *Mol. Microbiol.* **18**:685–701.
 40. Sambrook, J., and D. W. Russell. 2001. Molecular cloning: a laboratory manual, 3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
 41. Schild, S., A. K. Lamprecht, and J. Reidl. 2005. Molecular and functional characterization of O antigen transfer in *Vibrio cholerae*. *J. Biol. Chem.* **280**:25936–25947.
 42. Scholl, D., S. Adhya, and C. Merrill. 2005. *Escherichia coli* K1's capsule is a barrier to bacteriophage T7. *Appl. Environ. Microbiol.* **71**:4872–4874.
 43. Scholl, D., and C. Merrill. 2005. The genome of bacteriophage K1F, a T7-like phage that has acquired the ability to replicate on K1 strains of *Escherichia coli*. *J. Bacteriol.* **187**:8499–8503.
 44. Simon, R., U. Priefer, and A. Puhler. 1983. A broad host range mobilization system for in vivo genetic engineering: transposon mutagenesis in gram negative bacteria. *Nat. Biotechnol.* **1**:784–791.
 45. Steven, A. C., B. L. Trus, J. V. Maizel, M. Unser, D. A. Parry, J. S. Wall, J. F. Hainfeld, and F. W. Studier. 1988. Molecular substructure of a viral receptor-recognition protein. The gp17 tail-fiber of bacteriophage T7. *J. Mol. Biol.* **200**:351–365.
 46. Tabor, S., H. E. Huber, and C. C. Richardson. 1987. *Escherichia coli* thioredoxin confers processivity on the DNA polymerase activity of the gene 5 protein of bacteriophage T7. *J. Biol. Chem.* **262**:16212–16223.
 47. Tetart, F., C. Desplats, and H. M. Krisch. 1998. Genome plasticity in the distal tail fiber locus of the T-even bacteriophage: recombination between conserved motifs swaps adhesin specificity. *J. Mol. Biol.* **282**:543–556.
 48. Thomassen, E., G. Gielen, M. Schutz, G. Schoehn, J. P. Abrahams, S. Miller, and M. J. van Raaij. 2003. The structure of the receptor-binding domain of the bacteriophage T4 short tail fibre reveals a knitted trimeric metal-binding fold. *J. Mol. Biol.* **331**:361–373.
 49. Waldor, M. K., and J. J. Mekalanos. 1996. Lysogenic conversion by a filamentous phage encoding cholera toxin. *Science* **272**:1910–1914.
 50. Werts, C., V. Michel, M. Hofnung, and A. Charbit. 1994. Adsorption of bacteriophage lambda on the LamB protein of *Escherichia coli* K-12: point mutations in gene J of lambda responsible for extended host range. *J. Bacteriol.* **176**:941–947.
 51. Yokota, S., T. Hayashi, and H. Matsumoto. 1994. Identification of the lipopolysaccharide core region as the receptor site for a cytotoxin-converting phage, ΦCTX, of *Pseudomonas aeruginosa*. *J. Bacteriol.* **176**:5262–5269.
 52. Yu, S. L., K. L. Ko, C. S. Chen, Y. C. Chang, and W. J. Syu. 2000. Characterization of the distal tail fiber locus and determination of the receptor for phage AR1, which specifically infects *Escherichia coli* O157:H7. *J. Bacteriol.* **182**:5962–5968.