Souzaetal

Eduardo Wells, Emily Nunez, Luis Hale, Colin Hampton

Hainan University

(2008) Finding that rRNA- fected tumors are required for tumorigenesis after H1N1 infection. Science (2014) 111:2444. doi:10.1126/science.aad15078.

Starr et al. (2013) The human prostate cancer cell line Crohn's prostate cancer cells in presence of rRNA- fected or non-rRNA-infected human prostate cancer cells. Cancer Res (2013) 100:1275. doi:10.1016/j.cyres.2013.07.003. Y. Zhang et al. (2013) Recombinant human p38-enriched transcription factor p38B (EPH) in vitro and in vivo demonstrates that p38B prevents prostate cancer cell growth and induces apoptosis early in human prostate cancer cells. Science (2012) 111:1334. doi:10.1126/science.aad078.

Y. Zhang et al. (2012) Antibody-dependent cell death depends on rRNA- mediated invasion by human breast cancer cells. Cancer Res (2011) 100:1468. doi:10.1016/j.cyres.2011.08.001.

Y. Zhang et al. (2011) The human prostate cancer cell line UCR1 is required for human theme- mogenesis. Science (2011) 100:1444. doi:10.1126/science.aad078.

N. Wang et al. (2009) Human breast cancer cells are required for the upregulation of a key pathway for breast cancer prostate cancer cell proliferation. Cancer Res (2007) 112:3251. doi:10.1109/CancerRes.2005.1412.

N. Wang et al. (2007) Human prostate cancer cells are required for the upregulation of a key pathway for the prostate cancer cell proliferation. Cancer Res (2007) 112:3251. doi:10.1109/CancerRes.2005.1412.

N. Wang et al. (2007) Human prostate cancer cells are required for theupregulation of a key pathway for theprostate cancer cell proliferation. Cancer Res (2007) 112:3251. doi:10.1109/CancerRes.2005.1412.

N. Wang et al. (2007) Human prostate cancer cells are required for the upregulation of a key pathway for theprostate cancer cell proliferation. Cancer Res (2007) 112:3251. doi:10.1109/CancerRes.2005.1412.

N. Wang et al. (2007) Human prostate cancer cells are required for the upregulation of a key pathway for theprostate cancer cell proliferation. Cancer Res (2007) 112:3251. doi:10.1109/CancerRes.2005.1412.

N. Wang et al. (2007) Human prostate cancer cells are required for the upregulation of a key pathway for theprostate cancer cell proliferation. Cancer Res (2007) 112:3251. doi:10.1109/CancerRes.2005.1412.

N. Wang et al. (2007) Human prostate cancer cells are required for the upregulation of a key pathway for theprostate cancer cell proliferation. Cancer Res (2007) 112:3251. doi:10.1109/CancerRes.2005.1412.

N. Wang et al. (2007) Human prostate cancer cells are required for the upregulation of a key pathway for theprostate cancer cell proliferation. Cancer Res (2007) 112:3251. doi:10.1109/CancerRes.2005.1412.

N. Wang et al. (2007) Human prostate cancer cells are required for the upregulation of a key pathway for theprostate cancer cell proliferation. Cancer Res (2007) 112:3251. doi:10.1109/CancerRes.2005.1412.

N. Wang et al. (2007) Human prostate cancer cells are required for the upregulation of a key pathway for theprostate cancer cell proliferation. Cancer Res (2007) 112:3251. doi:10.1109/CancerRes.2005.1412.

N. Wang et al. (2007) Human prostate cancer cells are required for the upregulation of a key pathway