

Nasalbindingsitesofthecaspase3geneCBA1are

Catherine Davis, Robert Johnson, Sherry Munoz, Andrew
Bender, Terri Swanson, Jody Fox, Debra Jones, Marilyn
West, Amanda Jones

University Hospital Regensburg

present in *N. sumatrana* (Apollonia sativa L. perlane A8, Edwardsieville, USC) and are especially related to the CBA1-CBA3 fusion protein (2). The CBA1-CBA3 fusion protein of the CBA1-CBA3 protein family is the most studied fusion protein in this plant pathogen. The CBA1-CBA3 fusion protein of the CBA1-CBA3 protein family is also the subject of a recent study [39]. It is generally recognized that high-affinity CBA2 and CBA3 protein complexes have been detected in *C. albicans* [40]. In this study, we counted 5-fold increased activation of CBA2 and CBA3 protein complexes and analyzed the effect of CBA2 and CBA3 on *N. sumatrana* cell migration. A detailed analysis of the effects of CBA2 and CBA3 was performed with the YAP assay. CBA2 and CBA3 significantly inhibited the transfection of the CBA1-CBA3 fusion protein of the CBA1-CBA3 protein family together with the inhibition of invasion of the *N. sumatrana* ($P < 0.05$ by Student's t-test). The transfection was conducted with lentiviral vectors with or without pNA (Sigma) or pNA (Sigma) that contained 25 ng of the CBA1-CBA3 protein family in their promoter regions. The result showed that the transfection of CBA1-CBA3 proteins with 25 ng of the CBA1-CBA3 protein family was sufficient to induce the transfection of CBA1-CBA3 proteins with 25 ng of CBA1-CBA3 protein family in the caspase-3 locus. CBA1-CBA3 fusion protein is a crucial component in the migration of *N. sumatrana* (Papienziata). To address the role of the CBA1-CBA3 fusion protein in the migration of *N. sumatrana*, we performed a crude morphological analysis of the CBA1-CBA3 protein family of the CBA1-CBA3 fusion protein using the YAP assay. We found that the CBA1-CBA3 fusion protein of the CBA1-CBA3 protein family had a significantly reduced expression in migration of the CBA1-CBA3 fusion protein compared to the CBA1-CBA3 protein family alone. The CBA1-CBA3 fusion protein of the CBA1-CBA3 protein family was highly translocated between the tail and the base of the Figure 1. CBA1-CBA3 fusion protein is involved in the recruitment and migration of *N. sumatrana* (Papienziata) migratory glands in the eye. (A) CBA1-CBA3 fusion protein expression in the *N. sumatrana* (Papienziata) migrating rodent glands of the eye (A1). (B) CBA1-CBA3 fusion protein expression in the *N. sumatrana* (Papienziata) migratory glands of the eye (A2). (B1) (D) CBA1-CBA3 fusion protein expression in the *N. sumatrana* (Papienziata) migrating rodent glands of the eye (A3). (B2) (E) CBA1-CBA3 fusion protein expression in the *N. sumatrana* (Papienziata) migrating rodent glands of the eye (A4). (E1) CBA1-CBA3 fusion protein expression in the *N. sumatrana* (Papienziata) migrating rodent glands of the eye. (E2) CBA1-CBA3 fusion protein expression in the *N. sumatrana* (Papienziata) migrating rodent glands of the eye. (E3) (E4) CBA1-CBA3 fusion protein expression in the *N. sumatrana* (Papienziata) migrating rodent glands of the eye. (E5) CBA1-CBA3 fusion protein expression in the *N. sumatrana* (Papienziata) migrating rodent glands of the eye. CBA1-CBA3 protein expression in the *N. sumatrana* (Papienziata) migrating rodent glands of the eye. CBA1-