

As reported previously²⁵ the tumor necrosis factor beta inducible

Justin Davis, Erin Ho, Amanda Rubio, Ryan Rich,
Sandra Wood, Nicole McDonald, Briana Owens, Amber
Cook, Heather Henderson, Leslie Shields

University of Cincinnati

FIG. 1. Expression of p-ERK and p38 in Rps7- and MHC-labeled Rps7-E2 cells. Rps7 cells were treated with 5 mM DMSO, 5 mM MgCl₂, 10 mM Na₂VO₄, 2 mM DTT, 0.1 mM m-FGF, and 10 mM Na₂VO₄ and were treated with 0.1 M of DMSO for 1 h. The cells were washed twice with PBS and then resuspended in RPMI 1640. The cells were then incubated with 10 mM DMSO for 1 h. After washing, cells were collected, and the cells were collected and analyzed for apoptosis. FIG. 2. Expression of p-ERK and p38 in Rps7- and MHC-labeled Rps7-E2 cells. Rps7 cells were treated with 5 mM DMSO, 5 mM MgCl₂, 10 mM Na₂VO₄, 2 mM DTT, 0.1 mM m-FGF, and 10 mM Na₂VO₄ and were treated with 0.1 M of DMSO for 1 h. The cells were washed twice with PBS and then resuspended in RPMI 1640. The cells were incubated with 10 mM DMSO for 1 h. The cells were detected by a fluorescence-microscopy. (A) Growth inhibition of Rps7 cells was measured by 3S-ADP-conjugated secondary antibody (Dako IV, Japan) using a Zeiss Axiofermeine (Zeiss, Co., Japan). The cells were treated with 25 mM DMSO for 1 h, and the cells were washed twice with PBS. The cells were then incubated with 10 mM DMSO for 1 h. The cells were collected, and analyzed for apoptosis. (B) Growth inhibition of Rps7 cells was measured by 3S-ADP-conjugated secondary antibody (Dako IV, Japan) using a Zeiss Axiofermeine (Zeiss, Co., Japan). The cells were treated with 25 mM DMSO for 1 h, and the cells were washed twice with PBS. The cells were then incubated with 10 mM DMSO for 1 h. The cells were collected, and analyzed for apoptosis. FIG. 3. Mapping of apoptosis on Rps7- and MHC-labeled Rps7-E2 cells. Cells were treated with 5 mM DMSO, 5 mM MgCl₂, 10 mM Na₂VO₄, 2 mM DTT, 0.1 mM m-FGF, and 10 mM Na₂VO₄ and were treated with 0.1 M of DMSO for 1 h. The cells were washed twice with PBS, and the cells were incubated with 10 mM DMSO for 1 h. (A) Mapping of apoptosis on Rps7- and MHC-labeled Rps7-E2 cells. The cells were treated with 5 mM DMSO, 5 mM MgCl₂, 10 mM Na₂VO₄, 2 mM DTT, 0.1 mM m-FGF, and 10 mM Na₂VO₄ and were treated with 0.1 M of DMSO for 1 h. The cells were washed twice with PBS, and the cells were incubated with 10 mM DMSO for 1 h. (B) Mapping of apoptosis on Rps7-E2 cells. The cells were treated with 5 mM DMSO, 5 mM MgCl₂, 10 mM Na₂VO₄, 2 mM DTT, 0.1 mM m-FGF, and 10 mM Na₂VO₄ and were treated with 0.1 M of DMSO for 1 h. The cells were washed twice with PBS, and the cells were incubated with 10 mM DMSO for 1 h. (C) Mapping of apoptosis on Rps7- and MHC-labeled Rps7-E2 cells. The cells were treated with 5 mM DMSO, 5 mM MgCl₂, 10 mM Na₂VO₄, 2 mM DTT, 0.1 mM m-FGF, and 10 mM Na₂VO₄ and were treated with 0.1 M of DMSO for 1 h. The cells were washed