

# INS\_IEP: A Matlab package for fitting Inelastic Neutron Scattering data.

Alban Bloor Riley<sup>1</sup>, Marcus Webb<sup>1</sup>, and Michael L. Baker<sup>2</sup>

<sup>1</sup> The University of Manchester, Department of Mathematics <sup>2</sup> The University of Manchester, Department of Chemistry ¶ Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

## Software

- Review
- Repository
- Archive

Editor: [Open Journals](#)

Reviewers:

- @openjournals

Submitted: 01 January 1970

Published: unpublished

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#))

## Summary

Inelastic neutron scattering (INS) is a spectroscopic technique used to measure the magnetic excitations in materials with interacting electron spins. Fitting the experimental data to a spin Hamiltonian model can be formulated as an inverse eigenvalue problem (IEP). INS\_IEP is a MATLAB package that uses deflated numerical optimisation techniques to find multiple solutions to this problem. The package requires and is fully compatible with easyspin (Stoll & Schweiger, 2006), a package for solving similar problems in electron paramagnetic resonance.

## Statement of need

Neutrons are an excellent bulk probe of material properties since they carry no charge and therefore penetrate deeply into matter. Neutrons also carry a quantum spin of a half, making them a sensitive probe of magnetism (?). Reactors and spallation sources with dedicated high-flux neutron sources serve the international research community with neutron scattering experiment capabilities for material research. One technique at neutron facilities is the measurement of inelastic neutron scattering (INS) to study magnetism. In an INS experiment, a sample under investigation is irradiated with a beam of neutrons and the scattered neutron energy and momentum transfer are detected. For samples composed of finite-size clusters of magnetic moment-carrying atoms, such as single ions or molecular-based magnets, the detected neutron energy transfer gives direct access to the quantum spin excitations (M. Baker & Mutka, 2012; Furrer & Waldmann, 2013; "Spectroscopy Methods for Molecular Nanomagnets," 2014). The energy of such excitations relates to the energy difference between eigenvalues of the Hamiltonian matrix that describes the quantum spin dynamics of the compound in question. Single-ion and molecular-based magnets are studied as prototype components (quantum bits, sensors) for quantum technologies. INS can therefore provide crucial information concerning the precise quantum properties of such systems. However, to relate the INS experimental results to the Hamiltonian that describes quantum spin dynamics requires parameterisation of matrix elements such that a set of eigenvalues and eigenstates matching the experiment are determined. This situation is known as the inverse eigenvalue problem. To date, this problem is addressed in an iterative process where parameters of the Hamiltonian are varied and the resultant eigenvalues compared with experiment. INS\_IEP presents an elegant solution to solving this problem, alleviating the necessity of iterative Hamiltonian matrix diagonalisation, providing a more robust method to reliably extract an accurate spin Hamiltonian model from INS experimental data.

## The Spin Hamiltonian

The Spin Hamiltonian,  $H$ , is an approximation of the Hamiltonian that uses spin coordinates instead of orbital coordinates, it is widely used to model data arising from many spectroscopy techniques (?). It can be modeled as a linear combination of interaction terms, we will focus on the zero field interaction,  $H_{ZFI}$ , and the electron-electron interaction,  $H_{EEI}$ :

$$H = H_{ZFI} + H_{EEI}$$

Both of these terms can themselves be modelled as the linear sum of other basis matrices. The zero field interaction can be written as:

$$H_{ZFI} = \sum_{-k \leq q \leq k} B_k^q O_k^q$$

where the  $O_k^q$  are Stevens Operators (Rudowicz & Chung, 2004), and  $B_k^q$  the associated parameter. When there are multiple spin centres it is necessary to take kronecker products of the operator with identity matrices of the appropriate for each other spin centre.

When there are multiple spin centres it is also necessary to include an electron-electron interaction term,  $H_{\{EEI\}}$ . This term will be the sum of interaction terms between each pair of spin centres:

$$H_{EEI} = - \sum_{i \neq j} J_{ij} S_i \cdot S_j$$

where  $S_i$  is the vector of spin operators  $S_i = [S_x, S_y, S_z]$  for the  $i$ -th spin centre, and  $J_{ij}$  is the unknown parameter. Note that in the isotropic case  $J$  can be thought of as a scalar value, but in the anisotropic case will be a matrix where the off diagonals are skew symmetric. While the summation is in theory over all spin centre combinations, in practice many of these contributions will be negligible - often only the nearest neighbour interactions are modeled.

## Mathematical Background

### Inverse Eigenvalue Problem

The INS experiments provide eigenvalues of the Spin Hamiltonian matrix of the sample, the task of calculating the matrix from the eigenvalues is an inverse eigenvalue problem:

Let  $A(x)$  be the affine family of matrices,

$$A(x) = A_0 + \sum_{i=1}^{\ell} x_i A_i,$$

where  $x \in \mathbb{R}^{\ell}$  and  $A_0, \dots, A_{\ell} \in \mathbb{R}^{n \times n}$  are linearly independent symmetric matrices, and denote the ordered eigenvalues of  $A(x)$  as  $\lambda_1(x) \leq \dots \leq \lambda_n(x)$ . Then the least squares inverse eigenvalue problem is to find the parameters  $x \in \mathbb{R}^{\ell}$  that minimises

$$F(x) = \frac{1}{2} \|r(x)\|_2^2 = \frac{1}{2} \sum_{i=1}^m (\lambda_i(x) - \lambda_i^*)^2$$

where  $\lambda_1^* \leq \dots \leq \lambda_m^*$  are the experimental eigenvalues. In the case of INS fitting the  $A_i$  basis matrices will be a combination of Stevens operators and electron-electron exchange terms.

66 The IEP described above is formulated as an least squares problem, this is due to the fact that  
67 the number of eigenvalues that can be probed by INS experiments is often a small subset of  
68 the full spectrum

69 As far as we are aware this is the first time that the fitting of INS data has been explicitly  
70 formulated as an IEP. The advantage of this formulation is that there are explicit formulas for  
71 the derivatives of  $r(x)$ , the first derivative (Jacobian) is:

$$J_r(x) = \begin{pmatrix} q_1(x)^T A_1 q_1(x) & \dots & q_1(x)^T A_\ell q_1(x) \\ \vdots & \ddots & \vdots \\ q_m(x)^T A_1 q_m(x) & \dots & q_m(x)^T A_\ell q_m(x) \end{pmatrix}.$$

72 And the second derivative (Hessian) is:

$$(H_r)_{ij} = 2 \sum_{k=1}^m (\lambda_k - \lambda_k^*) \sum_{\substack{t=1 \\ \lambda_t \neq \lambda_k}}^m \frac{(q_t^T A_i q_k)(q_t^T A_j q_k)}{\lambda_k - \lambda_t}.$$

73 Access to analytical forms of the derivatives means it is not necessary to use the finite difference  
74 approximation that other approaches use, making the three optimisation methods that INS\_IEP  
75 uses faster and more accurate.

## 76 Methods

77 All of the methods used are iterative schemes of the form  $x^{k+1} = x^k + p^k$  where the step  $p^k$   
78 uniquely defines each algorithm:


- 79 ■ Newton's method:  $p^k = (J_r^T J_r + H_r r) - 1 J_r^T r$  (Nocedal & Wright, 2006)
- 80 ■ Gauss-Newton method:  $p^k = (J_r^T J_r)^{-1} J_r^T r$  (Bloor Riley et al., 2025)
- 81 ■ Lift and Projection Method:  $p^k = B^{-1} J_r^T r$  (?)

82 Where the matrix  $B$  is the Gram matrix formed from the frobenius inner products of the  
83 basis matrices:  $B_{ij} = \langle A_i, A_j \rangle_F$ . The Lift and Projection method is a Riemannian Gradient  
84 descent method (?), inspired by the Lift and Projection method (Chen & Chu, 1996), specifically  
85 designed for solving IEPs. In (?) it is proven that the method is a strictly descending algorithm,  
86 that is it reduces the value of the objective function every step. Both the deflated Gauss-Newton  
87 method and the Riemannian Gradient descent Lift and Projection method are new methods  
88 designed for this package.


## 89 Deflation

90 The number of eigenvalues that can be probed via INS experiments varies depending on the  
91 equipment and sample in question, meaning that the fitting problem is often under (or even  
92 over) determined. The IEP is also highly nonlinear and due to the experimental nature of  
93 the data there is no guarantee that the problem is not ill-posed. One consequence of this is  
94 that the solution space may be very 'bumpy', that is there may exist many local minimisers to  
95 the problem. For example in Figure 1, there are clearly 4 distinct solutions (for more details  
96 see Example 1 and the file Example1\_Mn12.m in the examples folder). We seek to solve  
97 the problem of multiple local minima by the use of Deflation, a numerical technique used to  
98 find multiple solutions to systems of equations (P. E. Farrell et al., 2015; Patrick E. Farrell  
99 et al., 2020). Fortunately it is cheap to apply deflation for the above methods, it is simply a  
100 change to the length of the step - notably this means that the direction of each step does not  
101 change. It is proven in (Bloor Riley et al., 2025) that the deflated methods will not converge  
102 to deflated points. The usual requirements still apply to the convergence of the new methods -  
103 that the initial guess is close enough to the new minimum, and that the Jacobian is full rank  
104 in a neighbourhood around that minimum. The rate of convergence of the deflated methods is

also more complicated, although the number of iterations required to converge can go up with the number of deflations this is not a strict correlation, as can be seen in Figure 2.



**Figure 1:** Contour plot of how  $F$  varies with the two parameters  $B_2^2$  and  $B_4^4$  for the molecule Mn\_12 as described in Example 1.



**Figure 2:** Rate of converge for each deflation.

## Examples

### Example 1 - Mn12

The first example we will look at is Manganese-12-acetate. This is a well know example in the INS and magnetism community, as one of the first molecules that behaves like a nano-sized magnet with a molecular magnetic coercivity as well as its role in the research of quantum tunnelling of magnetisation (Friedman et al., 1996; Sessoli et al., 1993).

The Spin Hamiltonian of this system, using the giant spin approximation, can be represented as a  $21 \times 21$  matrix modelled using 4 Stevens operators (Bircher et al., 2004):

$$H = B_2^0 O_2^0 + B_4^0 O_4^0 + B_2^2 O_2^2 + B_4^4 O_4^4 \in \mathbb{R}^{21 \times 21}$$

We utilise the same spin system syntax as easyspin, so to set up the problem we first set up the model, along with initial guesses for the parameters:

```
%One spin centre (because giant spin approximation)
Sys0.S=10;
%Four Stevens operators
Sys0.B2 = [-100,0,-1000,0,0];
Sys0.B4 = [-1,0,0,0,-1,0,0,0,0];
```

Then we input the experimental eigenvalues, these will normally be translated such that the smallest eigenvalue is zero, and define which parameters to fit. Note that all values given must be in Hertz, so it may be useful to use conversions.

```
rcm = 29979.2458; meV = rcm*8.065; %Conversions values
%Input calculated eigenvalues:
Exp.ev = [0,0,1.24,1.24,2.3,2.3,3.18,3.18,3.91,3.91,4.5,4.5,4.97,4.97,5.32,5.32,5.54,5.54];
%Vary all non zero parameters (no Fixed parameters):
Vary = Sys0;
```

Then all that is required is to call INS\_IEP with these three inputs:

```
SysOut = INS_IEP(Sys0,Vary,Exp);
```

If we wish to find all four solutions as shown in figure Figure 1 then we use the additional option:

```
Opt.NDeflations = 4;
SysOut = INS_IEP(Sys0,Vary,Exp,Opt);
```

In this case SysOut will be an array of four spin structures each containing a distinct solution.

### Example 2 - Chromium(iii) Horseshoes

The second example concerns antiferro-magnetically coupled chromium(III) chains six atoms long (M. L. Baker et al., 2011), although different length chains are of interest and can also be modelled. Because there are multiple spin centres an electron-electron interaction term is required. The spin hamiltonian is a  $4096 \times 4096$  matrix composed of two Stevens operators and one interaction term:

```
Sys0.S = [1.5 1.5 1.5 1.5 1.5 1.5];
Sys0.B2 = [1 0 -1 0 0 0;
           1 0 -1 0 0 0;
           1 0 -1 0 0 0;
           1 0 -1 0 0 0;
           1 0 -1 0 0 0];
```

```
Sys0.J = [100,0,0,0,0,100,0,0,0,100,0,0,100,0,100];
Vary = Sys0;
Exp.ev = [0,0.355,0.457,0.497,1.576,1.577,1.592,1.629,1.632,2.97,2.98,3.002,3.004,
3.01,3.038,3.821,3.824,3.827,3.837,3.856,3.879,3.888,3.895,3.903];
```

Note that only 24 eigenvalues were found experimentally, so this will form a partial eigenvalue problem. To find the solution system is as simple as:

```
SysOut = INS_IEP(Sys0,Vary,Exp);
```

It is possible to find multiple minimising systems even if they do not make any sense physically, however due to the scaling of the problem a change in the default deflation parameters is necessary:

```
Opt = struct('NDeflations',4,'Sigma',1e-7);
SysOut= INS_IEP(Sys0,Vary,Exp,Opt);
```

The output contains four different spin systems that all have the same eigenvalues as input, one is the original solution up to a change of sign of the  $B_2^2$  parameter, and all have the same exchange term.

## Acknowledgements

ABR thanks the University of Manchester for a Dean's Doctoral Scholarship. MW thanks the Polish National Science Centre (SONATA-BIS-9), project no. 2019/34/E/ST1/00390, for the funding that supported some of this research.

## References

- Baker, M. L., Bianchi, A., Carretta, S., Collison, D., Docherty, R. J., McInnes, E. J. L., McRobbie, A., Muryn, C. A., Mutka, H., Piligkos, S., Rancan, M., Santini, P., Timco, G. A., Tregenna-Piggott, P. L. W., Tuna, F., Güdel, H. U., & Winpenny, R. E. P. (2011). Varying spin state composition by the choice of capping ligand in a family of molecular chains: Detailed analysis of magnetic properties of chromium(iii) horseshoes. *Dalton Transactions*, 40(12), 2725. <https://doi.org/10.1039/c0dt01243b>
- Baker, M., & Mutka, H. (2012). Neutron spectroscopy of molecular nanomagnets. *The European Physical Journal Special Topics*, 213, 53–68. <https://doi.org/10.1140/epjst/e2012-01663-6>
- Bircher, R., Chaboussant, G., Sieber, A., Güdel, H. U., & Mutka, H. (2004). Transverse magnetic anisotropy in Mn 12 acetate: Direct determination by inelastic neutron scattering. *Phys. Rev. B*, 70(21), 212413. <https://doi.org/10.1103/PhysRevB.70.212413>
- Bloor Riley, A., Webb, M., & Baker, M. L. (2025). *Deflation Techniques for Finding Multiple Local Minima of a Nonlinear Least Squares Problem*. arXiv. <https://doi.org/10.48550/arXiv.2409.14438>
- Chen, X., & Chu, M. T. (1996). On the least squares solution of inverse eigenvalue problems. *SIAM Journal on Numerical Analysis*, 33, 2417–2430. <https://doi.org/10.1137/S0036142994264742>
- Farrell, P. E., Á. Birkisson, & Funke, S. W. (2015). Deflation Techniques for Finding Distinct Solutions of Nonlinear Partial Differential Equations. *SIAM J. Sci. Comp.*, 37(4), A2026–A2045. <https://doi.org/10.1137/140984798>
- Farrell, Patrick E., Croci, M., & Surowiec, T. M. (2020). Deflation for semismooth equations. *Optim. Method. Softw.*, 35(6), 1248–1271.

- 165 Friedman, J. R., Sarachik, M. P., Tejada, J., & Ziolo, R. (1996). Macroscopic Measurement  
166 of Resonant Magnetization Tunneling in High-Spin Molecules. *Phys. Rev. Lett.*, 76(20),  
167 3830–3833. <https://doi.org/10.1103/physrevlett.76.3830>
- 168 Furrer, A., & Waldmann, O. (2013). Magnetic cluster excitations. *Reviews of Modern Physics*,  
169 85, 367–420. <https://doi.org/10.1103/RevModPhys.85.367>
- 170 Nocedal, J., & Wright, S. J. (2006). *Numerical optimization* (2nd ed). Springer. ISBN: 978-0-  
171 387-30303-1
- 172 Rudowicz, C., & Chung, C. Y. (2004). The generalization of the extended Stevens operators  
173 to higher ranks and spins, and a systematic review of the tables of the tensor operators  
174 and their matrix elements. *Journal of Physics: Condensed Matter*, 16(32), 5825–5847.  
175 <https://doi.org/10.1088/0953-8984/16/32/018>
- 176 Sessoli, R., Gatteschi, D., Caneschi, A., & Novak, M. A. (1993). Magnetic bistability in a  
177 metal-ion cluster. *Nature*, 365(6442), 141–143. <https://doi.org/10.1038/365141a0>
- 178 Spectroscopy Methods for Molecular Nanomagnets. (2014). In *Structure and Bonding* (pp.  
179 231–291). Springer Berlin Heidelberg. [https://doi.org/10.1007/430\\_2014\\_155](https://doi.org/10.1007/430_2014_155)
- 180 Stoll, S., & Schweiger, A. (2006). EasySpin, a comprehensive software package for spectral  
181 simulation and analysis in EPR. *Journal of Magnetic Resonance*, 178, 42–55. <https://doi.org/10.1016/j.jmr.2005.08.013>  
182

DRAFT