

Total No. of Questions—8]

[Total No. of Printed Pages—6

Seat No.	
-------------	--

[5352]-166

S.E. (Computer Engineering/Information Technology)

(II Sem.) EXAMINATION, 2018

ENGINEERING MATHEMATICS—III

(2012 PATTERN)

Time : Two Hours

Maximum Marks : 50

N.B. :— (i) Attempt Q. No. 1 or Q. No. 2, Q. No. 3 or Q. No. 4,
Q. No. 5 or Q. No. 6, Q. No. 7 or Q. No. 8.

(ii) Neat diagrams must be drawn wherever necessary.

(iii) Figures to the right indicate full marks.

(iv) Use of electronic non-programmable calculator is allowed.

(v) Assume suitable data, if necessary.

1. (a) Solve (any two) : [8]

(i) $(D^3 - 7D - 6)y = e^{2x}(1 + x)$

(ii) $(D^2 + 4)y = \frac{1}{1 + \cos 2x}$ by method of variation of parameters

(iii) $x^3 \frac{d^2 y}{dx^2} + 3x^2 \frac{dy}{dx} + xy = \sin(\log x)$

(b) Find Fourier sine transform of : [4]

$$f(x) = \frac{e^{-ax}}{x}, \quad x > 0.$$

P.T.O.

Or

2. (a) A circuit consists of inductance L and condenser of capacity C in series. An emf $E \sin pt$ is applied at $t = 0$, the initial change and current being zero. If $p^2 = \frac{1}{LC}$, find current in the circuit at time t . [4]

- (b) Find inverse z -transform of (any one) : [4]

(i) $F(z) = \frac{z^3}{(z-1)(z-2)^2}, \quad |z| > 2$

(ii) $F(z) = \frac{10z}{(z-1)(z-2)}$ by inversion integral method.

- (c) Solve difference equation : [4]

$$f(k+2) - 3f(k+1) + 2f(k), \quad f(0) = 0, \quad f(1) = 1.$$

3. (a) The first four moments of a distribution about the value 4 are 1, 4, 10 and 45. Obtain the first four central moments, mean, standard deviation and coefficient of skewness and kurtosis. [4]

- (b) A sample of 100 dry battery cells tested to find the length of life produced the following results; $\bar{X} = 12$ hours, $\sigma = 3$ hours. Assuming the data to be normally distributed, what percentage of battery cells are expected to have life between 10 and 14 hours ? [Given : For $z = 0.67$, area = 0.2487] [4]

- (c) If the directional derivative of : [4]

$$\phi = a(x + y) + b(y + z) + c(x + z)$$

has maximum value 12 in the direction of the line :

$$\frac{x-1}{1} = \frac{y-2}{2} = \frac{z-1}{3},$$

find the values of a , b , c .

Or

4. (a) Find the correlation coefficient for the following data : [4]

x	y
43	99
21	65
25	79
42	75
57	87
59	81

- (b) Prove (any one) : [4]

(i) $\nabla^2 \left[\nabla \cdot \left(\frac{\bar{r}}{r^2} \right) \right] = \frac{2}{r^4}$

(ii) $\nabla \times \left[\frac{1}{r} \left(r^2 \bar{a} + (\bar{a} \cdot \bar{r}) \bar{r} \right) \right] = 0.$

- (c) Show that : [4]

$$\bar{F} = (2xz^3 + 6y) i + (6x - 2yz) j + (3x^2z^2 - y^2) k$$

is irrotational. Find the scalar potential associated with it.

5. (a) Evaluate : [4]

$$\int_C \bar{F} \cdot d\bar{r}$$

where

$$\bar{F} = (x^2 + xy) \bar{i} + (x^2 + y^2) \bar{j}$$

and C is the square formed by the lines $x = \pm 1, y = \pm 1$.

- (b) Apply Stokes theorem to evaluate : [5]

$$\iint_S (\nabla \times \bar{F}) \cdot \hat{n} dS$$

where

$$\bar{F} = (x^3 - y^3) \bar{i} - xyz \bar{j} + y^3 \bar{k}$$

and S is the surface $x^2 + y^2 + z^2 - 2x = 1$ above the plane $x = 0$.

- (c) Show that : [4]

$$\iiint_V \frac{dV}{r^2} = \iint_S \frac{\bar{r} \cdot \hat{n}}{r^2} dS.$$

Or

6. (a) Using Green's theorem, evaluate : [4]

$$\int_C \bar{F} \cdot d\bar{r}$$

where

$$\bar{F} = x^2 \bar{i} + xy \bar{j}$$

and C is the region enclosed by $y = x^2$ and the line $y = x$.

- (b) Apply Stokes theorem to evaluate : [5]

$$\int_C \left((x^2 + y^2) \vec{i} - 2xy \vec{j} \right) \cdot d\vec{r}$$

where C is the rectangle bounded by the lines $x = \pm a$, $y = 0$ and $y = b$.

- (c) Evaluate : [4]

$$\iiint_S 2x^2 y \, dy \, dz - y^2 \, dz \, dx + 4xz^2 \, dx \, dy$$

over the curved surface of the cylinder $y^2 + z^2 = 16$, bounded by $x = 0$ and $x = 2$.

7. (a) If $u = x^3 - 3xy^2$, find value of v such that $w = u + iv$ is an analytic function. Write w in term of z . [4]

- (b) Evaluate : [4]

$$\oint_C \frac{2z^2 + z + 5}{\left(z - \frac{3}{2}\right)^2} dz$$

where C is the ellipse :

$$\frac{x^2}{9} + \frac{y^2}{16} = 1.$$

- (c) Find the image of straight lines $x = 1$ and $y = 1$ under the transformation $w = z^2$. [5]

Or

8. (a) Show that real and imaginary parts of an analytic function are always harmonic. [4]

- (b) Using Cauchy's residue theorem evaluate : [4]

$$\oint_C \frac{12z - 7}{(z - 1)^2 (2z + 3)} dz$$

where C is the circle :

$$|z + i| = \sqrt{3}.$$

- (c) Find the bilinear transformation which maps the points $z = 1, i, 2i$ on the points $w = -2i, 0, 1$ respectively. [5]