A nonabelian Fourier transform for unipotent representations of exceptional p-adic groups

Albert Lopez Bruch

Abstract

In this paper, we prove that the elliptic dual nonabelian Fourier transform defined by Aubert–Ciubotaru–Romano in the elliptic space of unipotent representations of a split, simple, simply connected p-adic group G of type F_4 or E_6 is compatible with Lusztig's non-abelian Fourier transform and restrictions to reductive quotients of maximal open compact subgroups. Our work naturally builds on previous work, where this conjectural property of the dual elliptic nonabelian Fourier transform was stated and proven for $G = \mathrm{SL}_n$, PGL_n, SO_{2n+1}, Sp_4 or G_2.

Contents

1 Introduction **2**

2 Unipotent representations for finite groups of Lie type **6**

3 Unipotent representations of p-adic groups **10**

3.1 The Unipotent Langlands classification 12

3.2 Chevalley involutions and the dual nonabelian Fourier transform 13

3.3 Action of the fundamental group 16

3.4 The $\mathrm{SL}_n(F)$ case 20

4 Parahoric restriction of unipotent representations **22**

4.1 Reduction to representations over reflection groups 22

4.2 The Iwahori-spherical case and the Springer correspondence 25

4.3 Non-Iwahori-spherical case and weight diagrams 28

4.4 Parahoric restriction for type G_2 30

5 The p-adic group of type F_4 **33**

5.1 Examples of parahoric restrictions 36

6 The p-adic group of type E_6 **39**

7 Parahoric restriction tables **43**

7.1 Parahoric restriction tables for F_4 44

7.2 Parahoric restriction tables for E_6 50

8 Explicit parahoric restriction of $\pi(D_4(71), -)$ wrt K_4 **63**

1 Introduction

The overarching aim of the Local Langlands Correspondence (LLC) is to classify the smooth admissible complex representations of reductive p-adic groups in terms of geometric and number theoretic data. More concretely, if G is a reductive p-adic group over some non-archimedean field F, the Langlands philosophy predicts that the irreducible (smooth admissible) complex representations $\mathrm{Irr}(G)$ should be controlled by the Weil group $\mathcal{W}_F$ of the field F, together with the geometry of the complex dual group $G^\vee$ of G. In [Lus95a], Lusztig defined the subset of irreducible unipotent representations $\mathrm{Irr}_{\mathrm{un}}(G)$. This is the smallest subset of $\mathrm{Irr}(G)$ containing all irreducible representations containing an Iwahori-fixed vector (also denoted Iwahori-spherical) and closed under the partition of $\mathrm{Irr}(G)$ into L-packets. More concretely, the Local Langlands Correspondence predicts the existence of a bijection,

$$\begin{aligned} \mathrm{LLC}_{\mathrm{un}} : \{(x,\phi) \mid x \in G^\vee, \phi \in \widehat{A_x}\} &\longleftrightarrow \mathrm{Irr}_{\mathrm{un}}(G), \\ (x,\phi) &\longmapsto \pi(x,\phi), \end{aligned} \tag{1}$$

where A_x is the component group of the centralizer $Z_{G^\vee}(x)$, and the left hand side can be seen to be in bijection with unramified Langlands parameters (trivial on the inertia subgroup). In [KL87], Kazhdan–Lusztig had already observed that some of the labels $\{(x,\phi) \mid x \in G^\vee, \phi \in \widehat{A_x}\}$ naturally parametrized Iwahori-spherical representations. The Langlands parametrization for the set of unipotent representations has come after many years of development. Lusztig's work [Lus95a], [Lus02] settled it for adjoint p-adic groups, Reeder [Ree02] for Iwahori-spherical representations of split groups of arbitrary isogeny, and Solleveld for all reductive p-adic groups [Sol23a],[Sol23b]. In addition, Lusztig characterized unipotent representations in terms of the structure of G and representation theory of reductive quotients of its parahoric subgroups, which we revisit in Sections 2 and 3. Deligne–Lusztig theory provides a powerful tool to study representations of finite groups of Lie type and, a posteriori, of p-adic groups.

While many questions in the unipotent LLC have now been resolved in full generality, some of them remain open. One such question relates to the stability of L-packets. Moeglin and Waldspurger tackled this problem for $G = \mathrm{SO}_{2n+1}(F)$ in [MW03] by considering the restriction of unipotent representations to maximal open compact subgroups. These subgroups provide a passage from unipotent representations of the p-adic group to unipotent representations of certain finite reductive groups. Work of

Lusztig, Deligne and others in the late 20th century revealed remarkable structure on unipotent characters of these finite reductive groups, which Moeglin and Waldspurger exploited. Lusztig defined an involution on the space of virtual unipotent characters known as *Lusztig's non-abelian transform* taking the irreducible characters to the so-called *almost characters*. These characters can be obtained independently via the theory of character sheaves and, as a result, have nice geometrical and stability properties.

The hope is that, if we can lift the Fourier transform to the setting of p-adic groups with analogous properties, one could use the resulting virtual characters to prove stability properties of L-packets. Moreover, it is believed by Lusztig that an analogous theory of character sheaves for p-adic groups should exist. Work of Ciubotaru [Ciu20] and Aubert–Ciubotaru–Romano [ACR25] has already considered this problem and formulated precise conjectures. We consider the same setting and similar notation to [ACR25], which we now describe in the simplified setting when G is of simply connected type.

Let $\max(G)$ denote the set of conjugacy classes of maximal open compact subgroups of G, which coincides with the set of conjugacy classes of maximal parahoric subgroups since G is simply connected. For each $K \in \max(G)$, we define the *parahoric restriction map* Does it make much sense to avoid this in the introduction? Feels like I need to introduce U_K, $\overline{K}$ and the representation spaces anyway.

$$\begin{aligned} \mathrm{res}^K_{\mathrm{un}} : R_{\mathrm{un}}(G) &\longrightarrow R_{\mathrm{un}}(\overline{K}) \\ V &\longmapsto V^{U_K}, \end{aligned}$$

where U_K is the pro-p unipotent radical K, $\overline{K} := K/U_K$ is a connected finite group of Lie type and $R_{\mathrm{un}}(G)$ and $R_{\mathrm{un}}(\overline{K})$ are the complexification of the Grothendieck groups spanned by the unipotent representations of G and $\overline{K}$, respectively. Secondly, Lusztig's nonabelian Fourier transforms on $R_{\mathrm{un}}(\overline{K})$ for each $K \in \max(G)$ gives an involution

$$\mathrm{FT}^{\mathrm{cpt}}_{\mathrm{un}} : \bigoplus_{K \in \max(G)} R_{\mathrm{un}}(\overline{K}) \longrightarrow \bigoplus_{K \in \max(G)} R_{\mathrm{un}}(\overline{K}).$$

The remaining ingredient to formally state Aubert–Ciubotaru–Romano's conjecture [ACR25, Conjecture 1.2] is the construction of the *dual nonabelian Fourier transform* $\mathrm{FT}^\vee : R_{\mathrm{un}}(G) \longrightarrow R_{\mathrm{un}}(G)$ compatible with the unipotent restriction maps $\mathrm{res}^K_{\mathrm{un}}$ and Lusztig's nonabelian Fourier transform $\mathrm{FT}^{\overline{K}}$. The existence of such a map is conjectured, but so far no explicit general construction has been achieved on the infinite-dimensional space $R_{\mathrm{un}}(G)$. Instead, Ciubotaru–Opdam [CO17] realized that, for applications, it was often sufficient to study the quotient space $\overline{R}_{\mathrm{un}}(G)$ of $R_{\mathrm{un}}(G)$ by the span of all properly parabolically induced representations, since parabolic induction is generally well understood. The upshot of this simplification is double: firstly, the quotient space $\overline{R}_{\mathrm{un}}(G)$ is finite dimensional, in contrast to the infinite-dimensional space $R_{\mathrm{un}}(G)$. Secondly, $\overline{R}_{\mathrm{un}}(G)$ is canonically isomorphic (in fact,

isometric with respect to some canonical inner products) to the elliptic unipotent representation space $R_{\mathrm{un,ell}}(G)$, a certain subspace of $R_{\mathrm{un}}(G)$. Moreover, the Local Langlands Correspondence provides an explicit description for a natural orthogonal basis of $R_{\mathrm{un,ell}}(G)$. To describe this basis, let $u \in G^\vee$ be a unipotent element and let Γ_u be the reductive part of the centralizer $Z_{G^\vee}(u)$. We then consider the set of elliptic pairs

$$\mathcal{Y}(\Gamma_u)_{\mathrm{ell}} = \{(s,h) \in \Gamma_u \mid sh = hs \text{ and } Z_{\Gamma_u}(s,h) \text{ finite}\}$$

and, for each such elliptic pair (s,h), we define the virtual characters

$$\Pi(u,s,h) := \sum_{\phi \in A_{us}} \phi(h)\pi(us,\phi).$$

The Local Langlands Correspondence induces an isomorphism

$$\mathrm{LLC}^{\mathrm{ell}}_{\mathrm{un}} : \bigoplus_{u \in G^\vee} \mathbb{C}[\mathcal{Y}(\Gamma_u)]^{\Gamma_u} \longrightarrow R_{\mathrm{un,ell}}(G), \qquad (u,s) \longmapsto \Pi(u,s,h),$$

where the sum in the left hand side is taken over conjugacy classes of unipotent elements in $G^\vee$, so in particular $R_{\mathrm{un,ell}}(G)$ is the span of $\{\Pi(u,s,h) \mid u \in G^\vee \text{ unipotent}, (s,h) \in \mathcal{Y}(\Gamma_u)_{\mathrm{ell}}\}$, where the parameters are taken up to $G^\vee$-conjugation.

The space of elliptic pairs has an obvious involution given by the flip $(s,h) \mapsto (h,s)$, and this defines an involution

$$\mathrm{FT}^\vee_{\mathrm{ell}} : R_{\mathrm{un,ell}}(G) \longrightarrow R_{\mathrm{un,ell}}(G),$$

the dual elliptic nonabelian Fourier transform. We are finally ready to state the main conjecture of this paper, in the simplified setting of split, simply connected p-adic groups.

Conjecture 1.1. [ACR25, Conjecture 1.2] Let G be a split semisimple simply connected p-adic group. Then the diagram

$$\begin{array}{ccc} R_{\mathrm{un,ell}}(G) & \xrightarrow{\mathrm{FT}^\vee_{\mathrm{ell}}} & R_{\mathrm{un,ell}}(G) \\ \downarrow \mathrm{res}^{\mathrm{cpt}}_{\mathrm{un}} & & \downarrow \mathrm{res}^{\mathrm{cpt}}_{\mathrm{un}} \\ \bigoplus_{K \in \max(G)} R_{\mathrm{un}}(\overline{K}) & \xrightarrow{\mathrm{FT}^{\mathrm{cpt}}_{\mathrm{un}}} & \bigoplus_{K \in \max(G)} R_{\mathrm{un}}(\overline{K}), \end{array}$$

commutes, up to certain roots of unity.

In [ACR25], Aubert–Ciubotaru–Romano stated a more general conjecture including all isogeny classes, but the technical details are unnecessary for our purposes. Moreover, they showed that this conjecture (or the more general version) holds for $G = \mathrm{SL}_n(F)$, $\mathrm{PGL}_n(F)$ or $\mathrm{Sp}_4(F)$. In previous work, Ciubotaru and Opdam [Ciu20] showed it holds for $G_2(F)$, and Waldspurger [Wal18] showed it for $G = \mathrm{SO}_{2n+1}(F)$. In this document we settle the above conjecture for two further exceptional cases.

Theorem 1.2. *Let G be a simple, split, simply connected p-adic group of type F_4 or E_6. Then Conjecture 1.1 holds for G.*

Structure of the document

The document is organized as follows. We devote Section 2 to finite groups of Lie type and Deligne–Lusztig theory, with a focus on the construction of Lusztig's nonabelian Fourier transform. In Section 3 we discuss representation theoretic aspects of p-adic groups and their relation to representations of finite groups of Lie type. We recall the definition of unipotent representations and construct the dual nonabelian Fourier transform. Section 4 is the most technical; there we discuss in detail how to explicitly compute the restriction functors $\mathrm{res}^{\mathrm{cpt}}_{\mathrm{un}}$, a method developed by Reeder in [Ree00] for adjoint groups and which we adapt for simply connected groups. Sections 5 and 6 contain geometric information about the complex reductive groups $G^\vee = F_4(\mathbb{C})$ or $E_6(\mathbb{C})_{\mathrm{sc}}$ necessary to explicitly describe the virtual representations $\Pi(u, s, h)$ spanning the elliptic unipotent representation space $R_{\mathrm{un},\ell}(G)$.

In Section 7, we display all tables containing the parahoric restrictions of unipotent representations with respect to maximal parahoric subgroups. From these tables, the proof of Conjecture 1.1 is a simple verification.

2 Unipotent representations for finite groups of Lie type

In this first section, we recall the necessary structural and representation theoretic properties of finite groups of Lie type leading to Lusztig's construction of the nonabelian Fourier transform described in the introduction. To this aim, suppose that $\mathbb{G}$ is a connected reductive group over $\overline{\mathbb{F}}_p$ with Frobenius map $\mathrm{Fr} : \mathbb{G} \to \mathbb{G}$. The group $\mathbb{G}^{\mathrm{Fr}}$ is a finite group of Lie type and we are interested in studying the set of irreducible unipotent representations of $\mathbb{G}^{\mathrm{Fr}}$ denoted by $\mathrm{Irr}_{\mathrm{un}}(\mathbb{G}^{\mathrm{Fr}})$. We say that $\mathbb{G}^{\mathrm{Fr}}$ is *split* if the induced Frobenius action on the set of simple roots of $\mathbb{G}$ is trivial. We shall assume throughout that $\mathbb{G}^{\mathrm{Fr}}$ is a split group. We recall that Deligne–Lusztig induction associates, to each pair $(\mathbb{T}, \theta)$ where $\mathbb{T}$ is a Fr-stable maximal tori of $\mathbb{G}$ and θ is a character of $\mathbb{T}^{\mathrm{Fr}}$, a virtual character $R_{\mathbb{T},\theta}$. Then we define

$$\mathrm{Irr}_{\mathrm{un}}(\mathbb{G}^{\mathrm{Fr}}) = \{\rho \in \mathrm{Irr}(\mathbb{G}^{\mathrm{Fr}}) \mid \langle \rho, R_{\mathbb{T},\mathbf{1}} \rangle \neq 0 \text{ for some Fr-stable maximal torus } \mathbb{T} \text{ of } \mathbb{G}\}.$$

If the Fr-stable maximal torus $\mathbb{T}$ is maximally split and therefore contained in some Fr-stable Borel subgroup $\mathbb{B}$, then $R_{T_e,\mathbf{1}} = \mathrm{Ind}_{\mathbb{B}^{\mathrm{Fr}}}^{\mathbb{G}^{\mathrm{Fr}}} \mathbf{1}$, so the set of unipotent characters contain the principal series representations of $\mathbb{G}^{\mathrm{Fr}}$. By results of Lusztig, the classification of $\mathrm{Irr}_{\mathrm{un}}(\mathbb{G}^{\mathrm{Fr}})$ is reduced to the case when $\mathbb{G}$ is adjoint simple.

Let $W = N_{\mathbb{G}}(\mathbb{T})/\mathbb{T}$ be the Weyl group associated to $\mathbb{G}$. When the group $\mathbb{G}^{\mathrm{Fr}}$ is split, irreducible representations ϕ of W are known to to parametrize principal series representations χ_ϕ of $\mathbb{G}^{\mathrm{Fr}}$, and we shall make use of this bijection throughout. maybe mention here that other representation series are also in bijection with weyl groups of smaller types?. In addition, for each $\phi \in \widehat{W}$, define the *almost character*

$$R_\phi := \frac{1}{|W|} \sum_{w \in W} \phi(w) R_{\mathbb{T}_w,1},$$

arising as traces of certain character sheaves on the group $\mathbb{G}$. One can then partition $\mathrm{Irr}_{\mathrm{un}}(\mathbb{G}^{\mathrm{Fr}})$ into families by defining two representations ρ_1, ρ_2 to be related if

$$\langle \rho_1, R_\phi \rangle \neq 0 \quad \text{and} \quad \langle \rho_2, R_\phi \rangle \neq 0 \quad \text{for some } \phi \in \widehat{W},$$

and then extending by transitivity. To each family $\mathcal{U} \subset \mathrm{Irr}_{\mathrm{un}}(\mathbb{G}^{\mathrm{Fr}})$, Lusztig associated a finite group $\Gamma_{\mathcal{U}}$, scalars $\Delta(x,\sigma) \in \{\pm 1\}, (x,\sigma) \in \mathcal{M}(\Gamma_{\mathcal{U}})$ defined in [Lus84] and a bijection

$$\begin{aligned} \mathcal{U} &\longleftrightarrow \mathcal{M}(\Gamma_{\mathcal{U}}) := \{(x,\sigma) \mid x \in \Gamma_{\mathcal{U}}, \sigma \in \mathrm{Irr}(Z_{\Gamma_{\mathcal{U}}}(x))\}/\Gamma_{\mathcal{U}} \\ \rho_{(x,\sigma)} &\longleftrightarrow (x,\sigma). \end{aligned}$$

The indexing set $\mathcal{M}(\Gamma_{\mathcal{U}})$ carries a natural pairing

$$\{(x,\sigma),(y,\tau)\} = \frac{1}{|Z_{\Gamma_{\mathcal{U}}}(x)||Z_{\Gamma_{\mathcal{U}}}(y)|} \sum_{\substack{g \in \Gamma_{\mathcal{U}} \\ xgyg^{-1} = gygx^{-1}}} \sigma(gyg^{-1})\tau(g^{-1}x^{-1}g),$$

and the above bijection satisfies the property that

$$\langle \rho_{(x,\rho)}, R_\phi \rangle_{\mathbb{G}^{\mathrm{Fr}}} = \begin{cases} \Delta(x,\sigma)\,\{(x,\rho),(y,\tau)\} & \text{if } \chi_\phi = \rho_{(y,\tau)} \in \mathcal{U}, \\ 0 & \text{if } \chi_\phi \notin \mathcal{U}. \end{cases} \tag{2}$$

We recall that if W is an irreducible Weyl group, then $\Delta(x,\rho) = 1$ except for the families containing $\phi_{512,11}$ in E_7, $\phi_{4096,11}$ in E_8 or $\phi_{4096,26}$ in E_8. For any family $\mathcal{U}$, $\rho_{(e,1)} = \chi_\phi$ is a principal series representation corresponding to some representation ϕ of W. Such representations of W are called *special*, and they are the only ones satisfying the property that $\langle \chi, R_\phi \rangle \geq 0$ for all unipotent characters χ (see [Car85, §12.3]). We define the *unipotent almost characters* of $\mathbb{G}^{\mathrm{Fr}}$ to be the set of orthonormal class functions

$$R_{(x,\sigma)} := \sum_{(y,\tau)} \Delta(y,\tau)\{(y,\tau),(x,\sigma)\}\rho_{(y,\tau)}, \quad (x,\sigma) \in \mathcal{M}(\Gamma_{\mathcal{U}}).$$

In particular, if $\chi_\phi = \rho_{(x,\rho)}$, then $R_\phi = R_{(x,\rho)}$. Lusztig also defined the map

$$\begin{aligned} \mathrm{FT}_{\Gamma_{\mathcal{U}}} : \mathbb{C}[\mathcal{U}] &\longrightarrow \mathbb{C}[\mathcal{U}] \\ \rho_{(x,\sigma)} &\longmapsto \sum_{(y,\tau)\in\mathcal{M}(\Gamma_{\mathcal{U}})} \{(x,\sigma),(y,\tau)\}\rho_{(y,\tau)} \end{aligned}$$

The *unipotent nonabelian Fourier transform* of $\mathbb{G}^{\mathrm{Fr}}$ on the space $R_{\mathrm{un}}(\mathbb{G}^{\mathrm{Fr}}) = \oplus_{\mathcal{U}\subseteq \mathrm{Irr}_{\mathrm{un}}(\mathbb{G}^{\mathrm{Fr}})}\mathbb{C}[\mathcal{U}]$ is the map

$$\mathrm{FT}^{\mathbb{G}^{\mathrm{Fr}}} := \bigoplus_{\mathcal{U}\subset \mathrm{Irr}_{\mathrm{un}}(\mathbb{G}^{\mathrm{Fr}})} \mathrm{FT}_{\Gamma_{\mathcal{U}}},$$

and it satisfies many interesting properties. Most importantly, it is a self-Hermitian involution with respect to the standard inner product of characters and, in addition, $\mathrm{FT}^{\mathbb{G}^{\mathrm{Fr}}}(\chi_\phi) = R_\phi$. More generally, it takes unipotent representations to the almost characters arising as traces of character sheaves on $\mathbb{G}$. Let us discuss some important examples that will be used in our tables in Section 7.

Example 2.1. When $\mathbb{G} = \mathrm{PGL}_n(\overline{\mathbb{F}_p})$ and Fr is the standard q-th power Frobenius, then $\mathbb{G}^{\mathrm{Fr}} = \mathrm{PGL}_n(\mathbb{F}_q)$ and $W \cong S_n$. In this case, one can show that $R_\phi = \chi_\phi$ for all $\phi \in \widehat{S_n}$ so, in particular, all families have a unique element and unipotent representations coincide with the principal series. Thus, unipotent representations of $\mathbb{G}^{\mathrm{Fr}}$ are naturally by partitions of n, and we shall use this notation in the tables from Section 7.

Example 2.2. Suppose $\mathbb{G}^{\mathrm{Fr}} = \mathrm{SO}_{2n+1}(\mathbb{F}_q)$ or $\mathbb{G}^{\mathrm{Fr}} = \mathrm{PGSp}_{2n}(\mathbb{F}_q)$ (split, types B_n or C_n), and with Weyl group $W \cong S_n \rtimes C_2^n$. Its representations are parametrized by double partitions of n, and we shall use this notation to label principal series representations. In turn, double partitions of n are in bijection with symbols

$$\begin{pmatrix} \lambda_1, & \lambda_2, & \lambda_3, \ldots, & \lambda_{k+s} \\ & \mu_1, & \mu_2, & \ldots, \mu_k \end{pmatrix}$$

with $0 \leq \lambda_1 < \lambda_2 < \cdots < \lambda_{k+s}$, $0 \leq \mu_1 < \mu_2 < \cdots < \mu_k$, with $\lambda_1\mu_1 \neq 0$, rank n (should define rank

or refer to a definition) and defect $s = 1$. To account for the remaining unipotent representations of $\mathbb{G}^{\mathrm{Fr}}$, we allow for symbols with positive odd defect s. Remarkably, Lusztig showed that two unipotent representations lie in the same family if and only if the unordered set $\{\lambda_1, \dots, \lambda_{k+s}, \mu_1, \dots, \mu_k\}$ coincide. See [Car85, §11.4 and §13.8] for a complete overview of the theory.

For example, there are 10 double partitions of $n = 3$ (see Table 12) and the two symbols $\theta_1 = \left(\begin{smallmatrix} 0 & 1 & 2 & 3 \\ & 1 & & \end{smallmatrix}\right)$ and $\theta_2 = \left(\begin{smallmatrix} 0 & 1 & 3 \\ & - & \end{smallmatrix}\right)$ are the only two rank-3 symbols with higher defect (see Table 13). The 12 unipotent representations are partitioned in 4 trivial families and 2 families of size 4 containing the symbols

$$\left\{[1,11] = \left(\begin{smallmatrix} 0 & 1 & 3 \\ 1 & 2 & \end{smallmatrix}\right),\ [\cdot,12] = \left(\begin{smallmatrix} 0 & 1 & 2 \\ 1 & 3 & \end{smallmatrix}\right),\ [1^3,\cdot] = \left(\begin{smallmatrix} 1 & 2 & 3 \\ 0 & 1 & \end{smallmatrix}\right),\ \theta_1 = \left(\begin{smallmatrix} 0 & 1 & 2 & 3 \\ & 1 & & \end{smallmatrix}\right)\right\} \quad \text{and}$$
$$\left\{[2,1] = \left(\begin{smallmatrix} 0 & 3 \\ 1 & \end{smallmatrix}\right),\ [\cdot,3] = \left(\begin{smallmatrix} 0 & 1 \\ 3 & \end{smallmatrix}\right),\ [12,\cdot] = \left(\begin{smallmatrix} 1 & 3 \\ 0 & \end{smallmatrix}\right),\ \theta_2 = \left(\begin{smallmatrix} 0 & 1 & 3 \\ & - & \end{smallmatrix}\right)\right\}$$

When $n = 4$, there are 20 double partitions (see Table 5) and 5 rank-4 symbols with higher defect, namely $\theta_1 = \left(\begin{smallmatrix} 0 & 1 & 2 & 3 & 4 \\ & 1 & 2 & & \end{smallmatrix}\right)$, $\theta_2 = \left(\begin{smallmatrix} 0 & 1 & 2 & 3 \\ & & 2 & & \end{smallmatrix}\right)$, $\theta_3 = \left(\begin{smallmatrix} 0 & 1 & 2 & 4 \\ & 1 & & \end{smallmatrix}\right)$, $\theta_4 = \left(\begin{smallmatrix} 0 & 2 & 3 \\ & - & \end{smallmatrix}\right)$ and $\theta_5 = \left(\begin{smallmatrix} 0 & 1 & 4 \\ & - & \end{smallmatrix}\right)$ (see Table 6). These unipotent representations all lie in a family of size 4 whose associated group is $\Gamma \cong C_2$. As an example, the Fourier transform for these families is

$$\begin{pmatrix} 1/2 & 1/2 & 1/2 & 1/2 \\ 1/2 & 1/2 & -1/2 & -1/2 \\ 1/2 & -1/2 & 1/2 & -1/2 \\ 1/2 & -1/2 & -1/2 & 1/2 \end{pmatrix}.$$

Example 2.3. Consider the split finite group of Lie type $\mathbb{G}^{\mathrm{Fr}} = F_4(\mathbb{F}_q)$. The Weyl group W is a Coxeter group of order 1152 and 25 irreducible representations labelled as $\phi_{d,k}$, where d is the dimension and k is its b-invariant. The outer automorphism of W swapping long and short roots acts on $\widehat{W}$ while preserving these invariants, so for some pairs d, k one has two inequivalent representations labelled by $\phi'_{d,k}$ and $\phi''_{d,k}$. In addition to the principal series representations, $\mathrm{Irr}_{\mathrm{un}}(\mathbb{G}^{\mathrm{Fr}})$ contains 7 cuspidal representations labelled by $F_4^I[1], F_4^{II}[1], F_4[-1], F_4[\theta], F_4[\theta^2], F_4[i]$ and $F_4[-i]$. Finally, it contains 5 more unipotent representations arising as irreducible components of the parabolic induction of the unique unipotent cuspidal representation of a Levi of type B_2, labelled as $B_{2,\mathbf{1}}$, $B_{2,\varepsilon}$, $B_{2,\varepsilon'}$, $B_{2,\varepsilon''}$ and $B_{2,r}$.

The set $\mathrm{Irr}_{\mathrm{un}}(\mathbb{G}^{\mathrm{Fr}})$ is then partitioned in 11 families: there are 8 families consisting of a single representation, 2 families $\{\phi_{4,1}, \phi'_{2,4}, \phi''_{2,4}, B_2, \mathbf{1}\}$, $\{\phi_{4,13}, \phi'_{2,16}, \phi''_{2,16}, B_{2,\varepsilon}\}$ whose associated group is $\Gamma \cong C_2$ and labels $(1,1), (1,\varepsilon), (g_2,1), (g_2,\varepsilon)$, and one large family of size 21 whose associated group is $\Gamma \cong S_4$. See Table 1 for the representations and their labels in $\mathcal{M}(S_4)$.

$\phi_{12,4}$	$\phi''_{9,6}$	$\phi'_{9,6}$	$\phi''_{1,12}$	$\phi'_{1,12}$	$\phi''_{4,7}$	$\phi'_{4,7}$	$\phi_{4,8}$	$\phi'_{6,6}$	$\phi''_{6,6}$	$\phi_{16,5}$	$B_{2,r}$	$B_{2,\varepsilon'}$	$B_{2,\varepsilon''}$	$F_4[\theta]$	$F_4[\theta^2]$	$F_4[i]$	$F_4[-i]$	$F_4^I[1]$	$F_4^{II}[1]$	$F_4[-1]$
$(1,1)$	$(g_2',1)$	$(1,\lambda^1)$	(g_2',ε')	$(1,\lambda^2)$	$(g_4,1)$	(g_2,ε'')	(g_2',ε'')	$(g_3,1)$	$(1,\sigma)$	$(g_2,1)$	(g_2',r)	$(g_4,-1)$	(g_2,ε')	(g_3,θ)	(g_3,θ^2)	(g_4,i)	$(g_4,-i)$	(g_2',ε)	$(1,\lambda^3)$	(g_2,ε)

Table 1: Largest $F_4(\mathbb{F}_q)$-family of unipotent representations and their labels

When $\mathbb{G}^{\mathrm{Fr}}$ is split (as it is the case throughout this document) it is convenient to express Lusztig's nonabelian Fourier transform in term of stable combinations of unipotent representations. This makes the verification of Conjecture 1.1 cleaner and more straightforward. Let $\mathcal{U}$ be a family of unipotent representations and for each $x \in \Gamma_{\mathcal{U}}$, define the virtual combinations of unipotent characters

$$\Pi_{\mathcal{U}}(x,y) = \sum_{\sigma \in \widehat{Z_{\Gamma_{\mathcal{U}}}(x)}} \sigma(y^{-1})\rho_{(x,\sigma)}, \quad \text{where } y \in Z_{\Gamma_{\mathcal{U}}}(x). \tag{3}$$

Dually, when $\Gamma_{\mathcal{U}}$ is abelian, define the virtual combinations

$$\Pi_{\mathcal{U}}(\sigma,\tau) = \sum_{y \in \Gamma_{\mathcal{U}}} \tau(y)\rho_{(y,\sigma)}, \quad \text{if } \sigma,\tau \in \widehat{\Gamma_{\mathcal{U}}}. \tag{4}$$

Lemma 2.4 ([ACR25, Lemma 5.1])**.** *With the notation of* (3) *and* (4)*,*

$$\mathrm{FT}_{\Gamma_{\mathcal{U}}}(\Pi_{\mathcal{U}}(x,y)) = \Delta(x,y)\Pi_{\mathcal{U}}(y,x), \quad \mathrm{FT}_{\Gamma_{\mathcal{U}}}(\Pi_{\mathcal{U}}(\sigma,\tau)) = \Pi_{\mathcal{U}}(\tau,\sigma),$$

the latter when $\Gamma_{\mathcal{U}}$ *is abelian, where* $\Delta(x,y) = 1$ *unless* $\mathcal{U}$ *is one of the families described in* (2)*, in which case* $\Delta(1,g_2) = \Delta(g_2,1) = -1$ *and* 1 *otherwise.*

3 Unipotent representations of p-adic groups

Let F be a finite extension of $\mathbb{Q}_p$ with residue field $\mathbb{F}_q$, and let $\mathcal{G}$ be a connected simple algebraic F-group split over F and write $G = \mathcal{G}(F)$. Let $\mathcal{T}$ be a F-split maximal torus of $\mathcal{G}$ and let $T = \mathcal{T}(F)$ be a maximal torus of G. We assume throughout that $\mathcal{G}$ is in addition of simply-connected type.

Let $\mathrm{Irr}(G)$ be the set of irreducible smooth representations of G and $\mathrm{Irr}_{\mathrm{un}}(G) \subseteq \mathrm{Irr}(G)$ the subset of unipotent representations. Under the Local Langlands correspondence, there is a bijection

$$\mathrm{LLC}_{\mathrm{un}} : \left\{ \begin{array}{c} \text{Unipotent representations} \\ \text{of } G(F) \end{array} \right\}_{/\cong} \longleftrightarrow \left\{ \begin{array}{c} \text{Pairs } (x, \phi): \\ x \in G^\vee,\ \phi \in \widehat{A_x} \end{array} \right\}_{/G^\vee},$$

By the work of Lusztig, unipotent representations of G can also be characterized directly in terms of the topology of G and the representation theory of finite groups of Lie type. For each facet c in the Bruhat–Tits building $\mathcal{B}(G, F)$, consider the parahoric subgroup $K_c := \mathrm{Stab}_G(c)$. Since, by assumption, G is simply connected, K_c is a connected open compact subgroup of G fitting in a short exact sequence

$$1 \longrightarrow U_c \longrightarrow K_c \longrightarrow \overline{K}_c \longrightarrow 1,$$

where U_c is the pro-unipotent radical of K_c and $\overline{K}_c$ is the group of $\mathbb{F}_q$-rational points of a connected reductive group $\overline{\mathbf{K}}_{\mathbf{c}}$. Since $K_{g\cdot c} = gK_cg^{-1}$ for all facets c of $\mathcal{B}(G, F)$ and $g \in G$, conjugacy classes of parahoric subgroups are in bijection with G-orbits of facets in $\mathcal{B}(G)$. Furthermore, after fixing a set of simple affine roots S_{aff} of G with corresponding fundamental alcove $\mathcal{C}_0$, there is a bijection

$$\{\text{Subsets } J \subsetneq S_{\mathrm{aff}}\} \longleftrightarrow \{G\text{-orbits of facets in } \mathcal{B}(G)\} \longleftrightarrow \{\text{Conj. classes of parahoric subgroups}\}$$
$$J \longmapsto c_J = \{x \in \overline{\mathcal{C}_0} \mid \langle \alpha, x\rangle \in \mathbb{Z} \iff \alpha \in J, \text{ for any } \alpha \in S_{\mathrm{aff}}\} \longmapsto K_{c_J}$$

satisfying that $J_1 \subseteq J_2$ if and only if $c_{J_2} \subset \overline{c_{J_1}}$ if and only if $K_{c_{J_1}} \subseteq K_{c_{J_2}}$. Throughout, we denote by $(K_J, U_J, \overline{K}_J)$ the parahoric subgroup corresponding to the facet c_J. In particular, maximal parahoric subgroups correspond to facets of minimal dimension and maximal subsets $J \subsetneq S_{\mathrm{aff}}$. When G is simply connected, we recall that the set of maximal parahoric subgroups coincides with the set of maximal open compact groups. We define

$$\max(G) := \{K_x \mid \{x\} \subset \mathcal{B}(G) \text{ facet of minimal dimenson}\}$$

to be the set containing each orbit of maximal parahoric subgroups.

Next, let (π, V) a smooth admissible complex representation of G and let $(K, U, \overline{K})$ be a parahoric subgroup. The space V^{U_K} of fixed points under the pro-unipotent radical is naturally a representation

of $\overline{K} = K/U_K$. This induces the *parahoric restriction functor*

$$\mathrm{res}_K : R(G) \longrightarrow R(\overline{K}), \quad V \longmapsto V^{U_K}, \quad \text{for all } V \in \mathrm{Irr}(G), \tag{5}$$

where $R(G)$ and $R(\overline{K})$ are the complexification of the Grothendieck group of $\mathrm{Irr}(G)$ and $\mathrm{Irr}(\overline{K})$, respectively. If (σ, E) is a cuspidal representation of $\overline{K}$, we define

$$\mathrm{Irr}(G, K, E) = \{(\pi, V) \in \mathrm{Irr}(G) \mid \text{ the } \overline{K}\text{-module } V^{U_K} \text{ contains } E\}.$$

Definition 3.1. We say that an irreducible representation (π, V) of G is *unipotent* if there is a parahoric subgroup $(K, U_K, \overline{K})$ such that V^{U_K} contains a cuspidal unipotent representation of $\overline{K}$; that is, if $(\pi, V) \in \mathrm{Irr}(G, K, E)$ for some pair (K, E) where E is unipotent.

Therefore, the set of unipotent representations, up to isomorphism, is

$$\mathrm{Irr}_{\mathrm{un}}(G) = \bigcup_{\substack{J \subsetneq S_{\mathrm{aff}} \\ E \text{ cusp. unip. } \overline{K}_J\text{-rep}}} \mathrm{Irr}(G, K_J, E).$$

Each set $\mathrm{Irr}(G, K, E)$ generates a distinct *Bernstein block* of representations of G. If $\mathcal{I}$ is an Iwahori subgroup, $\mathcal{I}/U_{\mathcal{I}} \cong (\mathbb{F}_q^\times)^r$ and the only unipotent representation is the trivial character. Thus, Iwahori-spherical representations are all unipotent.

Moreover, unipotent representations behave well under the parahoric restriction functors res^K. The first important result is due to Moy and Prasad. A similar result holds for non-unipotent representations, but it is more complicated.

Proposition 3.2. [Ree00, Lemma 5.1] *Assume G is simply connected. Suppose $I \subsetneq S_{\mathrm{aff}}$ and that V^{U_I} contains a cuspidal unipotent representation σ of $\overline{K}_I$. If $J \subsetneq S_{\mathrm{aff}}$ with $V^{U_J} \neq 0$, and J is minimal with respect to this property $I = J$.*

The following is a direct consequence of the above result.

Corollary 3.3. *For any two pairs (K, E) and (K', E') of a parahoric subgroup and a cuspidal unipotent representation of the reductive quotient,* $\mathrm{Irr}(G, K, E)$ *and* $\mathrm{Irr}(G, K', E')$ *are either disjoint or equal.*

Proposition 3.2 together with a standard Harish–Chandra style argument yields the following corollary.

Corollary 3.4. *Let (π, V) be a unipotent representation of G and let $(K, U_K, \overline{K})$ be a parahoric subgroup. Then the $\overline{K}$-irreducible components of V^{U_K} are all unipotent.*

Consequently, for any parahoric subgroup K, the parahoric restriction functor res^K naturally restricts to the *unipotent parahoric restriction function*

$$\mathrm{res}^K_{\mathrm{un}} : R_{\mathrm{un}}(G) \longrightarrow R_{\mathrm{un}}(\overline{K}), \quad V \longmapsto V^{U_K},$$

where $R_{\mathrm{un}}(G)$ and $R_{\mathrm{un}}(\overline{K})$ are the complexification of the Grothendieck groups spanned by $\mathrm{Irr}_{\mathrm{un}}(G)$ and $\mathrm{Irr}_{\mathrm{un}}(\overline{K})$, respectively. Finally, we let

$$\mathrm{res}^{\mathrm{cpt}}_{\mathrm{un}} := \bigoplus_{K\in\max(G)} \mathrm{res}^{K}_{\mathrm{un}} : R_{\mathrm{un}}(G) \longrightarrow \bigoplus_{K\in\max(G)} R_{\mathrm{un}}(\overline{K}).$$

3.1 The Unipotent Langlands classification

The Local Langlands provides a beautiful classification of the unipotent representations of G in terms of the geometry of the dual group $G^\vee$ of G, of adjoint type. This result is widely known as Lusztig's arithmetic-geometric correspondence.

Theorem 3.5 (The arithmetic-geometric correspondence [Lus95a])**.** *There is a natural bijection*

$$\begin{aligned} G^\vee\backslash\{(x,\phi) \mid x\in G^\vee, \phi\in\widehat{A_x}\} &\longleftrightarrow \mathrm{Irr}_{\mathrm{un}}(G)\\ (x,\phi) &\longmapsto \pi(x,\phi). \end{aligned}$$

In particular, there is an isomorphism

$$\mathrm{LLC}_{\mathrm{un}} : \bigoplus_{x\in G^\vee} R(A_x) \longrightarrow R_{\mathrm{un}}(G), \tag{6}$$

where the sum on the left hand side is taken over conjugacy classes of $G^\vee$ and $R(A_x)$ is the space of virtual characters of A_x.

Given a Langlands parameter (x,ϕ), it is a natural question to ask inside which unipotent block does the representation $\pi(x,\phi)$ lies in. This can be answered using the generalized Springer correspondence, which we will not revise in detail here (see [CMBO25, §4.3]). We only state a fundamental result giving a geometric characterization for a representation $\pi(x,\phi)$ to be Iwahori-spherical, and thus lying in the principal block generated by $\mathrm{Irr}(G,\mathcal{I},\mathbf{1})$. Let $\mathcal{B}^x$ the partial flag variety containing all Borel subgroups of $G^\vee$ containing x. This variety carries a natural action of A_x given by conjugation. In particular, the cohomology complex $H^*(\mathcal{B}^x)$ inherits an action of A_x too.

Theorem 3.6. [KL87, Theorem 7.12] *The representation $\pi(x,\phi)$ is Iwahori-spherical if and only if $H^*(\mathcal{B}^x)^\phi \neq 0$. In particular, for any $x\in G^\vee$, the representation $\pi(x,\mathbf{1})$ is Iwahori-spherical.*

Consequently, unipotent representations can also be described as the smallest class of representations of G containing Iwahori-spherical representations and closed under L-packets.

In later sections, we display parahoric restriction tables for representations of G parametrized by several elements $x\in G^\vee$. To do this, we use the following standard notation for the Langlands parameters as in [Ree00], which we now explain.

Notation 3.7. Given a representation $\pi(x,\phi)$, let $x = us$ be the Jordan decomposition. Then u is a unipotent element of the connected pseudo-Levi subgroup $Z_{G^\vee}(s)^0$, and ϕ is an irreducible representation of the component group A_{us}. The representation $\pi(x,\phi)$ will be labelled by the type of unipotent element u is inside $Z_{G^\vee}(s)^0$, together with the representation ϕ. For example, if $G^\vee = F_4(\mathbb{C})$ and $u \in G^\vee$ is a unipotent element of type $F_4(a_3)$, there is a semisimple element $s \in Z_{G^\vee}(u)$ of order 2 such that $Z_{G^\vee}(s)$ is connected of type $C_3 \times A_1$. The unipotent element u is then subregular in the first component and regular in the second, labelled by $C_3(42) \times A_1$, and $\widehat{A_{us}} = C_2 \times C_2$. Thus, the four representations of $F_4(F)$ in the L-packet parametrized by $x = su \in F_4(\mathbb{C})$ are labelled $[C_3(42)A_1, \pm\pm]$.

3.2 Chevalley involutions and the dual nonabelian Fourier transform

In this section we define a modified version of the *dual elliptic nonabelian Fourier transform*, constructed in [ACR25, §9] Add citations. The missing ingredient to the conjecture is the role played by a Chevalley involution on the Langlands parameters, which we now describe. As usual, we let $G^\vee$ be the complex dual group of G.

Definition 3.8 (cite this better)**.** An automorphism $C : G^\vee \to G^\vee$ is called a *Chevalley involution* if there is some maximal torus $T^\vee$ of $G^\vee$ such that $C(t) = t^{-1}$ for all $t \in T^\vee$.

Let $\mathrm{Aut}(G^\vee)$ be the set of automorphisms of $G^\vee$, $\mathrm{Int}(G^\vee)$ the subset of inner automorphisms and $\mathrm{Out}(G^\vee) = \mathrm{Aut}(G^\vee)/\mathrm{Int}(G^\vee)$. We denote inner automorphisms by $\mathrm{int}(x) : g \mapsto xgx^{-1}$. Using based root data and pinnings, one can show that any Chevalley involution $C : G^\vee \to G^\vee$ exists, satisfies $C^2 = 1$ and for every semisimple element $g \in G$, $C(g)$ is conjugate to g. In addition, if C_1 and C_2 are Chevalley involutions with respect to $T_1^\vee$ and $T_2^\vee$, then they are conjugate by an inner automorphism of of $G^\vee$ (cite here!!). Thus, Chevalley involutions all lie in a distinguished class of $\mathrm{Out}(G^\vee)$ of order dividing 2. The following result characterizes the order of this class depending on the type of $G^\vee$. We insist that $G^\vee$ is assumed to be simple.

Lemma 3.9. *The Chevalley involutions $C : G^\vee \to G^\vee$ are inner if and only if -1 is in the Weyl group, in which case, $C = \mathrm{int}(n_0)$, for some $n_0 \in N_{G^\vee}(T^\vee)$ representing the longest element $w_0 = -1$. In particular, Chevalley involutions are inner for types $A_1, B_n, C_n, D_{2k}, E_7, E_8, F_4$ and G_2, and outer for the other types.*

The *dual nonabelian Fourier transform* is a conjectural involution $\mathrm{FT}^\vee : R_{\mathrm{un}}(G) \to R_{\mathrm{un}}(G)$ that lifts Lusztig's non-abelian Fourier transform and is compatible with the parahoric restriction maps $\mathrm{res}^K_{\mathrm{un}}$. While such a map has not been constructed in the literature on the infinite dimensional space $R_{\mathrm{un}}(G)$, we consider the finite-dimensional quotient $\overline{R}_{\mathrm{un}}(G)$ of $R_{\mathrm{un}}(G)$ by the radical of the Euler–Poincare pairing

$$EP_G(\ ,\ ) : R_{\mathrm{un}}(G) \times R_{\mathrm{un}}(G) \longrightarrow \mathbb{C}, \quad (V, V') \longmapsto \sum_{i \geq 0} (-1)^i \mathrm{Ext}^i(V, V'), \text{ where } V, V' \in \mathrm{Irr}_{\mathrm{un}}(G),$$

and extended by linearity, and where $\mathrm{Ext}^i(V, V')$ are calculated in the category of smooth complex G-representations. Thus, the space $\overline{R}_{\mathrm{un}}(G)$ carries a natural non-degenerate pairing $EP_G(\ ,\ )$. Similarly, for each $x \in G^\vee$ the space of virtual representations $R(A_x)$ carries a natural elliptic pairing $(\ ,\ )_{\mathrm{ell}}^{A_x}$ (see [ACR25, §8.1] for the definition), and we let $\overline{R}(A_x)$ be the quotient of $R(A_x)$ by the radical of the elliptic pairing. The arithmetic-geometric (6) induces an isometric isomorphism on the elliptic representation spaces

$$\overline{\mathrm{LLC}}_{\mathrm{un}} : \bigoplus_{x \in G^\vee} \overline{R}(A_x) \longrightarrow \overline{R}_{\mathrm{un}}(G), \tag{7}$$

where the sum on the left is again over conjugacy classes of elements in $G^\vee$. The left hand side of (7) can be further decomposed into

$$\bigoplus_{x \in G^\vee} \overline{R}(A_x) = \bigoplus_{\substack{u \in G^\vee \\ \text{unipotent}}} \bigoplus_{\substack{s \in Z_{G^\vee}(u) \\ \text{semisimple}}} \overline{R}(A_{us}).$$

Next, fix some unipotent element $u \in G^\vee$ and let Γ_u be the reductive part of the centralizer $Z_{G^\vee}(u)$. We define the set of elliptic pairs associated to u to be

$$\mathcal{Y}(\Gamma_u)_{\mathrm{ell}} = \{(s, h) \in \Gamma_u \times \Gamma_u \mid s, h \text{ semisimple}, sh = hs, Z_{\Gamma_u}(s, h) \text{ is finite}\}.$$

It is not hard to show that this set is always finite (see [ACR25, Lemma 8.1]). For every $(s, h) \in \mathcal{Y}(\Gamma_u)$, define

$$\Pi(s, h) = \sum_{\phi \in \widehat{A_{\Gamma_u}(s)}} \phi(\overline{h})\phi \in R(A_{\Gamma_u}(s)), \tag{8}$$

where $\overline{h}$ is the image of h in $A_{\Gamma_u}(s)$. Moreover, let $\overline{\Pi}(s, h)$ denote the image in $\overline{R}(A_{\Gamma_u}(s))$ and let $\mathbb{C}[\mathcal{Y}(\Gamma_u)_{\mathrm{ell}}]^{\Gamma_u} \cong \mathbb{C}[\Gamma_u \backslash \mathcal{Y}(\Gamma_u)_{\mathrm{ell}}]$ be the space of Γ_u-invariant functions $\mathcal{Y}(\Gamma_u)$. Let $\mathbf{1}_{[(s,h)]}$ be the indicator function of the Γ_u-orbit of (s, h).

Proposition 3.10. [ACR25, Proposition 8.3] *Let $u \in G^\vee$ be unipotent. Then the correspondence $\mathbf{1}_{[(s,h)]} \mapsto \overline{\Pi}(s, h)$ induces an isomorphism*

$$\mathbb{C}[\mathcal{Y}(\Gamma_u)_{\mathrm{ell}}]^{\Gamma_u} \longrightarrow \bigoplus_{\substack{s \in Z_{G^\vee}(u) \\ \text{semisimple}}} \overline{R}(A_{us})$$

Finally, for each $[(s, h)] \in \Gamma_u \backslash \mathcal{Y}(\Gamma_u)_{\mathrm{ell}}$, define the virtual combinations

$$\Pi(u, s, h) := \sum_{\phi \in \widehat{A_{us}}} \phi(h)\pi(us, \phi) \in R_{\mathrm{un}}(G), \tag{9}$$

and let $\overline{\Pi}(u, s, h)$ be the corresponding images in $\overline{R}_{\mathrm{un}}(G)$.

Definition 3.11. The elliptic unipotent representation space $R_{\mathrm{un,ell}}(G)$ is the finite-dimensional subspace of $R_{\mathrm{un}}(G)$ spanned by virtual characters $\{\Pi(u, s, h) \mid u \text{ unipotent}, (s, h) \in \mathcal{Y}(\Gamma_u)_{\mathrm{ell}}\}$.

Thus, the isomorphism (7) together with Proposition 3.10 yields an isometric isomorphism

$$R_{\mathrm{un,ell}}(G) \longrightarrow \overline{R}_{\mathrm{un}}(G), \quad \Pi(u,s,h) \mapsto \overline{\Pi}(u,s,h) \tag{10}$$

We now define the dual elliptic nonabelian Fourier transform on $R_{\mathrm{un,ell}}(G)$.

Definition 3.12. The *dual elliptic nonabelian Fourier transform* is the involutive linear map $\mathrm{FT}^{\vee}_{\mathrm{ell}} : R_{\mathrm{un,ell}} \to R_{\mathrm{un,ell}}$ defined by

$$\mathrm{FT}^{\vee}_{\mathrm{ell}}(\Pi(u,s,h)) = \Pi(C(u), C(h), C(s)), \quad u \in G^{\vee} \text{ unipotent}, \ (s,h) \in \Gamma_u \backslash \mathcal{Y}(\Gamma_u)_{\mathrm{ell}},$$

where $C : G^{\vee} \to G^{\vee}$ is a Chevalley involution of the complex dual group.

We note that $\mathrm{FT}^{\vee}_{\ell}$ is a well-defined involution. A Chevalley involution induced an isomorphism $C : \Gamma_u \to \Gamma_{C(u)}$ and therefore $(C(h), C(s)) \in \Gamma_{C(u)} \backslash \mathcal{Y}(\Gamma_{C(u)})_{\mathrm{ell}}$. Moreover, any two Chevalley involutions differ by an inner automorphism, so $\mathrm{FT}^{\vee}_{\mathrm{ell}}$ is independent on the choice of Chevalley involution. Finally,

$$(\mathrm{FT}^{\vee}_{\mathrm{ell}})^2(\Pi(u,s,h)) = \Pi(C^2(u), C^2(s), C^2(h)) = \Pi(u,s,h).$$

Conjecture 3.13. [ACR25, Conjecture 1.2] Let G be a split semisimple p-adic group of simply connected type. Then the following diagram commutes up to roots of unity,

$$\begin{array}{ccc} R_{\mathrm{un,ell}}(G) & \xrightarrow{\mathrm{FT}^{\vee}_{\mathrm{ell}}} & R_{\mathrm{un,ell}}(G) \\ \downarrow{\scriptstyle \mathrm{res}^{\mathrm{cpt}}_{\mathrm{un}}} & & \downarrow{\scriptstyle \mathrm{res}^{\mathrm{cpt}}_{\mathrm{un}}} \\ \bigoplus_{K \in \max(G)} R_{\mathrm{un}}(\overline{K}) & \xrightarrow{\mathrm{FT}^{\mathrm{cpt}}_{\mathrm{un}}} & \bigoplus_{K \in \max(G)} R_{\mathrm{un}}(\overline{K}), \end{array} \tag{11}$$

where $\mathrm{FT}^{\mathrm{cpt}}_{\mathrm{un}} := \bigoplus_{K \in \max(G)} \mathrm{FT}^{\overline{K}}_{\mathrm{un}}$ on each coordinate.

Remark 3.14. Since Chevalley involutions are inner automorphisms for types $A_1, B_n, C_n, D_{2k}, E_7$, E_8, F_4 and G_2, the above conjecture remains the same statement as in [ACR25, Conjecture 9.7]. Thus, Conjecture 3.13 holds for $G = \mathrm{SO}_{2n+1}$ by the work of Moeglin–Waldspurger [MW03], for $G = G_2$ by the work of Ciubotaru–Opdam [CO17] and for $G = \mathrm{Sp}_4$ by the work of Aubert–Ciubotaru–Romano [ACR25].

In general, the main difficulty of the above conjecture relies on the computation of the unipotent parahoric restriction maps $\mathrm{res}^{K}_{\mathrm{un}}$. The space $R_{\mathrm{un}}(G)$ has a natural basis in terms of the geometry of the dual group $G^{\vee}$, while irreducible characters $R_{\mathrm{un}}(\overline{K})$ are parametrized by Lusztig's labels, often of combinatorial flavour. The connection between these parametrizations is deep, and we devote the next section to discuss this in detail for Iwahori-spherical representations. We also give a brief discussion on the computation of this map for non-Iwahori-spherical representations. We closely follow Reeder's discussion in [Ree00, §8,9].

3.3 Action of the fundamental group

Let $\mathcal{G}_{\mathrm{ad}}$ be the adjoint group of the simple algebraic F-group $\mathcal{G}$ split over F and let $G_{\mathrm{ad}} = \mathcal{G}_{ad}(F)$ be the adjoint group of F. The natural short exact sequence of algebraic groups

$$1 \longrightarrow Z(\mathcal{G}) \longrightarrow \mathcal{G} \xrightarrow{\eta} \mathcal{G}_{\mathrm{ad}} \longrightarrow 1$$

gives rise, via Galois cohomology, to an exact sequence

$$1 \longrightarrow Z(\mathcal{G})(F) \longrightarrow G \xrightarrow{\eta} G_{\mathrm{ad}} \xrightarrow{\delta} H^1(F, Z(\mathcal{G})) \longrightarrow H^1(F, \mathcal{G}) = 1,$$

by the Kneser–Bruhat–Tits theorem. In particular, $\eta : G \to G_{\mathrm{ad}}$ is not surjective, and $G_{\mathrm{ad}}/G := G_{\mathrm{ad}}/\eta(G)$ is isomorphic to the Galois cohomology group $H^1(F, Z(\mathcal{G}))$. Let $\mathcal{T}$ be an F-split maximal torus of $\mathcal{G}$ and let $\mathcal{T}_{\mathrm{ad}} := \mathcal{T}/Z(\mathcal{G})$ be an F-split maximal torus of $\mathcal{G}_{\mathrm{ad}}$, with $T = \mathcal{T}(F)$ and $T_{\mathrm{ad}} = \mathcal{T}_{\mathrm{ad}}(F)$. It is known [Spr98, Lemma 16.3.6] that the group $T_{\mathrm{ad}}/T := T_{\mathrm{ad}}/\eta(T)$ is naturally isomorphic to G_{ad}/G, and this happens to be a very useful characterization.

The standard adjoint action of G_{ad} on G induces a natural action on $\mathrm{Irr}_{\mathrm{un}}(G)$ given by pullback

$$(\mathrm{Ad}(g)^*\pi)(h) = (\pi \circ \mathrm{Ad}(g))(h) = \pi(ghg^{-1}) \text{ for } g \in G_{\mathrm{ad}} \text{ and } h \in G.$$

For any $g \in \eta(G)$, $\mathrm{Ad}(g)$ is an inner automorphism, so the action above factors through G_{ad}/G but is generally non-trivial since G_{ad}/G acts on G by outer automorphisms. In this subsection, we describe the effect of this action on the set of unipotent Langlands parameters of G explicitly. We closely follow Solleveld's results in [Sol20, §2.1] and use Kato's notation in [Kat83].

Let $x = su$ be the Jordan decomposition of $x \in G^\vee$ and let $M^\vee := Z_{G^\vee}(s)^0$ be a connected complex reductive group. We then consider the finite component group

$$R_s := Z_{G^\vee}(s)/M^\vee = \pi_0(Z_{G^\vee}(s)).$$

Using Galois cohomology, Solleveld constructs a group homomorphism

$$\tau_s : G_{\mathrm{ad}}/G \longrightarrow \mathrm{Hom}(R_s, \mathbb{C}^\times).$$

Next, we note that $Z_{G^\vee}(x) = Z_{Z_{G^\vee}(s)}(u)$ and therefore $A_x = \pi_0(Z_{Z_{G^\vee}(s)}(u))$. There is a natural map

$$Z_{G^\vee}(x) \hookrightarrow Z_{G^\vee}(s) \twoheadrightarrow R_s, \quad \text{descending to the map} \quad A_x \xrightarrow{\lambda_x} R_s,$$

whose kernel is $A_u^s := \pi_0(Z_{M^\vee}(u))$ and whose image is the subgroup

$$R_s(u) = \left\{ r \in R_s \mid r \cdot \mathcal{O}_u^{M^\vee} = \mathcal{O}_u^{M^\vee} \right\}.$$

We therefore obtain a (split) short exact sequence

$$1 \longrightarrow A_u^s \longrightarrow A_x \xrightarrow{\lambda_x} R_s(u) \longrightarrow 1,$$

and by composing with λ_x we obtain a map

$$\begin{aligned} \tau_x : G_{\mathrm{ad}}/G &\longrightarrow \mathrm{Hom}(A_x, \mathbb{C}^\times), \\ g &\longmapsto \tau_s(g) \circ \lambda_x. \end{aligned} \tag{12}$$

We note that, by construction, $\tau_x(g)|_{A_u^s} = 1$ for all $g \in G_{\mathrm{ad}}/G$. By [Sol20, Theorem 2.10], the action of G_{ad}/G on the Langlands parameters is given by

$$\mathrm{Ad}(g)^*\pi(x, \phi) = \pi(x, \phi \otimes \tau_x(g)^{-1}).$$

Next, we study in more detail the structure of the group G_{ad}/G. By the discussion above, it is naturally isomorphic to T_{ad}/T. Let $\nu_F : F^\times \to \mathbb{Z}$ be the normalizer valuation on F, and consider the canonical homomorphism

$$\begin{aligned} \nu_{T_{\mathrm{ad}}} : T_{\mathrm{ad}} &\longrightarrow \mathrm{Hom}(X^*(T_{\mathrm{ad}}), \mathbb{Z}) = X_*(T_{\mathrm{ad}}) = P^\vee, \\ t &\longmapsto (\chi \longmapsto \nu_F(\chi(t))), \end{aligned}$$

where $P^\vee$ is the coweight lattice. The fundamental group of G_{ad} is defined as $\Omega := P^\vee/Q^\vee$, where $Q^\vee = X_*(T)$ is the coroot lattice, and the map $\nu_{T_{\mathrm{ad}}}$ naturally induces the Kottowitz homomorphism

$$\kappa_T : T_{\mathrm{ad}}/T \twoheadrightarrow \Omega$$

fitting in the (split) short exact sequence

$$1 \longrightarrow T_{\mathrm{ad}}(\mathcal{O}_F)/\eta(T(\mathcal{O}_F)) \longrightarrow T_{\mathrm{ad}}/T \xrightarrow{\kappa_T} \Omega \longrightarrow 1.$$

The splitting on this short exact sequence is not canonical; it depends on a choice of uniformizer. Moreover, the action of any $t \in T_{\mathrm{ad}}(\mathcal{O}_F)$ on the Langlands parameters is trivial by [Sol20, Lemma 7.5], and therefore the map τ_x factors through the Kottowitz homomorphism $\kappa : G_{\mathrm{ad}}/G \twoheadrightarrow \Omega$, finally giving a map

$$\overline{\tau}_x : \Omega \longrightarrow \mathrm{Hom}(A_x, \mathbb{C}^\times)$$

such that $\mathrm{Ad}(\omega)^*\pi(x, \phi) = \pi(x, \phi \otimes \overline{\tau}_x(\omega)^{-1})$ for all $\omega \in \Omega$.

Lemma 3.15. *Let $u \in G^\vee$ unipotent and let $(s, h) \in \mathcal{Y}(\Gamma_u)_{\mathrm{ell}}$ be an elliptic pair. For every $\omega \in \Omega$,*

$$\mathrm{Ad}(\omega)^*\Pi(u, s, h) = \overline{\tau}_{us}(\omega)(h)\Pi(u, s, h).$$

Proof. This follows by direct computation; indeed

$$\begin{aligned}\mathrm{Ad}(\omega)^*\Pi(u,s,h) &= \sum_{\phi\in\widehat{A_{us}}} \phi(h)\mathrm{Ad}(w)^*\pi(us,\phi) = \\ &= \sum_{\phi\in\widehat{A_{us}}} \phi(h)\pi(us,\phi\otimes\overline{\tau}_{us}(\omega)^{-1}) = \\ &= \sum_{\psi\in\widehat{A_{us}}} (\psi\otimes\overline{\tau}_{us}(\omega))(h)\pi(us,\psi) = \overline{\tau}_{us}(\omega)(h)\Pi(u,s,h).\end{aligned}$$

□

In [Sol20, §2.1], Solleveld gives a general construction for the homomorphism $\overline{\tau}_x$ for each $x \in G^\vee$. In this setting of unramified Langlands parameters, we can give an explicit formulas that are very useful. Firstly, let $\eta^\vee : G^\vee_{\mathrm{sc}} \to G^\vee$ be the universal covering map of $G^\vee$. Fix some $u \in G^\vee$ unipotent and $(s,h) \in \mathcal{Y}(\Gamma_u)_{\mathrm{ell}}$. While u has a unique unipotent lift $\tilde{u} \in G^\vee_{\mathrm{sc}}$, the semisimple elements s and h do not. Nevertheless, for any lifts $\tilde{s}, \tilde{h} \in G^\vee_{\mathrm{sc}}$, the value of

$$c_s(h) := [\tilde{s},\tilde{h}] = \tilde{s}\tilde{h}\tilde{s}^{-1}\tilde{h}^{-1} \in Z(G^\vee_{\mathrm{sc}})$$

is independent of the choice of lifts. Finally, there is a natural identification

$$\Omega \longleftrightarrow X^*(Z(G^\vee_{\mathrm{sc}})) = \mathrm{Hom}(Z(G^\vee_{\mathrm{sc}}),\mathbb{C}^\times),$$

so each $\omega \in \Omega$ induces a character $\chi_\omega : Z(G^\vee_{\mathrm{sc}}) \to \mathbb{C}^\times$. Then, Solleveld constructs

$$\overline{\tau}_{us}(\omega)(h) = \chi_\omega(c_s(h)). \tag{13}$$

With this expression, we can prove an important identity.

Proposition 3.16. *Fix a unipotent element $u \in G^\vee$ and let $(s,h) \in \mathcal{Y}(\Gamma_u)_{\mathrm{ell}}$ be an elliptic pair. If $C : G^\vee \to G^\vee$ is a Chevalley involution, then for all $\omega \in \Omega$, the identity*

$$\overline{\tau}_{us}(\omega)(h) = \overline{\tau}_{C(uh)}(\omega)(C(s)) \tag{14}$$

holds.

Proof. □

Furthermore, the fundamental group Ω can be naturally embedded inside the automorphism group of the extended Dynkin diagram of G. By composing with inner automorphisms if necessary, we may assume that $\mathrm{Ad}(\omega)$ acts on G by permuting simple root spaces according to the automorphism of the Dynkin diagram it is associated with. In particular, if $J \subsetneq S_{\mathrm{aff}}$ and $\omega \in \Omega$, the map $\mathrm{Ad}(\omega)^*$ induces

an isomorphism $\overline{K}_J \xrightarrow{\sim} \overline{K}_{\omega(J)}$ and

$$\mathrm{res}_{\mathrm{un}}^{K_J} \mathrm{Ad}(\omega)^* \Pi(u,s,h) \cong \mathrm{res}_{\mathrm{un}}^{K_{\omega(J)}} \Pi(u,s,h). \tag{15}$$

Proposition 3.17. *Let G be a simple, split p-adic group of simply connected type. Suppose that the diagram* (11) *commutes for the parahoric subgroup K_J where $J \subsetneq S_{\mathrm{aff}}$. Then, for any $\omega \in \Omega$, the diagram also commutes for the parahoric $K_{\omega(J)}$.*

Proof. Again, this follows directly from Lemma 3.15, Proposition 3.16 and (15) since

$$\begin{aligned}
\mathrm{res}_{\mathrm{un}}^{K_{\omega(J)}} \circ \mathrm{FT}_{\mathrm{ell}}^{\vee} \Pi(u,s,h) &= \mathrm{res}_{\mathrm{un}}^{K_{\omega(J)}} \Pi(C(u), C(h), C(s)) = \\
&= \mathrm{res}_{\mathrm{un}}^{K_J} \mathrm{Ad}(\omega)^* \Pi(C(u), C(h), C(s)) = \\
&= \overline{\tau}_{C(uh)}(\omega)(C(s))\ \mathrm{res}_{\mathrm{un}}^{K_J} \circ \mathrm{FT}_{\mathrm{ell}}^{\vee} \Pi(u,s,h) = \\
&= \overline{\tau}_{us}(\omega)(h)\ \mathrm{FT}_{\mathrm{un}}^{K_J} \circ \mathrm{res}_{\mathrm{un}}^{K_J} \Pi(u,s,h) = \\
&= \mathrm{FT}_{\mathrm{un}}^{K_J} \circ \mathrm{res}_{\mathrm{un}}^{K_J} \mathrm{Ad}(\omega)^* \Pi(u,s,h) = \mathrm{FT}_{\mathrm{un}}^{K_{\omega(J)}} \circ \mathrm{res}_{\mathrm{un}}^{K_{\omega(J)}} \Pi(u,s,h).
\end{aligned}$$

□

3.4 The $\mathrm{SL}_n(F)$ case

In this section we prove that Conjecture 3.13 holds for $G = \mathrm{SL}_n(F)$. Since the dual group is $G^\vee = \mathrm{PGL}_n(\mathbb{C})$, via Jordan's normal form each unipotent conjugacy class corresponds to a partition of n $\lambda = (\underbrace{1,\dots,1}_{r_1},\underbrace{2,\dots,2}_{r_2},\dots,\underbrace{\ell,\dots,\ell}_{r_\ell})$, with $r_1 + 2r_2 + \cdots + \ell r_\ell = n$. Given $u \in \mathrm{PGL}_n(\mathbb{C})$ unipotent with corresponding partition λ_u of n, the reductive part Γ_u of the centralizer of u in $G^\vee$ contains elliptic pairs if and only if λ_u is rectangular; that is, if $\lambda_u = (\underbrace{m,\dots,m}_{d})$ for some $n = md$ (see [ACR25, Proposition 13.2]). In that case

$$\mathrm{PGL}_d(\mathbb{C}) \cong \Gamma_u, \quad g \mapsto g \otimes I_m.$$

Let $W = S_d$ be the Weyl group of $\mathrm{PGL}_d(\mathbb{C})$ (type A_{d-1}) and let $\dot{w}_d$ be the permutation matrix corresponding to the d-cycle $w_d = (1,2,3,\dots,d) \in S_d$. In addition, let $s_d = \mathrm{diag}(1,\zeta_d,\zeta_d^2,\dots,\zeta_d^{d-1}) \in \mathrm{PGL}_d(\mathbb{C})$ (note that s_d is a regular semisimple element). Then a standard group theory argument shows that

$$\mathcal{Y}(\Gamma_u)_{\mathrm{ell}} = \bigsqcup_{k \in (\mathbb{Z}/d\mathbb{Z})^\times} \Gamma_u \cdot (s_d, \dot{w}_d^k)$$

and moreover the pairs $(\dot{w}_d^k, s_d)$ and $(s_d, \dot{w}_d^{-k})$ are Γ_u-conjugate. In particular, the number of orbits of elliptic pairs in $G^\vee = \mathrm{PGL}_n(\mathbb{C})$ is

$$\sum_{u \text{ unipotent class}} |\Gamma_u \backslash \mathcal{Y}(\Gamma_u)_{\mathrm{ell}}| = \sum_{d|n} |(\mathbb{Z}/d\mathbb{Z})^\times| = \sum_{d|n} \varphi(d) = n.$$

For a detailed proof of these facts, see [ACR25, §13.1].

For the remainder of the subsection, fix some $d \mid n$, let $m = n/d$ and let $u_{d\times m} \in \mathrm{PGL}_n(\mathbb{C})$ be the standard unipotent element with Jordan blocks $\lambda_u = (\underbrace{m,\dots,m}_{d})$. Fix a primitive d-th root of unity ζ_d and let

$$s_{d\times m} = s_d \otimes I_m = \mathrm{diag}(\underbrace{1,\dots,1}_{m},\underbrace{\zeta_d,\dots,\zeta_d}_{m},\dots,\underbrace{\zeta_d^{d-1},\dots,\zeta_d^{d-1}}_{m}) \quad \text{and} \quad \dot{w}_{d\times m} = \dot{w}_d \otimes I_m.$$

By the discussion above, $(s_{d\times m}, \dot{w}_{d\times m}^k)$ is an elliptic pair for $k \in (\mathbb{Z}/d\mathbb{Z})^\times$, so we consider the virtual representations

$$\Pi(u_{d\times m}, s_{d\times m}, \dot{w}_{d\times m}^k) = \sum_{\ell \in \mathbb{Z}/d\mathbb{Z}} \zeta_d^{k\ell} \pi(x_{d\times m}, \phi_\ell),$$

where $x_{d\times m} = u_{d\times m} s_{d\times m}$ and $\phi_l(\dot{w}_{d\times m}) = \zeta_d^\ell$. Note that we have used that $A_{x_{d\times m}} = \langle \dot{w}_{d\times m} \rangle \cong \mathbb{Z}/d\mathbb{Z}$.

The standard Chevalley involution on $\mathrm{PGL}_n(\mathbb{C})$ is given by the inverse-transpose map $C(g) = (g^T)^{-1}$, and in particular $C(s_{d\times m}) = s_{d\times m}^{-1}$ and $C(w_{d\times m}) = w_{d\times m}$. Let J_m be the standard m by m unipotent

block in $PGL_m(\mathbb{C})$ and let $Y \in \mathrm{PGL}_m(\mathbb{C})$ be a matrix such that $Y(J_m^T)^{-1}Y^{-1} = J_m$. Then the matrix

$$X = \begin{pmatrix} Y & & & \\ & & & Y \\ & & \iddots & \\ & Y & & \end{pmatrix} \in \mathrm{PGL}_n(\mathbb{C})$$

satisfies that $XC(u_{d\times m})X^{-1} = u_{d\times m}$, $Xs_{d\times m}^{-1}X^{-1} = s_{d\times m}$ and $X\dot{w}_{d\times m}X^{-1} = \dot{w}_{d\times m}^{-1}$. Consequently,

$$\begin{aligned} \mathrm{FT}^{\vee}_{\mathrm{ell}}(\Pi(u_{d\times m}, s_{d\times m}, \dot{w}_{d\times m}^k)) &= \Pi(C(u_{d\times m}), C(\dot{w}_{d\times m}^k), C(s_{d\times m})) \\ &= \Pi(u_{d\times m}, \dot{w}_{d\times m}^{-k}, s_{d\times m}) = \Pi(u_{d\times m}, s_{d\times m}, \dot{w}_{d\times m}^k). \end{aligned} \tag{16}$$

Proposition 3.18. *Conjecture 3.13 holds for $G = \mathrm{SL}_n(F)$, without the appearance of roots of unity.*

Proof. According to (16), the dual nonabelian Fourier transform $\mathrm{FT}^{\vee}_{\mathrm{ell}}$ is the identity automorphism on $R_{\mathrm{un,ell}}(G)$. On the other hand, for each $K \in \max(G)$, the reductive quotient $\overline{K}$ is isomorphic to $\mathrm{SL}_n(\mathbb{F}_q)$, so $\mathrm{FT}^{\mathrm{cpt}}_{\mathrm{un}}$ is also the trivial automorphism on each $R_{\mathrm{un}}(\overline{K})$. The result follows. □

4 Parahoric restriction of unipotent representations

Let G be a connected split simple group of simply connected type over a p-adic field F. In the previous section we stated the bijection

$$\begin{aligned} \mathrm{LLC}^p_{\mathrm{un}} : G^\vee \backslash \{(x,\phi) \mid x \in G^\vee, \phi \in \widehat{A_x}\} &\longleftrightarrow \mathrm{Irr}_{\mathrm{un}}(G) \\ (x,\phi) &\longmapsto \pi(x,\phi), \end{aligned}$$

parametrizing unipotent representations of G in terms of the geometry of its complex dual group $G^\vee$. For a fixed $J \subsetneq S_{\mathrm{aff}}$ and corresponding parahoric subgroup K_J, in this section we discuss how to explicitly compute the parahoric restriction maps

$$\mathrm{res}^{K_J}_{\mathrm{un}} : R_{\mathrm{un}}(G) \longrightarrow R_{\mathrm{un}}(\overline{K}_J), \quad V \longmapsto V^{U_J}$$

from unipotent representations of G to unipotent representations of $\overline{K}_J$. We closely follow the discussion from [Ree00, §4-8], which we recall for convenience. However, Reeder assumes G to be of adjoint type, and not all results translate as cleanly when G is of simply connected type.

Let us fix throughout a unipotent representation $V = \pi(x,\phi)$ of G, with $x \in G^\vee$ and $\phi \in \widehat{A_x}$. For each irreducible unipotent $\overline{K}_J$-representation χ, we want to calculate the value of $\langle \chi, V^{U_J} \rangle_{K_J}$, where we view the $\overline{K}_J$-representations as K_J-representations trivial on U_J.

In the first part, we state two reduction results that simply this problem to a question about restrictions of representations over reflection groups associated to V and K_J. These groups are much smaller and easier to compute things with. This part does not require G to be of adjoint type. For example, if V is Iwahori-spherical, we obtain a representation $V^{\mathbf{1}}_{q=1}$ over the affine Weyl group $W_{\mathrm{aff}} = W \ltimes X(T^\vee)$ of G, and the parahoric restriction of V with respect to K_J can be determined by studying the representation $V^{\mathbf{1}}_{q=1}|_{\widetilde{W}_J}$ where $\widetilde{W}_J$ is the subgroup of W_{aff} generated by the reflections in J.

The action of W_{aff} on $V^{\mathbf{1}}_{q=1}$ can be explained via Springer theory. We devote the second part of the section to discuss this. However, since not every unipotent representation of G is Iwahori-spherical, we discuss in the end a more general method to calculate the parahoric restriction of non Iwahori-spherical representations with the use of weight diagrams developed in [Ree97]. These have been computed for all square-integrable unipotent representations of adjoint exceptional groups in [Ree00, §9].

4.1 Reduction to representations over reflection groups

Given the unipotent representation $V = \pi(x,\phi)$ of G, by Corollary 3.3 there is a unique proper subset I of S_{aff} with corresponding standard parahoric $K := K_I$, such that V^{U_K} contains a cuspidal unipotent

representation σ of $\overline{K}$. In particular, the vector space

$$V^{\sigma} := \mathrm{Hom}_K(\sigma, V^{U_K}) = \mathrm{Hom}_K(\sigma, V) = \mathrm{Hom}_G(c\text{-}\mathrm{Ind}_K^G \sigma, V) \tag{17}$$

is a non-zero irreducible $\mathcal{H}(G, K, \sigma) = \mathrm{End}_G(c\text{-}\mathrm{Ind}_K^G \sigma)$-module. The passage $V \mapsto V^{\sigma}$ is known to lose no information about the representation V ([BK98],[Lus95a]).

Similarly, if χ is an irreducible unipotent representation of $\overline{K}_J$, there is a unique proper subset I' of S_{aff} contained in J such that χ is a subrepresentation of $\mathrm{Ind}_{K_{I'}}^{K_J} \sigma'$, where σ' is a cuspidal unipotent representation of $\overline{K}_{I'}$. In particular, the vector space

$$\chi^{\sigma'} := \mathrm{Hom}_{K_{I'}}(\sigma', \chi) = \mathrm{Hom}_{K_J}(\mathrm{Ind}_{K_{I'}}^{K_J} \sigma', \chi) \neq 0$$

is naturally a $\mathcal{H}(K_J, K_{I'}, \sigma')$-module, where $\mathcal{H}(K_J, K_{I'}, \sigma') \cong \mathrm{End}_{K_J}(\mathrm{Ind}_{K_{I'}}^{K_J} \sigma')$ is the subalgebra of functions of $\mathcal{H}(G, K_{I'}, \sigma')$ supported on K_J.

Lemma 4.1 (First reduction)**.** *(*[Ree00, §4.2]*) The multiplicity of the simple K_J-module χ in V^{U_J} is given by*

$$\langle \chi, V^{U_J} \rangle_{K_J} = \langle \chi^{\sigma'}, V^{\sigma'} \rangle_{\mathcal{H}(K_J, K_{I'}, \sigma')}.$$

Moreover, $\langle \chi, V^{U_J} \rangle = 0$ unless $I = I'$. In particular, if J does not contain I, then $V^{U_J} = 0$.

Proof. Since $U_J \subseteq U_{I'}$, we have that

$$\begin{aligned} V^{\sigma'} = \mathrm{Hom}_{K_{I'}}(\sigma', V) = \mathrm{Hom}_{K_{I'}}(\sigma', V^{U_J}) &\simeq \mathrm{Hom}_{K_J}(\mathrm{Ind}_{K_{I'}}^{K_J} \sigma', V^{U_J}) \\ &\simeq \bigoplus_{\eta} \langle \eta, V^{U_J} \rangle_{K_J} \mathrm{Hom}_{K_J}(\mathrm{Ind}_{K_{I'}}^{K_J} \sigma', \eta) \\ &\simeq \bigoplus_{\eta} \langle \eta, V^{U_J} \rangle_{K_J} \eta^{\sigma'}, \end{aligned}$$

and therefore

$$\langle \chi^{\sigma'}, V^{\sigma'} \rangle_{\mathcal{H}(K_J, K_{I'}, \sigma')} = \sum_{\eta} \langle \chi^{\sigma'}, \eta^{\sigma'} \rangle_{\mathcal{H}(K_J, K_{I'}, \sigma')} \langle \eta, V^{U_J} \rangle_{K_J} = \langle \chi, V^{U_J} \rangle_{K_J}.$$

To prove the second statement, we note that $V^{\sigma'} \neq 0$ if and only if $(K, \sigma) = (K_{I'}, \sigma')$ by Corollary 3.3. In other words, $I = I'$ and the last assertion follows immediately from the fact that J contains I'. □

Example 4.2. Let us briefly explore the two extreme cases of the above result.

- If V is a Iwahori-spherical representation of G, then $V^{\mathcal{I}} \neq 0$, $I = \emptyset$ and the pair (K, σ) is given by $(\mathcal{I}, \mathbf{1})$. In this case, Lemma 4.1 states that

$$\langle \chi, V^{U_J} \rangle_{K_J} = \langle \chi^{\mathcal{I}}, V^{\mathcal{I}} \rangle_{\mathcal{H}(K_J, \mathcal{I}, \mathbf{1})}.$$

Thus, V has a non-zero parahoric restriction for any $J \subsetneq S_{\mathrm{aff}}$.

- If V is a supercuspidal unipotent representation of G, then I is a maximal proper subset of S_{aff} and $V^{U_J} = 0$ unless $J = I$.

For the second reduction step we consider $R = \mathbb{C}(v, v^{-1})$, where v is an indeterminate and define $\mathcal{H}(G, K, \sigma)_v$ to be the *generic Hecke algebra* defined over R with the same generators and relations as $\mathcal{H}(G, K, \sigma)$, but with q replaced by v^2 in the quadratic relations. By specializing this generic Hecke algebra we recover

$$\mathcal{H}(G, K, \sigma)_{\sqrt{q}} = \mathcal{H}(G, K, \sigma), \qquad \mathcal{H}(G, K, \sigma)_1 = \mathbb{C}[\tilde{W}(K, \sigma)],$$

where $\tilde{W}(K, \sigma)$ is an explicit affine reflection group associated to the pair (K, σ) (see [Lus95a] for an explicit list, and [Car85, §10] for a detailed construction). When $K = \mathcal{I}$ is the Iwahori subgroup and $\sigma = \mathbf{1}$ is the trivial representation, $\mathcal{H}(G, \mathcal{I}, \mathbf{1})_1 = \mathbb{C}[\widetilde{W}]$, where $\widetilde{W}$ is the affine Weyl group. Instead, if K is a maximal parahoric subgroup of G, then $\mathcal{H}(G, K, \sigma) \cong \mathbb{C}$ so $\tilde{W}(K, \sigma)$ is trivial.

Fact: For any simple $\mathcal{H}(G, K, \sigma)$-module E *considered in this document*, there is a $\mathcal{H}(G, K, \sigma)_v$-module E_v such that

$$E \simeq E_v \otimes_R \mathbb{C}, \text{ where } f \in R \text{ acts on } \mathbb{C} \text{ by } f(\sqrt{q}).$$

It then follows that we have a $q = 1$ operation that takes modules over Hecke algebras to modules over the corresponding reflection groups, obtained by setting $v = 1$ in all matrix coefficients of the generic module.

Proposition 4.3 (Second reduction)**.** *(*[Ree00, §6.1]*) Let $J \subsetneq S_{\mathrm{aff}}$ and let $(K, U_K, \overline{K})$ be a parahoric subgroup contained in $(K_J, U_J, \overline{K}_J)$ with cuspidal unipotent $\overline{K}$-representation σ. Then the diagram*

$$\begin{array}{ccc} \mathcal{H}(G, K, \sigma)\text{-}mod & \xrightarrow{q=1} & \tilde{W}(K, \sigma)\text{-}mod \\ \Big\downarrow{\scriptstyle \mathrm{Res}_{\mathcal{H}(K_J, K, \sigma)}} & & \Big\downarrow{\scriptstyle \mathrm{Res}_{\tilde{W}(K, \sigma)_J}} \\ \mathcal{H}(K_J, K, \sigma)\text{-}mod & \xrightarrow{q=1} & \tilde{W}(K, \sigma)_J\text{-}mod \end{array}$$

is commutative, and the bottom arrow is an isometry with respect to the usual inner product of character. That is, for any irreducible K_J-module χ, we have that

$$\langle \chi^\sigma, V^\sigma \rangle_{\mathcal{H}(K_J, K, \sigma)} = \langle \chi^\sigma_{q=1}, V^\sigma_{q=1} \rangle_{\tilde{W}(K, \sigma)_J}. \tag{18}$$

Combining both reduction steps, we obtain the identity

$$\langle \chi, V^{U_J} \rangle_{K_J} = \langle \chi^\sigma, V^\sigma \rangle_{\mathcal{H}(K_J, K, \sigma)} = \langle \chi^\sigma_{q=1}, V^\sigma_{q=1} \rangle_{\tilde{W}(K, \sigma)_J} \tag{19}$$

Thus, to compute the parahoric restriction V^{U_J}, it suffices to study the $\tilde{W}(K, \sigma)_J$-module $V^\sigma_{q=1}$. By

calculating its decomposition into $\tilde{W}(K,\sigma)$-irreducible representation, one can directly obtain the decomposition of V^{U_J} into irreducible $\overline{K}_J$-representations.

At this point, we restrict our attention to the parahoric restriction of Iwahori-spherical representations, where $\tilde{W}(\mathcal{I},\mathbf{1}) \cong W_{\text{aff}}$ and $\tilde{W}(\mathcal{I},\mathbf{1})_J = \widetilde{W}_J$ is the subgroup of $\widetilde{W}$ generated by the simple reflections of the roots in J.

4.2 The Iwahori-spherical case and the Springer correspondence

Given an Iwahori-spherical representation $V = \pi(x,\phi)$ of G and a proper subset $J \subsetneq S_{\text{aff}}$, in the previous subsection we proved that

$$\langle \chi, V^{U_J} \rangle_{\overline{K}_J} = \langle \chi^{\mathbf{1}}_{q=1}, V^{\mathbf{1}}_{q=1} \rangle_{\widetilde{W}_J}. \tag{20}$$

for any principal series representation χ of the reductive quotient $\overline{K}_J$. We use Springer theory to extract the structure of the $\widetilde{W}_J$-module $V^{\mathbf{1}}_{q=1}$ in terms of the data (x,ϕ). Let $x = us$ be the Jordan decomposition of x, with $u \in G^\vee$ unipotent and $s \in G^\vee$ semisimple. By conjugating if necessary, we assume that $s \in T^\vee$ for a fixed split maximal torus of $G^\vee$. Let $\mathcal{B}$ and $\mathcal{B}_s$ be the flag varieties of the complex reductive groups $G^\vee$ and $M^\vee := Z_{G^\vee}(s)^0$, respectively, and consider the Springer fibres $\mathcal{B}^x$ and $\mathcal{B}^u_s$ parametrizing the set of Borel subgroups of $G^\vee$ containing x and the set of Borel subgroups of $M^\vee$ containing u. Let W_M be the Weyl group of $M^\vee$, isomorphic to the subgroup of W generated by all reflections of roots trivial on s.

While the varieties $\mathcal{B}_x$ and $\mathcal{B}^u_s$ do not admit a natural action of W and W_M, respectively, their singular cohomology spaces $H^*(\mathcal{B}_x)$ and $H^*(\mathcal{B}^u_s)$ do. The W_M structure of $H^*(\mathcal{B}^u_s)$ can be easily read from the Springer correspondence for the reductive group $M^\vee$ via the Green tables, which are well understood. Moreover, the module structures are tightly related by the formula ([Kat83, Proposition 2.5]),

$$H^*(\mathcal{B}^x) \cong \text{Ind}_{W_M}^{W} H^*(\mathcal{B}^u_s).$$

This action can be naturally extended to an action of the affine Weyl group $W_{\text{aff}} = W \ltimes X(T^\vee)$, where $X = X(T^\vee)$ is the character lattice of $T^\vee$. The semisimple element $s \in T^\vee$ is naturally a character of X given by evaluation. Thus, the cohomology spaces $H^*(\mathcal{B}^x)$ carry a natural W_{aff}-action given by

$$H^*(\mathcal{B}^x) = \text{Ind}_{W_{M,\text{aff}}}^{W_{\text{aff}}} H^*(\mathcal{B}^u_s) \otimes s,$$

where $W_{M,\text{aff}} = W_M \ltimes X$ (see [Kat83, Proposition 3.2]).

Finally, the component group A_x acts naturally on the varieties $\mathcal{B}^x$ and $\mathcal{B}^u_s$ by conjugation, inducing a natural action on the cohomology spaces $H^*(\mathcal{B}^x)$ and $H^*(\mathcal{B}^u_s)$. While the A_x-action on $H^*(\mathcal{B}^x)$

commutes with the W_{aff}-action, thus giving an $A_x \times W_{\mathrm{aff}}$-module structure, the A_x-action on $H^*(\mathcal{B}_s^u)$ does not necessarily commute with the $W_{M,\mathrm{aff}}$-action. The precise structure of this action is subtle, and we closely follow the discussion in [Kat83, §6] to explain this.

Given $x = su$ as above, let W_s be the stabilizer of s in W. Then W_M is a normal subgroup of W_s, and it fits in the splitting short exact sequence

$$1 \longrightarrow W_M \longrightarrow W_s \longrightarrow R_s \longrightarrow 1,$$

where $R_s = Z_{G^\vee}(s)/M^\vee$ is the component group of the centralizer of s in $G^\vee$ and the splitting is given by the identification $R_s \cong \{w \in W_s \mid w(\Phi_M^+) = \Phi_M^+\}$, where Φ_M^+ are the positive roots of $M^\vee$. In particular, $W_s = W_M \rtimes R_s$ and, by choosing Kac coordinates for s, the group R_s can be seen as a subgroup of the fundamental group Ω. To introduce the role of the unipotent element $u \in G^\vee$, Kato defines the canonical subgroup

$$R_s(u) = \{r \in R_s \mid r \cdot \mathcal{O}_u^{M^\vee} = \mathcal{O}_u^{M^\vee}\} \cong \{w \in R_s \mid u^{n(w)} \text{ and } u \text{ are conjugate in } M^\vee\}.$$

Since $Z_{G^\vee}(x) = Z_{G^\vee}(s) \cap Z_{G^\vee}(u)$, we have that $Z_{M^\vee}(u) \triangleleft Z_{G^\vee}(x)$ and $Z_{G^\vee}(x)/Z_{M^\vee}(u) \cong R_s(u)$. In particular, we may regard $A_u^s := Z_{M^\vee}(u)/Z_{M^\vee}(u)^0$ as a normal subgroup of A_x, and we have the isomorphism $A_x/A_u^s \cong R_s(u)$.

Denote by $j : Z_{G^\vee}(s) \to R_s$ the canonical surjection and note that each $y \in Z_{G^\vee}(x) \leq Z_{G^\vee}(s)$ defines a map $H^i(\mathcal{B}_s^u) \to H^i(\mathcal{B}_s^u)$, which we also denote by y. Moreover, for each $w \in W_{M,\mathrm{aff}}$, it follows from the definitions of the $W_{M,\mathrm{aff}}$ that [Kat83, (6.1)]

$$y \cdot w = {}^{j(y)}w \cdot y : H^i(\mathcal{B}_s^u) \longrightarrow H^i(\mathcal{B}_s^u).$$

With this in mind, we consider the induced surjection $j_0 : A_x \to R_s(u)$ and, since $R_s(u)$ acts on $W_{M,\mathrm{aff}}$ by conjugation, we can construct the semidirect product $A_x \ltimes W_{M,\mathrm{aff}}$. This group embeds in $A_x \times W_{\mathrm{aff}}$ via the map $(a, w) \mapsto (a, j_0(a)w)$ (the image in fact lands inside $A_x \times W_{s,\mathrm{aff}}$), and Kato shows that the A_x-action on $H^*(\mathcal{B}_s^u)$ extends to an $A_x \ltimes W_{M,\mathrm{aff}}$-action. In particular, we have the following result.

Proposition 4.4 ([Kat83, Proposition 6.2])**.** *As $A_x \times W_{\mathrm{aff}}$-modules, we have*

$$H^*(\mathcal{B}^x) \cong \mathrm{Ind}_{A_x \ltimes W_{M,\mathrm{aff}}}^{A_x \times W_{\mathrm{aff}}} H^*(\mathcal{B}_s^u).$$

Let us now go back to representations of p-adic groups. Let $V_{(x,\phi)}$ be an Iwahori-spherical unipotent representation of G. Then $V_{(x,\phi)}^{\mathcal{I}}$ is a simple $\mathcal{H}(G, \mathcal{I}, \mathbf{1})$-module, while $V_{(x,\phi),q=1}^{\mathbf{1}}$ is a simple W_{aff}-module. We then have the following central result, due to Lusztig. We remark that up to this point we have not used the fact that G is of simply connected type, so this result holds more generally by swapping W_{aff} with an adequate subgroup of the extended affine Weyl group $\widetilde{W}$.

Theorem 4.5. [Lus95b] *Let $V_{x,\phi}$ be an Iwahori-spherical irreducible representation of G. After letting $q \to 1$ and then taking semisimplification, the simple $\mathcal{H}(G, I, \mathbf{1})$-module $V^{\mathbf{1}}_{x,\phi}$ becomes the W_{aff}-module $\varepsilon \otimes H^*(\mathcal{B}^x)^\phi$, where ε is the sign representation of W_{aff}.*

Remark 4.6. An important remark is due at this point. The above theorem relates the structure of $V^{\mathbf{1}}_{(x,\phi),q=1}$ with the structure of $H^*(\mathcal{B}^x)^\phi$ as W_{aff}-modules. However, the former object is naturally p-adic, while the latter is complex analytic. The Weyl groups of G and $G^\vee$ are canonically isomorphic, but not equal if the group is not simply laced, since the isomorphism swaps short with long roots. It is a standard abuse of notation to denote both Weyl groups with the same symbol, but one needs to be careful during explicit calculations inside which Weyl group one is working. We are ultimately interested in the p-adic side, but most computations involving cohomology groups and the Springer correspondence take place in the complex side. In particular, the reflection s_0 inside W will be a **short** reflection, corresponding to the highest short root. At the very end, we then trace the representations back to the p-adic side by swapping long and short roots.

Let us discuss some consequences of Theorem 4.5. When $R(u) = \{1\}$, the semidirect product $A_x \ltimes W_{M,\mathrm{aff}}$ is simply the direct product $A_x \times W_{M,\mathrm{aff}}$, and we have that

$$H^*(\mathcal{B}^x)^\phi = \left(\mathrm{Ind}_{A_x \times W_{M,\mathrm{aff}}}^{A_x \times W_{\mathrm{aff}}} H^*(\mathcal{B}^u_s)\right)^\phi = \mathrm{Ind}_{W_{M,\mathrm{aff}}}^{W_{\mathrm{aff}}} H^*(\mathcal{B}^u_s)^\phi. \tag{21}$$

From (20) and Theorem 4.5, the computation of the parahoric restriction of $V_{(x,\phi)}$ is then reduced to the computation of the $\widetilde{W}_J$-module structure of $H^*(\mathcal{B}^x)^\phi$. The difficulty of this case comes from simultaneously controlling the action of the character lattice X on $H^*(\mathcal{B}^u_s)^\phi$. In [Ree00], Reeder observes that the canonical projection from W_{aff} to W is injective on each $\widetilde{W}_J$ with image W_J. If $\psi_J : W_J \to \widetilde{W}_J$ is the inverse of this map, then we can consider the pullback ψ_J^* of representations of $\widetilde{W}_J$ to representations of W_J. By (21) and Mackey theory, we have that

$$\psi_J^*\left[\left(\mathrm{Ind}_{\widetilde{W}_M}^{\widetilde{W}} H^*(\mathcal{B}^u_s)^\phi\right)|_{\widetilde{W}_J}\right] \cong \bigoplus_{w \in W_M \backslash W / W_J} \mathrm{Ind}_{W_{J,s^w}}^{W_J} \chi^J_{s^w} \otimes \left[H^*(\mathcal{B}^u_s)^\phi\right]^w, \tag{22}$$

where $W_{J,s^w} = W_M^w \cap W_J$ and $\chi^J_{s^w} = \chi_{s^w} \circ \chi_J : W_{J,s^w} \to \mathbb{C}^\times$. To use this formula we need an understanding of the groups W_{J,s^w} and the characters $\chi^J_{s^w}$. The groups $W_{J,t}$ have been studied in the literature, and they can be decomposed into

$$W_{J,t} = W^0_{J,t} \rtimes R_{J,t},$$

where $W^0_{J,t}$ is a "large" reflection subgroup of W generated by the reflections in J that fix t, and $R_{J,t}$ is a "small" group that normalizes $W^0_{J,t}$. The character χ^J_t is trivial on $W^0_{J,t}$ and is determined by its restriction to $R_{J,t}$. As a result, the characters χ^J_t are trivial in many cases. We refer to [Ree00, §8] for a detailed discussion on the structure of these characters, where explicit formulas to compute them are provided. If all characters $\chi^J_{s^w}$ appearing in (22) are trivial, then computations are greatly simplified.

Lemma 4.7. *With the same notation as in* (22)*, if the characters* $\chi_{s^w}^J$ *are trivial for all* $w \in W_s\backslash W/W_J$*, then*

$$\psi_J^*\left[\left(\mathrm{Ind}_{\widetilde{W}_s}^{\widetilde{W}} H^*(\mathcal{B}_s^u)^\phi\right)|_{\widetilde{W}_J}\right] \cong \left(\mathrm{Ind}_{W_s}^{W} H^*(\mathcal{B}_s^u)^\phi\right)|_{W_J}$$

Proof. This is an immediate consequence of Mackey's formula applied to the right hand side of (22). □

4.3 Non-Iwahori-spherical case and weight diagrams

Let us now fix a general unipotent representation $V = V_{(x,\phi)}$ of G with associated parahoric $K := K_I$ and cuspidal unipotent representation σ as in (17). Then V^σ is a simple module over the Hecke algebra $\mathcal{H}(G, K, \sigma) = \mathrm{End}_G(c\text{-}\mathrm{Ind}_K^G \sigma)$. The structure of these Hecke algebras have been widely studied in the literature (see [Lus95a, §1] for a detailed discussion). In particular, one can associate a canonical extended affine Weyl group $\tilde{W}(K, \sigma)$ parametrizing a basis $\{T_w, w \in \tilde{W}(K, \sigma)\}$ of the algebra $\mathcal{H}(G, K, \sigma)$ satisfying explicit relations. Importantly, the basis elements $\{T_s \mid s \in S_{\mathrm{aff}}(K, \sigma)\}$ for some choice of simple reflections $S_{\mathrm{aff}}(K, \sigma)$ in $\tilde{W}(K, \sigma)$ satisfy the quadratic relations

$$(T_s + 1)(T_s - q^{c(s)}) = 0,$$

where $c(s)$ is an explicit positive integer depending on the simple root corresponding to s. The structure of a modules over $\mathcal{H}(G, K, \sigma)$ can then be largely understood by studying the action of the basis elements T_s corresponding to the simple reflections.

On the other hand, the extended affine Weyl group $\tilde{W}(K, \sigma)$ can be decomposed as a semidirect product $W(K, \sigma) \ltimes X(K, \sigma)$, where $W(K, \sigma)$ can be realized as the Weyl group of an associated complex reductive group $\mathbf{G}$ and $X(K, \sigma)$ as the character lattice of a maximal torus $\mathbf{T}$ of $\mathbf{G}$. The subalgebra spanned by $\{T_w, w \in W(K, \sigma)\}$ is naturally isomorphic to the subalgebra $\mathcal{H}(K_{J_0}, K, \sigma)$ of functions supported on a hyperspecial maximal compact subgroup K_{J_0} of G. We then have an explicit isomorphism ([Sol21, Theorem 1.10], [Lus89, §3])

$$\mathcal{H}(G, K, \sigma) \cong \mathcal{H}(K_{J_0}, K, \sigma) \,\tilde{\otimes}\, \mathcal{O}[\mathbf{T}], \tag{23}$$

where $\tilde{\otimes}$ is a twisted tensor product of $\mathbb{C}$-vector space with a twisted multiplication given by

$$(T_s - q^{c(s)})f - f^s(T_s - q^{c(s)}) = (f^s - f)\zeta_s \in \mathcal{O}[\mathbf{T}]$$

where $\zeta_s \in (f^s - f)^{-1}\mathcal{O}[\mathbf{T}]$ is a rational function on $\mathbf{T}$.

Example 4.8. Consider the case of $G = F_4(F)$, K a parahoric subgroup of type B_2 and σ the unique cuspidal unipotent representation of $B_2(\mathbb{F}_q)$. Then $\tilde{W}(K, \sigma)$ is the affine Weyl group of type C_2 and the

Hecke algebra $\mathcal{H}(G, K, \sigma)$ is generated by the elements $T_{s_0}, T_{s_1}, T_{s_2}$ satisfying the quadratic relations

$$(T_{s_0} + 1)(T_{s_0} - q) = 0, \quad (T_{s_i} + 1)(T_{s_i} - q^3) = 0, \quad \text{for } i = 1, 2.$$

Moreover, $\mathbf{G} = \mathrm{SO}_5(\mathbb{C})$ and $\mathbf{T} = (\mathbb{C}^\times)^2$, with simple roots

$$e_{\alpha_1}(t_1, t_2) = t_1/t_2, \quad e_{\alpha_2}(t_1, t_2) = t_2,$$
$$\zeta_{s_1} = \frac{q^3 - e_{\alpha_1}}{1 - e_{\alpha_1}}, \quad \zeta_{s_2} = \frac{(q^2 - e_{\alpha_2})(q + e_{\alpha_2})}{1 - e_{2\alpha_2}}.$$


The theory of weight diagrams developed by Reeder in [Ree97] studies first the restriction of the module V^σ to the large subalgebra $\mathcal{O}[\mathbf{T}]$ and then the action of the basis elements T_s for $s \in W(K, \sigma)$ a simple reflection on the resulting decomposition. Since $\mathcal{O}[\mathbf{T}]$ is a commutative algebra, there is a decomposition

$$V^\sigma|_{\mathcal{O}[\mathbf{T}]} = \bigoplus_{\tau \in \mathbf{T}} V^\sigma(\tau),$$

where $V^\sigma(\tau)$ consists of those vectors in V^σ killed by some power of the maximal ideal of $\mathcal{O}[\mathbf{T}]$ corresponding to τ. From this decomposition, we can determine the action of the basis elements $W(K, \sigma)$ on the space $V^\sigma_{q=1}$, which immediately gives the parahoric restriction of V with respect to the maximal hyperspecial parahoric subgroup K_{J_0}. See [Ree00, §9.5] for the explicit formulas. To obtain the parahoric restriction with respect to other parahoric subgroups, one needs to understand the action of the remaining simple reflection $s_0 \in \tilde{W}(K, \sigma)$. This can be done by tracing back through the isomorphism $\mathcal{H}(G, K, \sigma) \cong \mathcal{H}(K_{J_0}, K, \sigma) \tilde{\otimes} \mathcal{O}[\mathbf{T}]$ and using the explicit formula for the twisted multiplication. See Example 4.8 for an explicit calculation. The remaining cases are omitted and we provide the final results without proof.

4.4 Parahoric restriction for type G_2

We end this section with the explicit calculation of the parahoric restrictions for some representations $\pi(x,\phi)$ of the p-adic group $G = G_2(F)$, highlighting the methods described above with a concrete example. This group is both simply connected and adjoint, with dual group $G^\vee = G_2(\mathbb{C})$. Let α_1, α_2 be a choice of simple roots with α_1 short and let $\alpha_0 = 2\alpha_1 + \alpha_2,\ \alpha^0 = 3\alpha_1 + 2\alpha_2$ be the highest short and long root, respectively. The affine Dynkin diagram of $G_2(F)$ is



Thus $G_2(F)$ has 3 classes of maximal parahoric subgroups K_0, K_1, K_2 corresponding to the subsets of affine simple roots $J_0 = \{\alpha_1^\vee, \alpha_2^\vee\}, J_1 = \{-\alpha_0^\vee, \alpha_2^\vee\}$ and $J_2 = \{-\alpha_0^\vee, \alpha_1^\vee\}$ and with reductive types G_2, $A_1 \times \tilde{A}_1$ and A_2, respectively. We now compute these parahoric restrictions for all representations $\pi(us,\phi)$ of $G = G_2(F)$, where u is a subregular unipotent element of $G_2(\mathbb{C})$, labelled by $G_2(a_1)$. This is an interesting case since u is a distinguished unipotent element and $\Gamma_u = A_u \cong S_3$. There are 8 such representations, given by

$$\pi(G_2(a_1), 1, \phi), \phi \in \{\mathbf{1}, \varepsilon, r\}, \quad \pi(G_2(a_1), g_2, \phi), \phi \in \{\mathbf{1}, \varepsilon\}, \quad \pi(G_2(a_1), g_3, \phi), \phi \in \{\mathbf{1}, \theta, \theta^2\}.$$

Lusztig's arithmetic-geometric correspondence directly implies that 4 of them are unipotent supercuspidal representations of $G_2(F)$ given by

$$\pi(G_2(a_1), 1, \epsilon) = c\text{-Ind}_{K_0}^G G_2[1], \quad \pi(G_2(a_1), g_2, \epsilon) = c\text{-Ind}_{K_0}^G G_2[-1] \quad \text{and}$$
$$\pi(G_2(a_1), g_3, \theta^k) = c\text{-Ind}_{K_0}^G G_2[\theta^k], k = 1, 2,$$

while the remaining 4 are Iwahori-spherical. For the supercuspidal representations, their restrictions can be easily computed. By Frobenius reciprocity, their parahoric restrictions to K_0 are irreducible, while the remaining parahoric restrictions are 0.

To compute parahoric restriction of Iwahori-spherical representations, we recall that we work inside *complex* Weyl group W and consequently s_0 is as a short reflection along α_0, while s^0 is the long reflection along α^0. The unipotent element $u \in G^\vee = G_2(\mathbb{C})$ is subregular, so by Springer theory

$$H^*(\mathcal{B}^u)^{\mathbf{1}} = H^0(\mathcal{B}^u)^{\mathbf{1}} + H^2(\mathcal{B}^u)^{\mathbf{1}} = \phi_{1,0} + \phi_{2,1}, \quad \text{and} \quad H^*(\mathcal{B}^u)^r = H^0(\mathcal{B}^u)^r + H^2(\mathcal{B}^u)^r = \phi'_{1,3}$$

as W-modules. On the other hand, u is a regular unipotent element in $G_{g_2}^\vee = \mathrm{GL}_2(\mathbb{C})$ and $G_{g_3}^\vee = \mathrm{SL}_3(\mathbb{C})$, so $H^0(\mathcal{B}_{g_2}^u)$ and $H^0(\mathcal{B}_{g_3}^u)$ afford the trivial representations of $W_{g_2} = \langle s^0, s_1\rangle$ and $W_{g_3} = \langle s^0, s_2\rangle$ respectively, and

$$\mathrm{Ind}_{W_{g_2}}^W \mathbf{1} = \phi_{1,0} + \phi''_{1,3} \quad \text{and} \quad \mathrm{Ind}_{W_{g_3}}^W \mathbf{1} = \phi_{1,0} + \phi_{2,2}.$$

To obtain the parahoric restrictions to K_0, we note that all characters $\chi_t^{J_0}$ are trivial so it remains to twist by $\varepsilon = \phi_{1,6}$ (swapping $\phi_{1,0} \leftrightarrow \phi_{1,6}$ and $\phi'_{1,3} \leftrightarrow \phi''_{1,3}$) and then by the outer automorphism of W exchanging short and long roots (swapping $\phi'_{1,3} \leftrightarrow \phi''_{1,3}$ only).

To compute the parahoric restrictions with respect to K_1 and K_2 in Table 2 we compute the remaining characters χ_t^J. Using their properties above, one can check that all characters are trivial except for the pairs $(s, J) = (g_3, J_2)$ and $(s, J) = (g_2, J_1)$. When the characters are trivial, Lemma 4.7 and Alvis' tables immediately yield the results. The last two cases must be dealt with separately.

- **Restriction of $\pi(G_2(a_1), g_3, 1)$ to K_{J_2}.** In this case, $W_{J_2} = \langle s_0, s_1 \rangle$ (where s_0 is the reflection along the highest short root) and $W_{g_3} = \langle s^0, s_2 \rangle$ (where s^0 is the reflection along the highest root). Then $W_{g_3} \backslash W / W_{J_2} = \{1\}$, $W_{J_2, g_3} \cong C_3$ and $\chi_{g_3}^{J_2}$ is a primitive character. Thus,

$$\mathrm{Ind}_{W_{J_2, g_3}}^{W_{J_2}} \chi_{g_3}^{J_2} = r.$$

- **Restriction of $\pi(G_2(a_1), g_2, 1)$ to K_{J_1}.** In this case, $W_{J_1} = \langle s_0, s_2 \rangle$ and $W_{g_2} = \langle s^0, s_1 \rangle$, so $W_{g_2} \backslash W / W_{J_1} = \{1, w\}$ has two elements. Then, $W_{J_1, g_2} \cong C_2$ with $\chi_{g_2}^{J_1}$ non-trivial character and $W_{J_1, g_2^w} = W_{J_1}$ with $\chi_{g_2^w}^{J_1}$ the trivial character. Thus,

$$\mathrm{Ind}_{W_{J_1, g_2}}^{W_{J_1}} \chi_{g_2}^{J_1} + \mathrm{Ind}_{W_{J_1, g_2^w}}^{W_{J_1}} \chi_{g_2^w}^{J_1} = \varepsilon \boxtimes \mathbf{1} + \mathbf{1} \boxtimes \varepsilon + \mathbf{1} \boxtimes \mathbf{1}.$$

Twisting by the sign character yields the desired result. The proof of Conjecture 1.1 is now just a matter of putting all the pieces together. The parahoric restrictions of the virtual characters $\Pi(u, s, h)$ can be computed by linearity from the parahoric restrictions of the representations $\pi(u, s, \phi)$, which are given in Table 2.

	$K_0 \longrightarrow G_2$	$K_1 \longrightarrow A_1 \times \tilde{A}_1$	$K_2 \longrightarrow A_2$
$\Pi(G_2, 1, 1)$	$\phi_{1,6}$	$\epsilon \boxtimes \epsilon$	ϵ
$\Pi(G_2(a_1), 1, 1)$	$\phi_{2,1} + G_2[1] + 2\phi'_{1,3} + \phi_{1,6}$	$\mathbf{1} \boxtimes \epsilon + 3\epsilon \boxtimes \mathbf{1} + \epsilon \boxtimes \epsilon$	$3\epsilon + r$
$\Pi(G_2(a_1), 1, g_2)$	$\phi_{2,1} - G_2[1] + \phi_{1,6}$	$\mathbf{1} \boxtimes \epsilon + \epsilon \boxtimes \mathbf{1} + \epsilon \boxtimes \epsilon$	$\epsilon + r$
$\Pi(G_2(a_1), g_2, 1)$	$\phi_{2,1} + \phi_{2,2} + G_2[-1] + \phi_{1,6}$	$\mathbf{1} \boxtimes \epsilon + \epsilon \boxtimes \mathbf{1} + \epsilon \boxtimes \epsilon$	$\epsilon + r$
$\Pi(G_2(a_1), 1, g_3)$	$\phi_{2,1} + G_2[1] - \phi'_{1,3} + \phi_{1,6}$	$\mathbf{1} \boxtimes \epsilon + \epsilon \boxtimes \epsilon$	r
$\Pi(G_2(a_1), g_3, 1)$	$\phi''_{1,3} + G_2[\theta] + G_2[\theta^2] + \phi_{1,6}$	$\mathbf{1} \boxtimes \epsilon + \epsilon \boxtimes \epsilon$	r
$\Pi(G_2(a_1), g_2, g_2)$	$\phi_{2,1} + \phi_{2,2} - G_2[-1] + \phi_{1,6}$	$\mathbf{1} \boxtimes \epsilon + \epsilon \boxtimes \mathbf{1} + \epsilon \boxtimes \epsilon$	$\epsilon + r$
$\Pi(G_2(a_1), g_3, g_3)$	$\phi''_{1,3} + \theta G_2[\theta] + \theta^2 G_2[\theta^2] + \phi_{1,6}$	$\mathbf{1} \boxtimes \epsilon + \epsilon \boxtimes \epsilon$	r
$\Pi(G_2(a_1), g_3, g_3^{-1})$	$\phi''_{1,3} + \theta^2 G_2[\theta] + \theta G_2[\theta^2] + \phi_{1,6}$	$\mathbf{1} \boxtimes \epsilon + \epsilon \boxtimes \epsilon$	r

Table 2: Restrictions of G_2-representations $\pi(G_2(a_1), s, \phi)$

Finally, Lusztig's nonabelian Fourier transform acts trivially on every representation, except for the $G_2(\mathbb{F}_q)$ family

$$\{\phi_{2,1}, \phi'_{1,3}, G_2[1], \phi_{2,2}, G_2[-1], \phi''_{1,3}, G_2[\theta], G_2[\theta^2]\},$$


where the action is given by the 8×8 matrix

$$\begin{pmatrix}
\frac{1}{6} & \frac{1}{3} & \frac{1}{6} & \frac{1}{2} & \frac{1}{2} & \frac{1}{3} & \frac{1}{3} & \frac{1}{3} \\
\frac{1}{3} & \frac{2}{3} & \frac{1}{3} & 0 & 0 & -\frac{1}{3} & -\frac{1}{3} & -\frac{1}{3} \\
\frac{1}{6} & \frac{1}{3} & \frac{1}{6} & -\frac{1}{2} & -\frac{1}{2} & \frac{1}{3} & \frac{1}{3} & \frac{1}{3} \\
\frac{1}{2} & 0 & -\frac{1}{2} & \frac{1}{2} & -\frac{1}{2} & 0 & 0 & 0 \\
\frac{1}{2} & 0 & -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} & 0 & 0 & 0 \\
\frac{1}{3} & -\frac{1}{3} & \frac{1}{3} & 0 & 0 & \frac{2}{3} & -\frac{1}{3} & -\frac{1}{3} \\
\frac{1}{3} & -\frac{1}{3} & \frac{1}{3} & 0 & 0 & -\frac{1}{3} & \frac{2}{3} & -\frac{1}{3} \\
\frac{1}{3} & -\frac{1}{3} & \frac{1}{3} & 0 & 0 & -\frac{1}{3} & -\frac{1}{3} & \frac{2}{3}
\end{pmatrix}$$


It is now a routine calculation to see that the square in Conjecture 1.1 commutes.

5 The p-adic group of type F_4

We now verify Conjecture 1.1 for the split, simply connected p-adic group $G = F_4(F)$. Since the fundamental group Ω is trivial, G is also adjoint. Its dual group is $G^\vee = F_4(\mathbb{C})$, and we let $\{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$ be the simple roots of $G^\vee$, with Dynkin diagram



We denote by α_0 and α^0 the highest short root and highest long root of the root system, respectively, so the simple affine roots of G are $S_{\text{aff}} = \{-\check{\alpha}_0, \check{\alpha}_4, \check{\alpha}_3, \check{\alpha}_2, \check{\alpha}_1\}$ with affine Dynkin diagram



The group G has five maximal open compact subgroups $\{K_0, K_4, K_3, K_2, K_1\}$ up to conjugacy, where K_i corresponds to $J_i := S_{\text{aff}} \setminus \{\alpha_i^\vee\}$ for $0 \leq i \leq 4$. The reductive quotients $\overline{K}_0, \overline{K}_4, \overline{K}_3, \overline{K}_2$ and $\overline{K}_1$ are finite reductive groups of type $F_4, A_1 \times C_3, A_2 \times \tilde{A}_2, A_3 \times \tilde{A}_1$ and B_4, respectively. In this section, we prove the following.

Theorem 5.1. *Let G be a simple p-adic group of type F_4. Then the diagram*

$$\begin{array}{ccc} R_{\text{un,ell}}(G) & \xrightarrow{\mathrm{FT}^\vee_{\text{ell}}} & R_{\text{un,ell}}(G) \\ \downarrow{\scriptstyle \mathrm{res}^{\text{cpt}}_{\text{un}}} & & \downarrow{\scriptstyle \mathrm{res}^{\text{cpt}}_{\text{un}}} \\ \bigoplus_{i=0}^4 R_{\text{un}}(\overline{K}_i) & \xrightarrow{(\mathrm{FT}^{K_i})_i} & \bigoplus_{i=0}^4 R_{\text{un}}(\overline{K}_i), \end{array}$$

commutes.

Since a Chevalley involution $C : F_4(\mathbb{C}) \to F_4(\mathbb{C})$ is an inner automorphism,

$$\mathrm{FT}^\vee_{\text{ell}}(\Pi(u, s, h)) = \Pi(C(u), C(h), C(s)) = \Pi(u, h, s).$$

The above result was already proven by Ciubotaru, in an unpublished paper [Ciu20] for the reduction to the hyperspecial parahoric subgroup K_0. We complete the proof for the remaining 4 parahoric subgroups.

Firstly, we compute the basis $\{\Pi(u, s, h) \mid u \in G^\vee \text{ unipotent}, (s, h) \in \Gamma_u \backslash \mathcal{Y}(\Gamma_u)_{\text{ell}}\}$ of the space $R_{\text{un,ell}}(G)$ (see Section 3.2 for the definitions). The classification of all elliptic pairs of $F_4(\mathbb{C})$ has been previously computed in the literature, and the list is given in Table 3. For each unipotent $u \in F_4(\mathbb{C})$, we also give the unipotent $F_4(\mathbb{F}_q)$-family $\mathcal{F}_u$ whose special element is the W-representation afforded by $\varepsilon \otimes H^{\text{top}}(\mathcal{B}^u)^{\mathbf{1}}$ under the Springer correspondence. Then $\Gamma_{\mathcal{F}_u}$ is the finite group canonically associated to the family $\mathcal{F}_u$.

Unipotent	Γ_u^0	A_u	$\lvert\Gamma_u\backslash\mathcal{Y}(\Gamma_u)_{\mathrm{ell}}\rvert$	$\mathcal{F}_u$	$\Gamma_{\mathcal{F}_u}$
F_4	1	1	1	$\phi_{1,24}$	1
$F_4(a_1)$	1	S_2	4	$\phi_{4,13}$	C_2
$F_4(a_2)$	1	S_2	4	$\phi_{9,10}$	1
B_3	PGL(2)	1	1	$\phi''_{8,9}$	1
$F_4(a_3)$	1	S_4	21	$\phi_{12,4}$	S_4

Table 3: Elliptic pairs for $F_4(\mathbb{C})$

For example, when u has type $F_4(a_3)$, $\Gamma_u \cong S_4$ and there are 21 virtual characters $\Pi(u,s,h)$ (below $\tau = g_2 \cdot g_2'$ and γ is conjugate to g_2' in S_4 but not in $Z_{S_4}(g_2') = D_8$):

$$\Pi(F_4(a_3),1,1) = [F_4(a_3),4] + 3[F_4(a_3),31] + 3[F_4(a_3),211] + [F_4(a_3),1^4] + 2[F_4(a_3),22];$$
$$\Pi(F_4(a_3),1,g_2) = [F_4(a_3),4] + [F_4(a_3),31] - [F_4(a_3),211] - [F_4(a_3),1^4];$$
$$\Pi(F_4(a_3),1,g_2') = [F_4(a_3),4] - [F_4(a_3),31] - [F_4(a_3),211] + [F_4(a_3),1^4] + 2[F_4(a_3),22];$$
$$\Pi(F_4(a_3),1,g_3) = [F_4(a_3),4] + [F_4(a_3),1^4] - [F_4(a_3),22];$$
$$\Pi(F_4(a_3),1,g_4) = [F_4(a_3),4] - [F_4(a_3),31] + [F_4(a_3),211] - [F_4(a_3),1^4];$$

$$\Pi(F_4(a_3),g_2,1) = [C(42)A_1,{+}{+}] + [C(42)A_1,{+}{-}] + [C(42)A_1,{-}{+}] + [C(42)A_1,{-}{-}];$$
$$\Pi(F_4(a_3),g_2,g_2) = [C(42)A_1,{+}{+}] - [C(42)A_1,{+}{-}] + [C(42)A_1,{-}{+}] - [C(42)A_1,{-}{-}];$$
$$\Pi(F_4(a_3),g_2,\tau) = [C(42)A_1,{+}{+}] + [C(42)A_1,{+}{-}] - [C(42)A_1,{-}{+}] - [C(42)A_1,{-}{-}];$$
$$\Pi(F_4(a_3),g_2,g_2') = [C(42)A_1,{+}{+}] - [C(42)A_1,{+}{-}] - [C(42)A_1,{-}{+}] + [C(42)A_1,{-}{-}];$$

$$\Pi(F_4(a_3),g_2',1) = [B_4(531),\mathbf{1}] + 2[B_4(531),r] + [B_4(531),\varepsilon'] + [B_4(531),\varepsilon''] + [B_4(531),\varepsilon];$$
$$\Pi(F_4(a_3),g_2',g_2') = [B_4(531),\mathbf{1}] - [B_4(531),\varepsilon'] + [B_4(531),\varepsilon''] - [B_4(531),\varepsilon];$$
$$\Pi(F_4(a_3),g_2',g_2'') = [B_4(531),\mathbf{1}] - 2[B_4(531),r] + [B_4(531),\varepsilon'] + [B_4(531),\varepsilon''] + [B_4(531),\varepsilon];$$
$$\Pi(F_4(a_3),g_2',g_4) = [B_4(531),\mathbf{1}] - [B_4(531),\varepsilon'] - [B_4(531),\varepsilon''] + [B_4(531),\varepsilon];$$
$$\Pi(F_4(a_3),g_2',\gamma) = [B_4(531),\mathbf{1}] + [B_4(531),\varepsilon'] - [B_4(531),\varepsilon''] - [B_4(531),\varepsilon];$$

$$\Pi(F_4(a_3),g_3,1) = [2A_2,\mathbf{1}] + [2A_2,\theta] + [2A_2,\theta^2];$$
$$\Pi(F_4(a_3),g_3,g_3) = [2A_2,\mathbf{1}] + \theta^2[2A_2,\theta] + \theta[2A_2,\theta^2];$$
$$\Pi(F_4(a_3),g_3,g_3^{-1}) = [2A_2,\mathbf{1}] + \theta[2A_2,\theta] + \theta^2[2A_2,\theta^2];$$

$$\Pi(F_4(a_3), g_4, 1) = [A_3A_1, 1] + [A_3A_1, i] + [A_3A_1, -1] + [A_3A_1, -i];$$
$$\Pi(F_4(a_3), g_4, g_4) = [A_3A_1, 1] - i[A_3A_1, i] - [A_3A_1, -1] - i[A_3A_1, -i];$$
$$\Pi(F_4(a_3), g_4, g_2') = [A_3A_1, 1] - [A_3A_1, i] + [A_3A_1, -1] - [A_3A_1, -i];$$
$$\Pi(F_4(a_3), g_4, g_4^{-1}) = [A_3A_1, 1] + i[A_3A_1, i] - [A_3A_1, -1] - i[A_3A_1, -i];$$

These virtual characters span a 21-dimensional space inside the 31-dimensional space $R_{\text{un,ell}}(G)$. All remaining 10 basis elements are easily computable. The proof of Theorem 5.1 relies on an explicit computation of the restriction map

$$\text{res}^{\text{cpt}}_{\text{un}} : R_{\text{un,ell}}(G) \longrightarrow \bigoplus_{i=0}^{5} R_{\text{un}}(\overline{K}_i).$$

Most of this work was completed by Reeder in [Ree00]. Nevertheless, his tables contain a few mistakes (systematically arising from the two representations $[F_4(a_3), 31]$ and $[F_4(a_3), 211]$) and do not include the representations whose unipotent label u has type B_3 (for this is not a distinguished unipotent element of $F_4(\mathbb{C})$). We tabulate these results at the end of the section, fixing the mistakes and adding the new restrictions.

Let us briefly discuss how the parahoric restrictions are computed. The unipotent representations of $F_4(F)$ fall into 9 unipotent blocks, together with their associated Hecke algebras. There are 7 supercuspidal blocks, one for each cuspidal unipotent representation of $F_4(\mathbb{F}_q)$. The corresponding Hecke algebras $\mathcal{H}(F_4(F), F_4(\mathcal{O}_F), \sigma)$ are all isomorphic to $\mathbb{C}$ so these blocks contain a unique unipotent representation. This information can be read from Lusztig's arithmetic-geometric correspondence tables in [Lus95a, pg. 577]. The Langlands parameters of these representations are

$$[F_4(a_3), 1^4],\ [C_3(42), --],\ [B_4(531), \varepsilon],\ [2A_2, \theta],\ [2A_2, \theta^2],\ [A_1A_3, i] \text{ and } [A_1A_3, -i],$$

where we are using Notation 3.7 for the Langlands parameters. The parahoric restrictions of these representations are straightforward to compute, for they are irreducible for the hyperspecial parahoric K_0 and vanish for the other ones.

The principal block contains all Iwahori-spherical representations, that is, all representations $\pi(us, \phi)$ such that $H(\mathcal{B}_s^u)^\phi \neq 0$. The corresponding Hecke algebra $\mathcal{H}(F_4(F), I, \mathbf{1})$ is the affine Hecke algebra of type F_4 with equal parameters. We compute the parahoric restrictions, using Springer theory, the Green tables of $F_4(\mathbb{F}_q)$ and the methods described in subsection 4.2.

Finally, there is an intermediate block whose Hecke algebra $\mathcal{H}(F_4(F), B_2, \sigma)$ is the affine Hecke algebra of type C_2 with unequal parameters that we introduced in Example 4.8. All Langlands parameters of the representations of interest in this block are *elliptic*, so it suffices to compute the square integrable representations of $\mathcal{H}(F_4(F), B_2, \sigma)$ to determine the desired parahoric restrictions. This can be achieved

using the theory of weight diagrams briefly discussed in Section 4.3 (see also [Ciu20, §9]). The Langlands parameters of these representations are

$$[B_4, -],\ [C_3A_1, -],\ [C_3(42)A_1, -+],\ [B_4(531), r] \text{ and } [A_1A_3, -1].$$

We explicitly discuss the $[A_1A_3, -1]$ case in Example 5.5, and we tabulate the rest of the parahoric restrictions in Section 7. The proof of Theorem 5.1 follows immediately from Tables 7, 9, 11 and 14.

Let us finish this section by discussing a few examples in detail to illustrate the main ideas involved in the calculations.

5.1 Examples of parahoric restrictions

The first example is the computation of a parahoric restrictions with respect to the maximal hyperspecial parahoric K_0 using Springer Theory and Alvis' tables ([Alv05]).

Example 5.2. Consider the representation $[B_4(531), \mathbf{1}]$, with u of type $F_4(a_3)$ and $s \in \Gamma_u$ a product of two 2-cycles. The centralizer $Z_{G^\vee}(s)$ has type B_4 and $u \in Z_{G^\vee}(s)$ is subregular. Thus, the Springer fibre $\mathcal{B}_s^u$ is 2 dimensional, and by reading the Green functions we obtain

$$H^*(\mathcal{B}_s^u)^{\mathbf{1}} = [2,2] + [3,1] + [4,\cdot] \quad \text{as } W_s\text{-modules.}$$

Since all characters $\chi_{s^w}^J$ are trivial when $J = J_0$, Alvis' tables yield

$$\mathrm{Ind}_{W_s}^W [2,2] + [3,1] + [4,\cdot] = \phi_{9,2} + \phi''_{9,6} + \phi''_{8,3} + \phi_{4,1} + \phi''_{2,4} + \phi_{1,0}.$$

To obtain the parahoric restriction of the representation $[B_4(531), \mathbf{1}]$ with respect to K_0, we twist by ε and by the outer automorphism of W swapping long and short roots, so

$$[B_4(531), \mathbf{1}] \xrightarrow{\mathrm{res}_{\mathrm{un}}^{K_0}} \phi_{9,10} + \phi''_{9,6} + \phi''_{8,9} + \phi_{4,13} + \phi''_{2,16} + \phi_{1,24}.$$

Once the parahoric restrictions for the maximal hyperspecial have been calculated, we can turn our attention towards computing the rest. The first step is to determine when the characters χ_t^J are trivial, which happens often and it can be easily determined from [Ree00, Lemma 8.3]. Whenever this is the case, we can use Lemma 4.7 and Alvis' tables to compute the restriction directly. Let us discuss another interesting example.

Example 5.3. Consider the parahoric restriction of the Iwahori-spherical representation $[A_1A_3, \mathbf{1}]$, where u is of type $F_4(a_3)$ and $s \in \Gamma_u$ is a 4-cycle, with respect to the parahoric subgroup K_2. Note that $W_{J_2} = \langle s_0, s_4, s_3, s_1 \rangle \cong W(A_3) \times W(A_1)$. The semisimple part s can be chosen to be $s = \check{\alpha}_3(-1)\check{\alpha}_4(i)$ so that $W_s = \langle s_4, s_2, s_1, s^0 \rangle \cong W(A_1) \times W(A_3)$. In addition, u is a regular unipotent element in

$Z_{G^\vee}(s)$, so $H^*(\mathcal{B}_s^u)^{\mathbf{1}} = \mathbf{1}$ as a W_s-module. Next,

$$W_s\backslash W/W_{J_2} = \{1, w\},$$

where $w \in W$ is explicitly computable with computer algebra programs such as GAP. In addition,

$$W_{J_2,s} = \langle s_4, s_1\rangle \cong C_2 \times C_2, \quad \text{while} \quad W_{J_2,s^w} = \langle d\rangle \cong C_4,$$

for some $d \in W$ of order 4. Since $W_{J_2,s}$ is a reflection subgroup of W_{J_2}, the character $\chi_s^{J_2}$ is trivial. On the other hand, W_{J_2,s^w} is not a reflection subgroup of W_{J_2}, and $\chi_{s^w}^{J_2} = \chi$ is a primitive character of C_4. The induced representations are then given by

$$\mathrm{Ind}_{W_{J_2,s}}^{W_{J_2}} \mathbf{1} = [[4] + 2[311] + [22] + [211]] \boxtimes [2] \quad \text{and} \quad \mathrm{Ind}_{W_{J_2,s^w}}^{W_{J_2}} \chi = [[31] + [211]] \boxtimes [[2] + [11]].$$

Adding these and twisting by sign gives the required parahoric restriction.

Example 5.4. A more challenging example is the calculation of the parahoric restriction of $[2A_2, \mathbf{1}]$, with u of type $F_4(a_3)$ and $s \in \Gamma_u$ is a 3-cycle, with respect to K_3. This is the only case where a characters $\chi_{s^w}^{J_3}$ may be non-trivial. Note that $W_{J_3} = \langle s_0, s_4, s_2, s_1\rangle \cong W(A_2) \times W(A_2)$. The semisimple part can be taken to be $s = \check{\alpha}_3(\xi_3)\check{\alpha}_4(\xi_3^2)$, where ξ_3 is a primitive third root of unity. In particular, $Z_{G^\vee}(s)$ is a pseudo-Levi subgroup of type $A_2 \times A_2$ generated by the affine simple roots $\{\alpha_4, \alpha_3, \alpha_1, -\alpha^0\}$ and $W_s = \langle s_4, s_3, s_1, s^0\rangle$. In addition, the unipotent element u is regular in $Z_{G^\vee}(s)$, so $H^*(\mathcal{B}_s^u)^{\mathbf{1}} = \mathbf{1}$ as W_s-modules. Next,

$$W_s\backslash W/W_{J_3} = \{w_1 = 1, w_2, w_3, w_4, w_5\},$$

where $w_i \in W$ are representatives of double cosets with sizes $324, 324, 36, 36$ and 432, respectively. The subgroups

$$W_{J_3,s^{w_3}} = W_{J_3,s^{w_4}} = W_{J_3} \quad \text{and} \quad W_{J_3,s} \cong W_{J_3,s^{w_2}} \cong C_2 \times C_2$$

are all reflections subgroups of W_{J_3}, so the characters $\chi_{s^{w_i}}^{J_3}$ are trivial for $i = 1, 2, 3, 4$. The remaining coset is more interesting since

$$W_{J_3,s^{w_5}} = \langle d\rangle \cong C_3,$$

is no longer a reflection subgroup, and $\chi_{s^{w_4}}^{J_3}$ is a primitive character χ of C_3.

The induced representations are then given by

$$\mathrm{Ind}_{W_{J_3,s}}^{J_3} \mathbf{1} = \mathrm{Ind}_{W_{J_3,s^{w_2}}}^{J_3} \mathbf{1} = [3,3] + [3,21] + [21,3] + [21,21], \quad \mathrm{Ind}_{W_{J_3,s^{w_3}}}^{J_3} \mathbf{1} = \mathrm{Ind}_{W_{J_3,s^{w_4}}}^{J_3} \mathbf{1} = [3,3]$$
$$\text{and} \quad \mathrm{Ind}_{W_{J_3,s^{w_5}}}^{J_3} \chi = [21,3] + [3,21] + [21,21] + [21,1^3] + [1^3,21].$$

Adding these and twisting by sign gives the required parahoric restriction.

Finally, let us discuss the parahoric restrictions of the non-Iwahori-spherical representation $[A_1A_3, -1]$.

Example 5.5. Let V be the representation labelled by $[A_1A_3, -1]$, K a parahoric subgroup of type B_2 and let σ be the unique cuspidal unipotent representation of $B_2(\mathbb{F}_q)$. Since the types of the reductive quotients of K_2 and K_3 ($A_3 \times \tilde{A}_1$ and $A_2 \times \tilde{A}_2$, respectively) do not contain B_2 as a factor, the parahoric restrictions of V to K_2 and K_3 vanish.

On the other hand, V^σ is a square integrable representation of $\mathcal{H}(F_4(F), B_2, \sigma) \cong \mathcal{H}(K_{J_0}, K, \sigma)\tilde{\otimes}\mathcal{O}[\mathbf{T}]$, whose restriction to $\mathcal{O}[\mathbf{T}]$ is

$$V^\sigma|_{\mathcal{O}[\mathbf{T}]} = V^\sigma(q^{-2}, -q^{-1}) \oplus V^\sigma(-q^{-1}, q^{-2}),$$

both subspaces being one-dimensional. By choosing a basis element in each subspace, the rational functions $e_{\alpha_1}, e_{\alpha_2} \in \mathcal{O}[\mathbf{T}]$ act on V^σ by $\mathrm{diag}(-q^{-1}, -q)$ and $\mathrm{diag}(-q^{-1}, q^{-2})$. Using the formula for the twisted multiplication in $\mathcal{H}(F_4(F), B_2, \sigma)$, the action of the generators T_1 and T_2 can be computed explicitly, and we obtain

$$T_{s_1} \longrightarrow \frac{1}{q+1}\begin{pmatrix} q^3-1 & q^3+q \\ q^4+1 & q^4-q \end{pmatrix} \quad \text{and} \quad T_{s_2} \longrightarrow \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Finally, one can trace back through the isomorphisms $\mathcal{H}(F_4(F), B_2, \sigma) \cong \mathcal{H}(K_{J_0}, K, \sigma) \tilde{\otimes} \mathcal{O}[\mathbf{T}]$ to obtain that $T_{\tilde{s}_0}$ acts by

$$T_{\tilde{s}_0} \longrightarrow \frac{1}{q^3+q^2}\begin{pmatrix} -q^6+q^3-q^2-1 & q^5+q^3-q^2-1 \\ -q^7+q^4-q^3+1 & q^6+q^4-q^3+1 \end{pmatrix}.$$

Thus, the action of $\tilde{W}(K, \sigma)$ on $V^\sigma_{q=1}$ is given by


$$s_1 \longrightarrow \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad s_2 \longrightarrow \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \quad \text{and} \quad \tilde{s}_0 \longrightarrow \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}.$$

The parahoric restrictions are now obtained by restricting to reflection subgroups. Restriction to the reflection subgroup $\langle s_1, s_2\rangle \cong W(B_2)$ gives the parahoric restriction to K_0, restriction to $\langle s_1, \tilde{s}_0\rangle \cong W(B_2)$ gives the parahoric restriction to K_1 and restriction to $\langle s_2, \tilde{s}_0\rangle \cong W(A_1 \times A_1)$ gives the parahoric restriction to K_4. In each case, there is a bijection between representations of the Weyl group and unipotent representations of $\overline{K}_i$ in the series generated by σ, which we use implicitly. We refer the reader to [Car85, §10.11 and §13.9] Throughout, we use Lusztig's notation for the unipotent representations. We obtain

- Firstly, $V^\sigma|_{\langle s_1, s_2\rangle} = [11, \emptyset] + [\emptyset, 11]$, so $[A_1A_3, -1] \xrightarrow{\mathrm{res}^{K_0}_{\mathrm{un}}} B_2[\varepsilon'] + B_2[\varepsilon]$.
- Secondly, $V^\sigma|_{\langle s_1, \tilde{s}_0\rangle} = [1, 1]$, so $[A_1A_3, -1] \xrightarrow{\mathrm{res}^{K_1}_{\mathrm{un}}} ({}^{0\ 1}_{1}{}^{2\ 4})$.
- Finally, $V^\sigma|_{\langle s_2, \tilde{s}_0\rangle} = [11] \boxtimes [2] + [11] \boxtimes [11]$, so $[A_1A_3, -1] \xrightarrow{\mathrm{res}^{K_4}_{\mathrm{un}}} ({}^{0\ 1}_{1}{}^{2\ 3}) \boxtimes \mathbf{1} + ({}^{0\ 1}_{1}{}^{2\ 3}) \boxtimes \mathrm{St}$.

6 The p-adic group of type E_6

In this section, we let $G = E_6(F)_{\mathrm{sc}}$ be the simple p-adic of type E_6 of simply connected type. Its affine Dynkin diagram is



and its fundamental group is $\Omega \cong \mathbb{Z}/3\mathbb{Z}$. The group G has 7 conjugacy classes of open maximal open compact subgroups $K_i,\ i = 0, 1, \ldots, 6$, one for each $J_i = S_{\mathrm{aff}} \setminus \{\check{\alpha}_i\},\ i = 0, 1, \ldots, 6$. In this section, we prove

Theorem 6.1. *Let G be a simple p-adic group of type E_6 of simply connected type. Then the diagram*

$$\begin{array}{ccc} R_{\mathrm{un,ell}}(G) & \xrightarrow{\mathrm{FT}^{\vee}_{\mathrm{ell}}} & R_{\mathrm{un,ell}}(G) \\ \downarrow{\scriptstyle \mathrm{res}^{\mathrm{cpt}}_{\mathrm{un}}} & & \downarrow{\scriptstyle \mathrm{res}^{\mathrm{cpt}}_{\mathrm{un}}} \\ \bigoplus_{i=0}^{6} R_{\mathrm{un}}(\overline{K}_i) & \xrightarrow{(\mathrm{FT}^{K_i})_i} & \bigoplus_{i=0}^{6} R_{\mathrm{un}}(\overline{K}_i), \end{array}$$

commutes, up to roots of unity.

The classes parametrized by J_0, J_1, J_6 and J_2, J_3, J_5 are conjugate under the action of Ω, respectively, and in particular they have the same reductive type. By Proposition 3.17, conditional to the verification of condition (**??**), it is enough to prove the conjecture for the conjugacy classes of maximal open compact subgroups corresponding to J_0, J_2 and J_4.

Next, we classify the elliptic pairs in $G^{\vee} = E_6(\mathbb{C})_{\mathrm{ad}}$. This classification is well known in the literature for $E_6(\mathbb{C})_{\mathrm{sc}}$, but for $E_6(\mathbb{C})_{\mathrm{ad}}$ is slightly more involved. For most unipotent elements $u \in G^{\vee}$, the reductive part Γ_u of its centralizer is connected. When this is the case, there exists a precise description of the elliptic pairs in Γ_u.

Proposition 6.2. [ACR25, Proposition 8.7] *Suppose that Γ_u is connected. Then there is a bijection*

$$\Gamma_u \backslash \mathcal{Y}(\Gamma_u)_{ell} \leftrightarrow W(\Gamma_u) \backslash \{(s, w) \mid s \in T \text{ is regular, } w \in W(\Gamma_u) \text{ is elliptic, } s \in T^w\},$$

where the map is obtained by conjugating an elliptic pair (s, h) so that s lies in a fixed torus T of Γ_u and h is the lift of some $w \in W(\Gamma_u)$ such that $s \in T^w$.

With this in mind we can give a precise description of the elliptic pairs inside $G^{\vee} = E_6(\mathbb{C})_{\mathrm{ad}}$. We use

the same notation as in Table 3.

Unipotent	Γ_u^0	A_u	$\lvert\Gamma_u\backslash\mathcal{Y}(\Gamma_u)_{\mathrm{ell}}\rvert$	$\mathcal{F}_u$	$\Gamma_{\mathcal{F}_u}$
E_6	1	1	1	$\phi_{1,36}$	1
$E_6(a_1)$	1	1	1	$\phi_{6,25}$	1
$E_6(a_3)$	1	S_2	4	$\phi_{30,15}$	C_2
D_4	PGL_3	1	2	$\phi_{24,12}$	1
$D_4(a_1)$	$(\mathbb{C}^\times)^2$	S_3	12	$\phi_{80,7}$	S_3
A_2	$(\mathrm{SL}_3\times\mathrm{SL}_3)/\mu_3$	1	2	$\phi_{30,3}$	C_2

Table 4: Elliptic pairs for $E_6(\mathbb{C})_{\mathrm{ad}}$

Let us briefly describe the elliptic pairs above explicitly. When $\Gamma_u^0 = \{1\}$ the description is straightforward. Next, consider the semisimple matrices

$$s_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \zeta_3 & 0 \\ 0 & 0 & \zeta_3^2 \end{pmatrix} \quad \text{and} \quad w_3 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} \quad \text{in } \mathrm{SL}_3(\mathbb{C}).$$

We also let $s_{3,3} = (s_3, s_3)$ and $w_{3,3} = (w_3, w_3)$ be matrices in $\mathrm{SL}_3 \times \mathrm{SL}_3$. We abuse notation and denote their images in central quotients with the same symbols as necessary.

When u has type D_4, then $\Gamma_u\backslash\mathcal{Y}(\Gamma_u)_{\mathrm{ell}} = \{(s_3, w_3), (s_3, w_3^{-1})\}$ and $A_{us_3} = R(u) \cong C_3$, while if u has type A_2, then $\Gamma_u\backslash\mathcal{Y}(\Gamma_u)_{\mathrm{ell}} = \{(s_{3,3}, w_{3,3}), (s_{3,3}, w_{3,3}^{-1})\}$ and $A_{us_{3,3}} = A^0_{us_{3,3}} \times R(u) \cong C_2 \times C_3$. The remaining case when u has type $D_4(a_1)$ is the most interesting. Then $\Gamma_u = (\mathbb{C}^\times)^2 \rtimes S_3$, where the 3-cycle (123) in S_3 acts by $(a, b) \mapsto (b, \frac{1}{ab})$ and the 2-cycle (12) acts by $(a, b) \mapsto (\frac{1}{b}, \frac{1}{a})$. A lengthy computation then shows that Γ_u has 12 elliptic pairs up to conjugacy. To describe them, let

$$1 = ((1,1), e),\ z = ((\zeta_3, \zeta_3), e),\ g_3 = ((1,1), (123)) \in \Gamma_u.$$

Then the 12 elliptic pairs up to conjugacy are $\{(z^k, g_3), (g_3, z^k), (g_3, z^k g_3), (g_3, z^k g_3^{-1}) \mid k = 0, 1, 2\}$, where

$$A_u = A_u^0 = S_3, \quad A_{uz} = R(u) = \langle g_3 \rangle \cong C_3 \quad \text{and} \quad A_{ug_3} = A^0_{ug_3} \times R(u) = \langle g_3 \rangle \times \langle z \rangle \cong C_3 \times C_3$$

Thus, $R_{\mathrm{un,ell}}(G)$ is a 22-dimensional vector space, and the proof of 6.1 relies again on an explicit computation of $\mathrm{FT}^\vee_{\mathrm{ell}}$ and the parahoric restriction maps

$$\mathrm{res}^{K_i}_{\mathrm{un}} : R_{\mathrm{un,ell}}(G) \longrightarrow R_{\mathrm{un}}(\overline{K}_i), \quad \text{for } i = 0, 2, 4.$$

Next, we discuss the action of $\mathrm{FT}^\vee_{\mathrm{ell}}$ on the basis elements $\Pi(u, s, h)$. To this aim, let $C : G^\vee \to G^\vee$ be a Chevalley involution and let $T^\vee \subset G^\vee$ a maximal torus such that $C(t) = t^{-1}$ for all $t \in T^\vee$. When u

has type E_6, $E_6(a_1)$ and $E_6(a_3)$, all automorphisms of Γ_u are inner, and therefore

$$\mathrm{FT}^\vee_{\mathrm{ell}}(\Pi(u,s,h)) = \Pi(u,h,s).$$

When u has type D_4 or $D_4(a_1)$, we fix $L^\vee \subset G^\vee$ to be a standard Levi subgroup of type D_4 containing $T^\vee$. Since maximal tori in Γ_u have rank two, the centre $Z(L^\vee)$ is a maximal torus of Γ_u contained in $T^\vee$. In addition, the Chevalley involution acts by $C(\mathcal{X}_\alpha) = \mathcal{X}_{-\alpha}$ on root spaces, so it preserves $L^\vee$. Inside $L^\vee$, the element u is either regular or subregular, so there exists some $l \in L^\vee$ such that $lC(u)l^{-1} = u$. Therefore, the map $\mathrm{int}(l)\circ C : \Gamma_u \to \Gamma_u$ is a Chevalley involution since $\mathrm{int}(l)(C(t)) = t^{-1}$ for all $t \in Z(L^\vee) \subset T^\vee$. When u has type A_2, an analogous argument shows that there is some l in a standard Levi $L^\vee$ of type A_2 containing $T^\vee$ such that $\mathrm{int}(l) \circ C$ is a Chevalley involution of Γ_u. In section 3.4, we showed that

$$\begin{aligned}\mathrm{FT}^\vee_{\mathrm{ell}} : \Pi(D_4, s_3, w_3^{\pm 1}) &\longmapsto \Pi(D_4, s_3, w_3^{\pm 1}),\\ \Pi(A_2, s_{3,3}, w_{3,3}^{\pm 1}) &\longmapsto \Pi(A_2, s_{3,3}, w_{3,3}^{\pm 1}).\end{aligned}$$

Finally, when u has type $D_4(a_1)$, $\Gamma_u = (\mathbb{C}^\times)^2 \rtimes S_3$ and the automorphism $C_u(((a,b),\tau)) = ((\frac{1}{a},\frac{1}{b}),\tau)$ is a Chevalley involution. Thus, we have that $\mathrm{FT}^\vee_{\mathrm{ell}}$ swaps the basis vectors

$$\begin{gathered}\Pi(u,1,g_3) \longleftrightarrow \Pi(u,g_3,1), \quad \Pi(u,z,g_3) \longleftrightarrow \Pi(u,g_3,z^2), \quad \Pi(u,z^2,g_3) \longleftrightarrow \Pi(u,g_3,z),\\ \Pi(u,g_3,z^k g_3^{\pm 1}) \longleftrightarrow \Pi(u,g_3,z^k g_3^{\pm 1}), \quad k = 0,1,2.\end{gathered}$$

Lemma 6.3. *Condition* (**??**) *holds for all* $\omega \in \Omega$, $u \in G^\vee$ *unipotent and* $(s,h) \in \mathcal{Y}(\Gamma_u)_{\mathrm{ell}}$. *In particular, if Diagram* (11) *commutes for parahoric subgroups* K_0, K_2 *and* K_4, *then Theorem 6.1 holds.*

Proof. If the ordered triple $(C(u), C(h), C(s))$ is conjugate to the ordered triple (u,s,h), then condition (**??**) holds trivially. On the other hand, by Remark **??**, condition (**??**) holds when either $s = 1$ or $h = 1$. By the classification of elliptic pairs above, it only remains to prove condition (**??**) when u has type $D_4(a_1)$ and $(s,h) \in \{(z,g_3),(z^2,g_3)\}$. One can directly prove that, up to a choice of primitive third root of unity,

$$\overline{\tau}_{uz}(\omega)(g_3) = \theta, \quad \text{and} \quad \overline{\tau}_{ug_3}(\omega)(z) = \theta^2,$$

from which the Lemma easily follows. □

Finally, let us describe the unipotent blocks and the Langlands parameters for G. There are 8 unipotent blocks, 6 of which are supercuspidal, one for each pair (K_0, σ), where $K_0 \cong E_6(\mathcal{O}_F)$ is a hyperspecial parahoric subgroup (there are three conjugacy classes in G) and $\sigma \in \{E_6[\theta], E_6[\theta^2]\}$ is a cuspidal unipotent representation of $E_6(\mathbb{F}_q)$. The Hecke algebras $\mathcal{H}(E_6(F), K_0, \sigma) \cong \mathbb{C}$ so the block contains a unique Langlands parameter, which can be read from Lusztig's arithmetic-geometric tables (when G is adjoint), combined with Solleveld's results ([Sol20, Lemma 7.3]).

The principal block contains all representations $\pi(us,\phi)$ such that $H(\mathcal{B}_s^u)^\phi \neq 0$ and the corresponding Hecke algebra $\mathcal{H}(E_6(F),\mathcal{I},\mathbf{1})$ is the affine Hecke algebra of type E_6 of equal parameters. The parahoric restrictions are computed using the methods described in Section 4.

Finally, there is one intermediate Hecke algebra $\mathcal{H}(E_6(F),D_4(\mathcal{O}_F),\sigma)$, where σ is the unique cuspidal unipotent representation of $D_4(\mathbb{F}_q)$. The only Langlands parameter we encounter corresponding to this Hecke algebra is $[A_1A_5,-]$, where u has type $E_6(a_3)$, s is the unique non-trivial semisimple element commuting with u (of order 2), and ϕ is non-trivial.

We conclude this section with an explicit expression of the virtual characters $\Pi(u,s,h)$.

$$
\begin{aligned}
\Pi(E_6,1,1) &= [E_6,1];\\
\Pi(E_6(a_1),1,1) &= [E_6(a_1),1];\\
\Pi(E_6(a_3),1,1) &= [E_6(a_3),+]+[E_6(a_3),-];\\
\Pi(E_6(a_3),1,g_2) &= [E_6(a_3),+]-[E_6(a_3),-];\\
\Pi(E_6(a_3),g_2,1) &= [A_1A_5,+]+[A_1A_5,-];\\
\Pi(E_6(a_3),g_2,g_2) &= [A_1A_5,+]-[A_1A_5,-];\\
\Pi(D_4,s_3,w_3) &= [D_4(71),1]+\theta[D_4(71),\theta]+\theta^2[D_4(71),\theta^2];\\
\Pi(D_4,s_3,w_3^{-1}) &= [D_4(71),1]+\theta^2[D_4(71),\theta]+\theta[D_4(71),\theta^2];\\
\Pi(D_4(a_1),1,g_3) &= [D_4(a_1),1]+[D_4(a_1),\varepsilon]-[D_4(a_1),r];\\
\Pi(D_4(a_1),z,g_3) &= [D_4(53),\mathbf{1}]+\theta[D_4(53),\theta]+\theta^2[D_4(53),\theta^2];\\
\Pi(D_4(a_1),z^2,g_3) &= \Pi(D_4(a_1),z,g_3^{-1}) = [D_4(53),\mathbf{1}]+\theta^2[D_4(53),\theta]+\theta[D_4(53),\theta^2];\\
\Pi(D_4(a_1),g_3,z^kg_3^l) &= \sum_{0\leq i,j\leq 2}\theta^{il+jk}[3A_2,\theta^i\boxtimes\theta^j],\ k=0,1,2,\ l\in\{\pm 1\};\\
\Pi(A_2,s_{3,3},w_{3,3}) &= \sum_{0\leq i\leq 2}\theta^i([D_4(3311),\mathbf{1}\boxtimes\theta^i]+[D_4(3311),\epsilon\boxtimes\theta^i]);\\
\Pi(A_2,s_{3,3},w_{3,3}^{-1}) &= \sum_{0\leq i\leq 2}\theta^i([D_4(3311),\mathbf{1}\boxtimes\theta^i]+[D_4(3311),\epsilon\boxtimes\theta^i]).
\end{aligned}
$$

7 Parahoric restriction tables

We now display the parahoric restriction tables obtained by applying the methods described in Section 4 together with the geometric data from $F_4(\mathbb{C})$ and $E_6(\mathbb{C})_{\mathrm{ad}}$ discussed in Sections 5 and 6. For each maximal parahoric subgroup K of $F_4(F)$ or $E_6(F)_{\mathrm{sc}}$, we display two tables. The first table contains the parahoric restrictions of the relevant unipotent representations arising in the virtual representations $\Pi(u,s,h)$. The second table then contains the parahoric restrictions of $\Pi(u,s,h)$ and is easily obtained from the first by computing the corresponding linear combinations. The first columns the second tables are the characters in trivial families; then we write each family with the special character always coming first. We omit the tables for the maximal hyperspecial subgroup K_0 of $F_4(F)$ for the tables are lengthy and the conjecture was already verified by Ciubotaru in [Ciu20]. From these tables, together with Lusztig's nonabelian Fourier transform, Theorems 5.1 and 6.1 can be easily verified

For all tables, we use the notation described in 3.7 for the Langlands labels parametrizing the unipotent representations of $G = F_4(F)$, coinciding with Reeder's notation in [Ree00]. The unipotent representations of finite reductive groups of Lie type are labelled as follows. For groups of type A_n, they are parametrized by partitions λ of $n+1$. For groups of type B_n/C_n, the principal series representations are parametrized by double partitions $[\lambda,\mu]$ of n. To understand these tables, it is enough to know that groups of type B_3/C_3 have two non-principal series representations θ_1 and θ_2, while groups of type B_4/C_4 have five non-principal series representations $\theta_1,\dots,\theta_5$. They all lie in the same representation series, arising from the unipotent cuspidal representation of their Levi subgroup of type B_2/C_2 and of all them are part of a family of size 4 of unipotent representations.

Finally, we use the following useful notation shorthand. A finite group of Lie type of type A_1 has only two unipotent characters; the trivial and the Steinberg. Thus, whenever the type of reductive quotient is the product of two simple types, one of which is A_1, we give the multiplicities for each character χ of the other type as a pair $a,\overline{b}$, where a and $\overline{b}$ are the multiplicities of $\chi\otimes\mathbf{1}_{A_1}$ and $\chi\otimes\mathrm{St}_{A_1}$, respectively.

7.1 Parahoric restriction tables for F_4

Unipotent B_4-types

Table 5: Parahoric restriction of Iwahori-spherical $F_4(F)$-representations with respect to K_1

	$[21^2,\cdot]$	$[1^4,\cdot]$	$[1^3,1]$	$[2,11]$	$[11,2]$	$[11,11]$	$[1,21]$	$[1,1^3]$	$[\cdot,31]$	$[\cdot,22]$	$[\cdot,21^2]$	$[\cdot,1^4]$
$[F_4,1]$	0	0	0	0	0	0	0	0	0	0	0	1
$[F_4(a_1),+]$	0	0	0	0	0	0	0	1	0	0	0	1
$[F_4(a_1),-]$	0	1	0	0	0	0	0	0	0	0	0	1
$[B_4,+]$	0	0	0	0	0	0	0	0	0	0	1	0
$[F_4(a_2),+]$	0	0	0	0	0	1	0	1	0	0	1	1
$[F_4(a_2),-]$	0	0	0	0	0	0	0	0	0	1	0	0
$[C_3A_1,+]$	0	0	1	0	0	0	0	1	0	0	1	1
$[B_4(711),+]$	0	0	0	0	0	0	1	1	0	0	1	0
$[B_4(711),-]$	0	0	0	0	0	1	0	0	0	1	0	1
$[F_4(a_3),4]$	0	0	1	1	1	1	1	2	0	0	1	1
$[F_4(a_3),31]$	1	1	1	0	0	1	0	1	0	0	0	1
$[F_4(a_3),22]$	0	0	0	1	0	0	0	1	0	0	0	0
$[F_4(a_3),211]$	0	1	0	0	0	0	0	0	0	0	0	0
$[C_3(42)A_1,++]$	1	0	1	1	1	1	1	2	0	0	2	1
$[C_3(42)A_1,+-]$	0	1	1	0	0	1	0	0	0	0	0	1
$[B_4(531),1]$	0	0	0	1	1	0	1	1	1	0	2	0
$[B_4(531),\varepsilon']$	0	0	0	0	0	0	0	0	1	0	0	0
$[B_4(531),\varepsilon'']$	1	0	0	0	0	0	0	0	0	0	1	0
$[2A_2,1]$	0	0	1	0	1	1	1	1	0	0	1	1
$[A_1A_3,1]$	0	0	0	0	1	0	1	1	1	0	1	0

Table 6: Parahoric restriction of non-Iwahori-spherical $F_4(F)$-representations with respect to K_1

	$\theta_1 = \left(\begin{smallmatrix} 0 & 1 & 2 & 3 & 4 \\ & 1 & 2 & & \end{smallmatrix}\right)$	$\theta_2 = \left(\begin{smallmatrix} 0 & 1 & 2 & 3 \\ & 2 & & \end{smallmatrix}\right)$	$\theta_3 = \left(\begin{smallmatrix} 0 & 1 & 2 & 4 \\ & 1 & & \end{smallmatrix}\right)$	$\theta_4 = \left(\begin{smallmatrix} 0 & 2 & 3 \\ & - & \end{smallmatrix}\right)$	$\theta_5 = \left(\begin{smallmatrix} 0 & 1 & 4 \\ & - & \end{smallmatrix}\right)$
$[B_4,-]$	1	0	0	0	0
$[C_3A_1,-]$	0	1	0	0	0
$[C_3(42)A_1,-+]$	1	0	0	0	0
$[B_4(531),r]$	1	0	1	0	0
$[A_1A_3,-]$	0	0	1	0	0

Table 7: Parahoric restrictions of the virtual characters $\Pi(u,s,h)$ with respect to K_1

	$[11,2]$	$[1,21]$	$[\cdot,1^4]$	$[1,1^3]$	$[1^4,\cdot]$	$[\cdot,21^2]$	θ_1	$[11,11]$	$[1^3,1]$	$[\cdot,22]$	θ_2	$[2,11]$	$[21^2,\cdot]$	$[\cdot,31]$	θ_3
$\Pi(F_4,1,1)$	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0
$\Pi(F_4(a_1),1,1)$	0	0	2	1	1	0	0	0	0	0	0	0	0	0	0
$\Pi(F_4(a_1),1,g_2)$	0	0	0	1	-1	0	0	0	0	0	0	0	0	0	0
$\Pi(F_4(a_1),g_2,1)$	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0
$\Pi(F_4(a_1),g_2,g_2)$	0	0	0	0	0	1	-1	0	0	0	0	0	0	0	0
$\Pi(F_4(a_2),1,1)$	0	0	1	1	0	1	0	1	0	1	0	0	0	0	0
$\Pi(F_4(a_2),1,g_2)$	0	0	1	1	0	1	0	1	0	-1	0	0	0	0	0
$\Pi(F_4(a_2),g_2,1)$	0	0	1	1	0	1	0	0	1	0	1	0	0	0	0
$\Pi(F_4(a_2),g_2,g_2)$	0	0	1	1	0	1	0	0	1	0	-1	0	0	0	0
$\Pi(B_4(711),s,h)$	0	1	-1	1	0	1	0	-1	0	-1	0	0	0	0	0
$\Pi(F_4(a_3),1,1)$	1	1	4	7	6	1	0	4	4	0	0	3	3	0	0
$\Pi(F_4(a_3),1,g_2)$	1	1	2	3	0	1	0	2	2	0	0	1	1	0	0
$\Pi(F_4(a_3),1,g_2')$	1	1	0	3	-2	1	0	0	0	0	0	3	-1	0	0
$\Pi(F_4(a_3),1,g_3)$	1	1	1	1	0	1	0	1	1	0	0	0	0	0	0
$\Pi(F_4(a_3),1,g_4)$	1	1	0	1	0	1	0	0	0	0	0	1	-1	0	0
$\Pi(F_4(a_3),g_2,1)$	1	1	2	2	1	2	1	2	2	0	0	1	1	0	0
$\Pi(F_4(a_3),g_2,g_2)$	1	1	2	2	1	2	-1	2	2	0	0	1	1	0	0
$\Pi(F_4(a_3),g_2,\tau)$	1	1	0	2	-1	2	1	0	0	0	0	1	1	0	0
$\Pi(F_4(a_3),g_2,g_2')$	1	1	0	2	-1	2	-1	0	0	0	0	1	1	0	0
$\Pi(F_4(a_3),g_2',1)$	1	1	0	1	0	3	2	0	0	0	0	1	1	2	2
$\Pi(F_4(a_3),g_2',g_2)$	1	1	0	1	0	3	0	0	0	0	0	1	1	0	0
$\Pi(F_4(a_3),g_2',g_2')$	1	1	0	1	0	3	-2	0	0	0	0	1	1	2	-2
$\Pi(F_4(a_3),g_2',\gamma)$	1	1	0	1	0	1	0	0	0	0	0	1	-1	2	0
$\Pi(F_4(a_3),g_2',g_4)$	1	1	0	1	0	1	0	0	0	0	0	1	-1	0	0
$\Pi(F_4(a_3),g_3,h)$	1	1	1	1	0	1	0	1	1	0	0	0	0	0	0
$\Pi(F_4(a_3),g_4,h), h\in\{1,g_2'\}$	1	1	0	1	0	1	0	0	0	0	0	0	0	1	1
$\Pi(F_4(a_3),g_4,g_4^{\pm 1})$	1	1	0	1	0	1	0	0	0	0	0	0	0	1	-1

Table 8: Parahoric restriction of $F_4(F)$-representations with respect to K_2

	$[1^4]$	$[211]$	$[22]$	$[31]$	$[4]$
$[F_4, 1]$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[F_4(a_1), +]$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[F_4(a_1), -]$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[B_4, +]$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[F_4(a_2), +]$	$1, \bar{2}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[F_4(a_2), -]$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[C_3A_1, +]$	$1, \bar{2}$	$1, \bar{2}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[B_4(711), +]$	$0, \bar{1}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$
$[B_4(711), -]$	$1, \bar{1}$	$0, \bar{1}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[F_4(a_3), 4]$	$2, \bar{3}$	$3, \bar{5}$	$1, \bar{1}$	$1, \bar{2}$	$0, \bar{0}$
$[F_4(a_3), 31]$	$3, \bar{2}$	$2, \bar{2}$	$1, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[F_4(a_3), 22]$	$1, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$
$[F_4(a_3), 211]$	$1, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[C_3(42)A_1, ++]$	$2, \bar{3}$	$4, \bar{6}$	$1, \bar{1}$	$1, \bar{2}$	$0, \bar{0}$
$[C_3(42)A_1, +-]$	$2, \bar{2}$	$1, \bar{1}$	$1, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[B_4(531), 1]$	$0, \bar{1}$	$2, \bar{4}$	$0, \bar{1}$	$1, \bar{3}$	$0, \bar{0}$
$[B_4(531), \varepsilon'']$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[B_4(531), \varepsilon']$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$
$[2A_2, 1]$	$1, \bar{3}$	$2, \bar{4}$	$1, \bar{1}$	$1, \bar{1}$	$0, \bar{0}$
$[A_1A_3, 1]$	$0, \bar{1}$	$1, \bar{3}$	$0, \bar{1}$	$1, \bar{2}$	$0, \bar{0}$

Table 9: Parahoric restrictions of the virtual characters $\Pi(u, s, h)$ with respect to K_2

	$[1^4]$	$[211]$	$[22]$	$[31]$	$[4]$
$\Pi(F_4, 1, 1)$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$\Pi(F_4(a_1), 1, 1)$	$2, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$\Pi(F_4(a_1), 1, g_2)$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$\Pi(F_4(a_1), g_2, 1)$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$\Pi(F_4(a_1), g_2, g_2)$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$\Pi(F_4(a_2), 1, 1)$	$1, \bar{2}$	$1, \bar{2}$	$0, \bar{2}$	$0, \bar{0}$	$0, \bar{0}$
$\Pi(F_4(a_2), 1, g_2)$	$1, \bar{2}$	$1, \bar{2}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$\Pi(F_4(a_2), g_2, 1)$	$1, \bar{2}$	$1, \bar{2}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$\Pi(F_4(a_2), g_2, g_2)$	$1, \bar{2}$	$1, \bar{2}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$\Pi(B_4(711), s, h)$	$-1, \bar{0}$	$1, \bar{1}$	$-1, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$
$\Pi(F_4(a_3), 1, 1)$	$16, \bar{9}$	$11, \bar{13}$	$4, \bar{1}$	$1, \bar{4}$	$0, \bar{0}$
$\Pi(F_4(a_3), 1, g_2)$	$4, \bar{5}$	$5, \bar{7}$	$2, \bar{1}$	$1, \bar{2}$	$0, \bar{0}$
$\Pi(F_4(a_3), 1, g_2')$	$0, \bar{1}$	$3, \bar{5}$	$0, \bar{1}$	$1, \bar{4}$	$0, \bar{0}$
$\Pi(F_4(a_3), 1, g_3)$	$1, \bar{3}$	$2, \bar{4}$	$1, \bar{1}$	$1, \bar{1}$	$0, \bar{0}$
$\Pi(F_4(a_3), 1, g_4)$	$0, \bar{1}$	$1, \bar{3}$	$0, \bar{1}$	$1, \bar{2}$	$0, \bar{0}$
$\Pi(F_4(a_3), g_2, 1)$	$4, \bar{5}$	$5, \bar{7}$	$2, \bar{1}$	$1, \bar{2}$	$0, \bar{0}$
$\Pi(F_4(a_3), g_2, g_2)$	$4, \bar{5}$	$5, \bar{7}$	$2, \bar{1}$	$1, \bar{2}$	$0, \bar{0}$
$\Pi(F_4(a_3), g_2, \tau)$	$0, \bar{1}$	$3, \bar{5}$	$0, \bar{1}$	$1, \bar{2}$	$0, \bar{0}$
$\Pi(F_4(a_3), g_2, g_2')$	$0, \bar{1}$	$3, \bar{5}$	$0, \bar{1}$	$1, \bar{2}$	$0, \bar{0}$
$\Pi(F_4(a_3), g_2', 1)$	$0, \bar{1}$	$3, \bar{5}$	$0, \bar{1}$	$1, \bar{4}$	$0, \bar{0}$
$\Pi(F_4(a_3), g_2', g_2)$	$0, \bar{1}$	$3, \bar{5}$	$0, \bar{1}$	$1, \bar{2}$	$0, \bar{0}$
$\Pi(F_4(a_3), g_2', \gamma)$	$0, \bar{1}$	$1, \bar{3}$	$0, \bar{1}$	$1, \bar{4}$	$0, \bar{0}$
$\Pi(F_4(a_3), g_2', g_2')$	$0, \bar{1}$	$3, \bar{5}$	$0, \bar{1}$	$1, \bar{4}$	$0, \bar{0}$
$\Pi(F_4(a_3), g_2', g_4)$	$0, \bar{1}$	$1, \bar{3}$	$0, \bar{1}$	$1, \bar{2}$	$0, \bar{0}$
$\Pi(F_4(a_3), g_3, h)$	$1, \bar{3}$	$2, \bar{4}$	$1, \bar{1}$	$1, \bar{1}$	$0, \bar{0}$
$\Pi(F_4(a_3), g_4, h)$	$0, \bar{1}$	$1, \bar{3}$	$0, \bar{1}$	$1, \bar{2}$	$0, \bar{0}$

Table 10: Parahoric restriction of $F_4(F)$-representations with respect to K_3

	$[1^3,1^3]$	$[1^3,21]$	$[1^3,3]$	$[21,1^3]$	$[21,21]$	$[21,3]$	$[3,1^3]$	$[3,21]$	$[3,3]$
$[F_4,1]$	1	0	0	0	0	0	0	0	0
$[F_4(a_1),+]$	1	1	0	1	0	0	0	0	0
$[F_4(a_1),-]$	0	1	0	0	0	0	0	0	0
$[B_4,+]$	1	0	0	1	0	0	0	0	0
$[F_4(a_2),+]$	2	2	0	2	1	0	0	0	0
$[F_4(a_2),-]$	0	0	0	1	0	0	0	0	0
$[C_3A_1,+]$	2	2	0	1	1	0	0	0	0
$[B_4(711),+]$	2	3	1	1	1	0	0	0	0
$[B_4(711),-]$	1	1	0	1	1	0	0	0	0
$[F_4(a_3),4]$	4	4	1	4	4	1	1	1	0
$[F_4(a_3),31]$	1	3	2	0	2	1	0	0	0
$[F_4(a_3),22]$	0	1	1	1	1	0	1	0	0
$[F_4(a_3),211]$	0	0	1	0	0	0	0	0	0
$[C_3(42)A_1,++]$	4	5	1	4	5	1	1	1	0
$[C_3(42)A_1,+-]$	1	2	1	0	1	1	0	0	0
$[B_4(531),1]$	3	2	0	5	3	0	2	1	0
$[B_4(531),\varepsilon'']$	0	1	0	0	1	0	0	0	0
$[B_4(531),\varepsilon']$	0	0	0	1	0	0	1	0	0
$[2A_2,1]$	4	3	0	3	3	1	0	1	0
$[A_1A_3,1]$	3	1	0	4	2	0	1	1	0

Table 11: Parahoric restrictions of the virtual characters $\Pi(u,s,h)$ with respect to K_3

	$[1^3,1^3]$	$[1^3,21]$	$[1^3,3]$	$[21,1^3]$	$[21,21]$	$[21,3]$	$[3,1^3]$	$[3,21]$	$[3,3]$
$\Pi(F_4,1,1)$	1	0	0	0	0	0	0	0	0
$\Pi(F_4(a_1),1,1)$	1	2	0	1	0	0	0	0	0
$\Pi(F_4(a_1),1,g_2)$	1	0	0	1	0	0	0	0	0
$\Pi(F_4(a_1),g_2,1)$	1	0	0	1	0	0	0	0	0
$\Pi(F_4(a_1),g_2,g_2)$	1	0	0	1	0	0	0	0	0
$\Pi(F_4(a_2),1,1)$	2	2	0	3	1	0	0	0	0
$\Pi(F_4(a_2),1,g_2)$	2	2	0	1	1	0	0	0	0
$\Pi(F_4(a_2),g_2,1)$	2	2	0	1	1	0	0	0	0
$\Pi(F_4(a_2),g_2,g_2)$	2	2	0	1	1	0	0	0	0
$\Pi(B_4(711),s,h)$	1	2	1	0	0	0	0	0	0
$\Pi(F_4(a_3),1,1)$	7	15	12	6	12	4	3	1	0
$\Pi(F_4(a_3),1,g_2)$	5	7	2	4	6	2	1	1	0
$\Pi(F_4(a_3),1,g_2')$	3	3	0	6	4	0	3	1	0
$\Pi(F_4(a_3),1,g_3)$	4	3	0	3	3	1	0	1	0
$\Pi(F_4(a_3),1,g_4)$	3	1	0	4	2	0	1	1	0
$\Pi(F_4(a_3),g_2,1)$	5	7	2	4	6	2	1	1	0
$\Pi(F_4(a_3),g_2,g_2)$	5	7	2	4	6	2	1	1	0
$\Pi(F_4(a_3),g_2,\tau)$	3	3	0	4	4	0	1	1	0
$\Pi(F_4(a_3),g_2,g_2')$	3	3	0	4	4	0	1	1	0
$\Pi(F_4(a_3),g_2',1)$	3	3	0	6	4	0	3	1	0
$\Pi(F_4(a_3),g_2',g_2)$	3	3	0	4	4	0	1	1	0
$\Pi(F_4(a_3),g_2',\gamma)$	3	1	0	6	2	0	3	1	0
$\Pi(F_4(a_3),g_2',g_2')$	3	3	0	6	4	0	3	1	0
$\Pi(F_4(a_3),g_2',g_4)$	3	1	0	4	2	0	1	1	0
$\Pi(F_4(a_3),g_3,h)$	4	3	0	3	3	1	0	1	0
$\Pi(F_4(a_3),g_4,h)$	3	1	0	4	2	0	1	1	0

Iwahori-spherical $A_1 \times C_3$-types

Table 12: Parahoric restriction of Iwahori-spherical $F_4(F)$-representations with respect to K_4

	$[\cdot, 1^3]$	$[1^3, \cdot]$	$[11, 1]$	$[1, 11]$	$[21, \cdot]$	$[\cdot, 21]$	$[2, 1]$	$[1, 2]$	$[\cdot, 3]$	$[3, \cdot]$
$[F_4, 1]$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[F_4(a_1), +]$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[F_4(a_1), -]$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[B_4, +]$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[F_4(a_2), +]$	$1, \bar{2}$	$0, \bar{0}$	$0, \bar{1}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[F_4(a_2), -]$	$0, \bar{1}$	$1, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[C_3A_1, +]$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[B_4(711), +]$	$2, \bar{2}$	$1, \bar{1}$	$0, \bar{1}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[B_4(711), -]$	$0, \bar{1}$	$0, \bar{0}$	$1, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[F_4(a_3), 4]$	$2, \bar{2}$	$0, \bar{1}$	$1, \bar{2}$	$2, \bar{3}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{1}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[F_4(a_3), 31]$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$1, \bar{2}$	$0, \bar{1}$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$
$[F_4(a_3), 22]$	$1, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[F_4(a_3), 211]$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$
$[C_3(42)A_1, ++]$	$2, \bar{2}$	$0, \bar{1}$	$1, \bar{2}$	$2, \bar{3}$	$0, \bar{1}$	$2, \bar{2}$	$0, \bar{1}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[C_3(42)A_1, +-]$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$
$[B_4(531), 1]$	$3, \bar{2}$	$1, \bar{2}$	$1, \bar{2}$	$2, \bar{2}$	$0, \bar{1}$	$1, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[B_4(531), \varepsilon'']$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{1}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[B_4(531), \varepsilon']$	$1, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[2A_2, 1]$	$1, \bar{2}$	$0, \bar{1}$	$1, \bar{2}$	$1, \bar{2}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[A_1A_3, 1]$	$2, \bar{2}$	$1, \bar{2}$	$1, \bar{2}$	$1, \bar{1}$	$0, \bar{0}$	$1, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$

Table 13: Parahoric restriction of non-Iwahori-spherical $F_4(F)$-representations with respect to K_4

	$\overline{\theta_1} = \left(\begin{smallmatrix} 0 & 1 & 2 & 3 \\ & 1 & & \end{smallmatrix}\right) \otimes \mathrm{St}$	$\theta_1 = \left(\begin{smallmatrix} 0 & 1 & 2 & 3 \\ & 1 & & \end{smallmatrix}\right) \otimes \mathbf{1}$	$\overline{\theta_2} = \left(\begin{smallmatrix} 0 & 1 & 3 \\ & - & \end{smallmatrix}\right) \otimes \mathrm{St}$	$\theta_2 = \left(\begin{smallmatrix} 0 & 1 & 3 \\ & - & \end{smallmatrix}\right) \otimes \mathbf{1}$
$[B_4, -]$	1	0	0	0
$[C_3A_1, -]$	0	1	0	0
$[C_3(42)A_1, -+]$	0	0	1	0
$[B_4(531), r]$	1	1	1	0
$[A_1A_3, -]$	1	1	0	0

Table 14: Parahoric restrictions of the virtual characters $\Pi(u,s,h)$ with respect to K_4

	$[1,2]$	$[11,1]$	$[\cdot,1^3]$	$[1,11]$	$[1^3,\cdot]$	$[\cdot,21]$	θ_1	$[2,1]$	$[21,\cdot]$	$[\cdot,3]$	θ_2
$\Pi(F_4,1,1)$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(F_4(a_1),1,1)$	$0,\bar{0}$	$0,\bar{0}$	$1,\bar{1}$	$0,\bar{1}$	$0,\bar{0}$	$0,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(F_4(a_1),1,g_2)$	$0,\bar{0}$	$0,\bar{0}$	$1,\bar{1}$	$0,\bar{1}$	$0,\bar{0}$	$0,-\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(F_4(a_1),g_2,1)$	$0,\bar{0}$	$0,\bar{0}$	$1,\bar{1}$	$0,\bar{0}$	$0,\bar{1}$	$0,\bar{0}$	$0,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(F_4(a_1),g_2,g_2)$	$0,\bar{0}$	$0,\bar{0}$	$1,\bar{1}$	$0,\bar{0}$	$0,\bar{1}$	$0,\bar{0}$	$0,-\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(F_4(a_2),1,1)$	$0,\bar{0}$	$0,\bar{1}$	$1,\bar{3}$	$1,\bar{1}$	$1,\bar{0}$	$0,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(F_4(a_2),1,g_2)$	$0,\bar{0}$	$0,\bar{1}$	$1,\bar{1}$	$1,\bar{1}$	$-1,\bar{0}$	$0,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(F_4(a_2),g_2,1)$	$0,\bar{0}$	$0,\bar{1}$	$1,\bar{1}$	$0,\bar{1}$	$0,\bar{0}$	$1,\bar{1}$	$1,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(F_4(a_2),g_2,g_2)$	$0,\bar{0}$	$0,\bar{0}$	$1,\bar{1}$	$0,\bar{1}$	$0,\bar{0}$	$1,\bar{1}$	$-1,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(B_4(711),s,h)$	$0,\bar{0}$	$-1,\bar{1}$	$2,\bar{1}$	$1,\bar{0}$	$1,\bar{1}$	$0,-\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(F_4(a_3),1,1)$	$4,\bar{4}$	$1,\bar{2}$	$4,\bar{2}$	$4,\bar{8}$	$0,\bar{1}$	$4,\bar{7}$	$0,\bar{0}$	$0,\bar{6}$	$0,\bar{0}$	$0,\bar{6}$	$0,\bar{0}$
$\Pi(F_4(a_3),1,g_2)$	$2,\bar{2}$	$1,\bar{2}$	$2,\bar{2}$	$2,\bar{4}$	$0,\bar{1}$	$2,\bar{3}$	$0,\bar{0}$	$0,\bar{2}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(F_4(a_3),1,g_2')$	$0,\bar{0}$	$1,\bar{2}$	$4,\bar{2}$	$4,\bar{4}$	$0,\bar{1}$	$0,-\bar{1}$	$0,\bar{0}$	$0,\bar{2}$	$0,\bar{0}$	$0,-\bar{2}$	$0,\bar{0}$
$\Pi(F_4(a_3),1,g_3)$	$1,\bar{1}$	$1,\bar{2}$	$1,\bar{2}$	$1,\bar{2}$	$0,\bar{1}$	$1,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(F_4(a_3),1,g_4)$	$0,\bar{0}$	$1,\bar{2}$	$2,\bar{2}$	$2,\bar{2}$	$0,\bar{1}$	$0,-\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(F_4(a_3),g_2,1)$	$2,\bar{2}$	$1,\bar{2}$	$2,\bar{2}$	$2,\bar{4}$	$0,\bar{1}$	$2,\bar{3}$	$0,\bar{0}$	$0,\bar{1}$	$0,\bar{1}$	$0,\bar{1}$	$0,\bar{1}$
$\Pi(F_4(a_3),g_2,g_2)$	$2,\bar{2}$	$1,\bar{2}$	$2,\bar{2}$	$2,\bar{4}$	$0,\bar{1}$	$2,\bar{3}$	$0,\bar{0}$	$0,\bar{1}$	$0,\bar{1}$	$0,\bar{1}$	$0,-\bar{1}$
$\Pi(F_4(a_3),g_2,\tau)$	$0,\bar{0}$	$1,\bar{2}$	$2,\bar{2}$	$2,\bar{2}$	$0,\bar{1}$	$2,\bar{1}$	$0,\bar{0}$	$0,\bar{1}$	$0,\bar{1}$	$0,-\bar{1}$	$0,\bar{1}$
$\Pi(F_4(a_3),g_2,g_2')$	$0,\bar{0}$	$1,\bar{2}$	$2,\bar{2}$	$2,\bar{2}$	$0,\bar{1}$	$2,\bar{1}$	$0,\bar{0}$	$0,\bar{1}$	$0,\bar{1}$	$0,-\bar{1}$	$0,-\bar{1}$
$\Pi(F_4(a_3),g_2',1)$	$0,\bar{0}$	$1,\bar{2}$	$4,\bar{2}$	$2,\bar{2}$	$2,\bar{3}$	$2,\bar{1}$	$2,\bar{2}$	$0,\bar{0}$	$0,\bar{2}$	$0,\bar{0}$	$0,\bar{2}$
$\Pi(F_4(a_3),g_2',g_2)$	$0,\bar{0}$	$1,\bar{2}$	$2,\bar{2}$	$2,\bar{2}$	$0,\bar{1}$	$2,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{2}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(F_4(a_3),g_2',\gamma)$	$0,\bar{0}$	$1,\bar{2}$	$4,\bar{2}$	$2,\bar{2}$	$2,\bar{3}$	$0,-\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(F_4(a_3),g_2',g_2')$	$0,\bar{0}$	$1,\bar{2}$	$4,\bar{2}$	$2,\bar{2}$	$2,\bar{3}$	$2,\bar{1}$	$-2,-\bar{2}$	$0,\bar{0}$	$0,\bar{2}$	$0,\bar{0}$	$0,-\bar{2}$
$\Pi(F_4(a_3),g_2',g_4)$	$0,\bar{0}$	$1,\bar{2}$	$2,\bar{2}$	$2,\bar{2}$	$0,\bar{1}$	$0,-\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(F_4(a_3),g_3,h)$	$1,\bar{1}$	$1,\bar{2}$	$1,\bar{2}$	$1,\bar{2}$	$0,\bar{1}$	$1,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(F_4(a_3),g_4,1/g_2')$	$0,\bar{0}$	$1,\bar{2}$	$2,\bar{2}$	$1,\bar{1}$	$1,\bar{2}$	$1,\bar{0}$	$1,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(F_4(a_3),g_4,g_4^{\pm 1})$	$0,\bar{0}$	$1,\bar{2}$	$2,\bar{2}$	$1,\bar{1}$	$1,\bar{2}$	$1,\bar{0}$	$-1,-\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$

7.2 Parahoric restriction tables for E_6

Unipotent E_6-types

Table 15: Parahoric restriction of unipotent $E_6(F)_{\mathrm{sc}}$ with respect to K_0, K_1 and K_6, where the triality action of Ω can be easily observed

	$J_0 \longrightarrow E_6$	$J_1 \longrightarrow E_6$	$J_6 \longrightarrow E_6$
$[E_6, 1]$	$\phi_{1,36}$		
$[E_6(a_1), 1]$	$\phi_{1,36} + \phi_{6,25}$		
$[E_6(a_3), +]$	$\phi_{1,36} + \phi_{6,25} + \phi_{20,20} + \phi_{30,15}$		
$[E_6(a_3)_,]$	$\phi_{6,25} + \phi_{15,17}$		
$[A_1A_5, +]$	$\phi_{1,36} + \phi_{15,16} + \phi_{20,20}$		
$[A_1A_5, -]$	$D_4[\epsilon]$		
$[D_4(71), 1]$	$\phi_{1,36} + \phi_{15,17} + \phi_{20,20} + \phi_{24,12} + \phi_{30,15}$	$\phi_{6,25} + \phi_{20,20} + \phi_{64,13}$	$\phi_{6,25} + \phi_{20,20} + \phi_{64,13}$
$[D_4(71), \theta]$	$\phi_{6,25} + \phi_{20,20} + \phi_{64,13}$	$\phi_{6,25} + \phi_{20,20} + \phi_{64,13}$	$\phi_{1,36} + \phi_{15,17} + \phi_{20,20} + \phi_{24,12} + \phi_{30,15}$
$[D_4(71), \theta^2]$	$\phi_{6,25} + \phi_{20,20} + \phi_{64,13}$	$\phi_{1,36} + \phi_{15,17} + \phi_{20,20} + \phi_{24,12} + \phi_{30,15}$	$\phi_{6,25} + \phi_{20,20} + \phi_{64,13}$
$[D_4(a_1), 1]$	$\phi_{1,36} + \phi_{6,25} + \phi_{15,16} + 3\phi_{20,20} + \phi_{24,12} + 2\phi_{30,15} + \phi_{80,7} + 2\phi_{60,11} + 2\phi_{64,13} + \phi_{81,10}$		
$[D_4(a_1), r]$	$\phi_{6,25} + \phi_{15,17} + \phi_{20,20} + \phi_{30,15} + \phi_{90,8} + 2\phi_{64,13} + \phi_{81,10}$		
$[D_4(a_1), \epsilon]$	$\phi_{20,10} + \phi_{15,17}$		
$[D_4(53), 1]$	$\phi_{1,36} + \phi_{6,25} + \phi_{20,10} + \phi_{15,17} + \phi_{15,16} + \phi_{20,20} + \phi_{24,12} + 2\phi_{30,15} + \phi_{80,7} + 2\phi_{64,13} + \phi_{81,10}$	short	short
$[D_4(53), \theta]$	$\phi_{6,25} + \phi_{15,17} + 2\phi_{20,20} + \phi_{30,15} + \phi_{90,8} + \phi_{60,11} + 2\phi_{64,13} + \phi_{81,10}$	short	long
$[D_4(53), \theta^2]$	$\phi_{6,25} + \phi_{15,17} + 2\phi_{20,20} + \phi_{30,15} + \phi_{90,8} + \phi_{60,11} + 2\phi_{64,13} + \phi_{81,10}$	long	short
$[3A_2, 1 \boxtimes 1]$	$\phi_{1,36} + \phi_{10,9} + \phi_{15,16} + \phi_{24,12} + \phi_{30,15}$	$\phi_{20,20} + \phi_{60,11}$	$\phi_{20,20} + \phi_{60,11}$
$[3A_2, \theta \boxtimes 1]$	$E_6[\theta]$	0	0
$[3A_2, \theta^2 \boxtimes 1]$	$E_6[\theta^2]$	0	0
$[3A_2, 1 \boxtimes \theta]$	$\phi_{20,20} + \phi_{60,11}$	$\phi_{1,36} + \phi_{10,9} + \phi_{15,16} + \phi_{24,12} + \phi_{30,15}$	$\phi_{20,20} + \phi_{60,11}$
$[3A_2, \theta \boxtimes \theta]$	0	$E_6[\theta]$	0
$[3A_2, \theta^2 \boxtimes \theta]$	0	$E_6[\theta^2]$	0
$[3A_2, 1 \boxtimes \theta^2]$	$\phi_{20,20} + \phi_{60,11}$	$\phi_{20,20} + \phi_{60,11}$	$\phi_{1,36} + \phi_{10,9} + \phi_{15,16} + \phi_{24,12} + \phi_{30,15}$
$[3A_2, \theta \boxtimes \theta^2]$	0	0	$E_6[\theta]$
$[3A_2, \theta^2 \boxtimes \theta^2]$	0	0	$E_6[\theta^2]$
	$J_0 \longrightarrow E_6$		
$[D_4(3311), \mathbf{1} \boxtimes \mathbf{1}]$	$\phi_{1,36} + \phi_{10,9} + \phi_{6,25} + \phi_{20,10} + \phi_{15,17} + 2\phi_{15,16} + 2\phi_{20,20} + \phi_{24,6} + 3\phi_{24,12} + \phi_{30,3} + 4\phi_{30,15} + 2\phi_{60,8} + 3\phi_{80,7} + 3\phi_{90,8} + \phi_{60,5} + 3\phi_{60,11} + 4\phi_{64,13} + 2\phi_{81,6} + 4\phi_{81,10}$		
$[D_4(3311), \mathbf{1} \boxtimes \theta^{\pm 1}]$	$\phi_{6,25} + \phi_{15,17} + \phi_{15,16} + 3\phi_{20,20} + \phi_{24,12} + 2\phi_{30,15} + 2\phi_{60,8} + 3\phi_{80,7} + 3\phi_{90,8} + \phi_{60,5} + 4\phi_{60,11} + \phi_{64,4} + 5\phi_{64,13} + 2\phi_{81,6} + 4\phi_{81,10}$		
$[D_4(3311), \varepsilon \boxtimes \mathbf{1}]$	$\phi_{6,25} + \phi_{20,10} + \phi_{15,5} + \phi_{15,17} + \phi_{15,16} + \phi_{24,6} + \phi_{24,12} + \phi_{30,15} + \phi_{80,7} + 2\phi_{90,8} + \phi_{60,5} + \phi_{60,11} + 2\phi_{64,13} + \phi_{81,6} + 2\phi_{81,10}$		
$[D_4(3311), \varepsilon \boxtimes \theta^{\pm 1}]$	$\phi_{20,10} + \phi_{15,17} + \phi_{20,20} + \phi_{30,15} + \phi_{60,8} + \phi_{80,7} + 2\phi_{90,8} + \phi_{60,11} + \phi_{64,4} + 2\phi_{64,13} + \phi_{81,6} + 2\phi_{81,10}$		

Table 16: Parahoric restrictions of the virtual characters $\Pi(u,s,h)$ with respect to K_0

	$\phi_{80,7}$	$\phi_{90,8}$	$\phi_{20,10}$	$\phi_{60,8}$	$D_4[r]$	$\phi_{10,9}$	$E_6[\theta]$	$E_6[\theta^2]$	$\phi_{30,15}$	$\phi_{15,17}$	$\phi_{15,16}$	$D_4[\epsilon]$	$\phi_{30,3}$	$\phi_{15,5}$	$\phi_{15,4}$	$D_4[1]$	Fourier-stable
$\Pi(E_6,1,1)$	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$\phi_{1,36}$
$\Pi(E_6(a_1),1,1)$	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	$\phi_{1,36}+\phi_{6,25}$
$\Pi(E_6(a_3),1,1)$	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	$\Phi_{E_6(a_3)}$
$\Pi(E_6(a_3),1,g_2)$	0	0	0	0	0	0	0	0	1	-1	0	0	0	0	0	0	$\phi_{1,36}+\phi_{20,20}$
$\Pi(E_6(a_3),g_2,1)$	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	$\phi_{1,36}+\phi_{20,20}$
$\Pi(E_6(a_3),g_2,g_2)$	0	0	0	0	0	0	0	0	0	0	1	-1	0	0	0	0	$\phi_{1,36}+\phi_{20,20}$
$\Pi(D_4,s_3,w_3^{\pm 1})$	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	Φ_{D_4}
$\Pi(D_4(a_1),1,g_3)$	1	-1	1	0	0	0	0	0	1	0	1	0	0	0	0	0	$\Phi^1_{D_4(a_1)}$
$\Pi(D_4(a_1),z,g_3)$	1	-1	1	0	0	0	0	0	1	0	1	0	0	0	0	0	$\Phi^2_{D_4(a_1)}$
$\Pi(D_4(a_1),z^2,g_3)$	1	-1	1	0	0	0	0	0	1	0	1	0	0	0	0	0	$\Phi^2_{D_4(a_1)}$
$\Pi(D_4(a_1),g_3,1)$	0	0	0	0	0	1	1	1	1	0	1	0	0	0	0	0	$\Phi^1_{D_4(a_1)}$
$\Pi(D_4(a_1),g_3,z)$	0	0	0	0	0	1	1	1	1	0	1	0	0	0	0	0	$\Phi^2_{D_4(a_1)}$
$\Pi(D_4(a_1),g_3,z^2)$	0	0	0	0	0	1	1	1	1	0	1	0	0	0	0	0	$\Phi^2_{D_4(a_1)}$
$\Pi(D_4(a_1),g_3,g_3^{\pm 1})$	0	0	0	0	0	1	-1	-1	1	0	1	0	0	0	0	0	$\Phi^1_{D_4(a_1)}$
$\Pi(D_4(a_1),g_3,zg_3^{\pm 1})$	0	0	0	0	0	1	-1	-1	1	0	1	0	0	0	0	0	$\Phi^2_{D_4(a_1)}$
$\Pi(D_4(a_1),g_3,z^2g_3^{\pm 1})$	0	0	0	0	0	1	-1	-1	1	0	1	0	0	0	0	0	$\Phi^2_{D_4(a_1)}$
$\Pi(A_2,s_{3,3},w_{3,3}^{\pm 1})$	0	0	1	-1	0	1	0	0	2	0	2	0	1	1	0	0	Φ_{A_2}

Where, in the above table, all Φ_- are Fourier-stable combinations of the irreducible characters of $E_6(F)$ that are easily calculated from Table 15.

Unipotent $A_1 \times A_5$-types

Table 17: Parahoric restriction of Iwahori-spherical $E_6(F)$-representations with respect to K_2

V	$[1^6]$	$[21^4]$	$[2^21^2]$	$[222]$	$[31^3]$	$[321]$	$[33]$	$[41^2]$	$[42]$	$[51]$	$[6]$
$[E_6, 1]$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[E_6(a_1), 1]$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[E_6(a_3), +]$	$2, \bar{2}$	$1, \bar{3}$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[E_6(a_3), -]$	$1, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[A_1A_5, +]$	$1, \bar{2}$	$0, \bar{2}$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(71), 1]$	$1, \bar{2}$	$2, \bar{2}$	$1, \bar{2}$	$0, \bar{1}$	$1, \bar{2}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(71), \theta]$	$1, \bar{1}$	$2, \bar{3}$	$1, \bar{2}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(71), \theta^2]$	$1, \bar{1}$	$2, \bar{3}$	$1, \bar{2}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(a_1), 1]$	$3, \bar{5}$	$6, \bar{10}$	$7, \bar{10}$	$2, \bar{5}$	$4, \bar{6}$	$4, \bar{6}$	$1, \bar{1}$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(a_1), \epsilon]$	$0, \bar{0}$	$1, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(a_1), r]$	$2, \bar{1}$	$5, \bar{5}$	$4, \bar{4}$	$1, \bar{1}$	$4, \bar{6}$	$2, \bar{4}$	$0, \bar{0}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(53), 1]$	$3, \bar{3}$	$5, \bar{6}$	$5, \bar{6}$	$2, \bar{3}$	$5, \bar{7}$	$2, \bar{4}$	$1, \bar{1}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(53), \theta]$	$2, \bar{2}$	$6, \bar{7}$	$5, \bar{6}$	$1, \bar{2}$	$4, \bar{6}$	$3, \bar{5}$	$0, \bar{0}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(53), \theta^2]$	$2, \bar{2}$	$6, \bar{7}$	$5, \bar{6}$	$1, \bar{2}$	$4, \bar{6}$	$3, \bar{5}$	$0, \bar{0}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[3A_2, 1 \boxtimes 1]$	$1, \bar{2}$	$0, \bar{1}$	$1, \bar{2}$	$1, \bar{2}$	$1, \bar{1}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[3A_2, 1 \boxtimes \theta]$	$0, \bar{1}$	$1, \bar{2}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[3A_2, 1 \boxtimes \theta^2]$	$0, \bar{1}$	$1, \bar{2}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(3311), \mathbf{1} \boxtimes \mathbf{1}]$	$5, \bar{5}$	$11, \bar{14}$	$17, \bar{20}$	$9, \bar{11}$	$17, \bar{19}$	$18, \bar{22}$	$6, \bar{7}$	$9, \bar{11}$	$6, \bar{8}$	$2, \bar{2}$	$0, \bar{1}$
$[D_4(3311), \mathbf{1} \boxtimes \theta]$	$3, \bar{4}$	$13, \bar{15}$	$17, \bar{20}$	$7, \bar{9}$	$15, \bar{18}$	$20, \bar{24}$	$4, \bar{5}$	$9, \bar{10}$	$6, \bar{8}$	$2, \bar{3}$	$0, \bar{0}$
$[D_4(3311), \mathbf{1} \boxtimes \theta^2]$	$3, \bar{4}$	$13, \bar{15}$	$17, \bar{20}$	$7, \bar{9}$	$15, \bar{18}$	$20, \bar{24}$	$4, \bar{5}$	$9, \bar{10}$	$6, \bar{8}$	$2, \bar{3}$	$0, \bar{0}$
$[D_4(3311), \epsilon \boxtimes \mathbf{1}]$	$2, \bar{1}$	$5, \bar{5}$	$7, \bar{7}$	$4, \bar{3}$	$9, \bar{9}$	$8, \bar{10}$	$2, \bar{2}$	$5, \bar{7}$	$3, \bar{4}$	$1, \bar{2}$	$0, \bar{0}$
$[D_4(3311), \epsilon \boxtimes \theta]$	$1, \bar{1}$	$6, \bar{5}$	$7, \bar{7}$	$3, \bar{3}$	$8, \bar{9}$	$9, \bar{10}$	$1, \bar{2}$	$5, \bar{7}$	$3, \bar{4}$	$1, \bar{2}$	$0, \bar{0}$
$[D_4(3311), \epsilon \boxtimes \theta^2]$	$1, \bar{1}$	$6, \bar{5}$	$7, \bar{7}$	$3, \bar{3}$	$8, \bar{9}$	$9, \bar{10}$	$1, \bar{2}$	$5, \bar{7}$	$3, \bar{4}$	$1, \bar{2}$	$0, \bar{0}$

Table 18: Parahoric restriction of the virtual characters $\Pi(u,s,h)$ with respect to K_2

	$[1^6]$	$[21^4]$	$[2^21^2]$	$[222]$	$[31^3]$	$[321]$	$[33]$	$[41^2]$	$[42]$	$[51]$	$[6]$
$\Pi(E_6,1,1)$	$0,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(E_6(a_1),1,1)$	$1,\bar{1}$	$0,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(E_6(a_3),1,1)$	$3,\bar{2}$	$2,\bar{4}$	$1,\bar{1}$	$0,\bar{1}$	$0,\bar{2}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(E_6(a_3),1,g_2)$	$1,\bar{2}$	$0,\bar{2}$	$1,\bar{1}$	$0,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(E_6(a_3),g_2,1)$	$1,\bar{2}$	$0,\bar{2}$	$1,\bar{1}$	$0,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(E_6(a_3),g_2,g_2)$	$1,\bar{2}$	$0,\bar{2}$	$1,\bar{1}$	$0,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(D_4,s_3,w_3^{\pm1})$	$0,\bar{1}$	$0,-\bar{1}$	$0,\bar{0}$	$0,\bar{1}$	$0,\bar{1}$	$0,-\bar{1}$	$0,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(D_4(a_1),1,g_3)$	$1,\bar{4}$	$2,\bar{5}$	$3,\bar{6}$	$1,\bar{4}$	$1,\bar{1}$	$2,\bar{2}$	$1,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(D_4(a_1),z,g_3)$	$1,\bar{1}$	$-1,-\bar{1}$	$0,\bar{0}$	$1,\bar{1}$	$1,\bar{1}$	$-1,-\bar{1}$	$1,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(D_4(a_1),z^2,g_3)$	$1,\bar{1}$	$-1,-\bar{1}$	$0,\bar{0}$	$1,\bar{1}$	$1,\bar{1}$	$-1,-\bar{1}$	$1,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(D_4(a_1),g_3,1)$	$1,\bar{4}$	$2,\bar{5}$	$3,\bar{6}$	$1,\bar{4}$	$1,\bar{1}$	$2,\bar{2}$	$1,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(D_4(a_1),g_3,z)$	$1,\bar{1}$	$-1,-\bar{1}$	$0,\bar{0}$	$1,\bar{1}$	$1,\bar{1}$	$-1,-\bar{1}$	$1,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(D_4(a_1),g_3,z^2)$	$1,\bar{1}$	$-1,-\bar{1}$	$0,\bar{0}$	$1,\bar{1}$	$1,\bar{1}$	$-1,-\bar{1}$	$1,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(D_4(a_1),g_3,g_3^{\pm1})$	$1,\bar{4}$	$2,\bar{5}$	$3,\bar{6}$	$1,\bar{4}$	$1,\bar{1}$	$2,\bar{2}$	$1,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(D_4(a_1),g_3,zg_3^{\pm1})$	$1,\bar{1}$	$-1,-\bar{1}$	$0,\bar{0}$	$1,\bar{1}$	$1,\bar{1}$	$-1,-\bar{1}$	$1,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(D_4(a_1),g_3,z^2g_3^{\pm1})$	$1,\bar{1}$	$-1,-\bar{1}$	$0,\bar{0}$	$1,\bar{1}$	$1,\bar{1}$	$-1,-\bar{1}$	$1,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$\Pi(A_2,s_{3,3},w_{3,3}^{\pm1})$	$3,\bar{1}$	$-3,-\bar{1}$	$0,\bar{0}$	$3,\bar{2}$	$3,\bar{1}$	$-3,-\bar{2}$	$3,\bar{2}$	$0,\bar{1}$	$0,\bar{0}$	$0,-\bar{1}$	$0,\bar{1}$

Table 19: Parahoric restriction of Iwahori spherical $E_6(F)$-representations with respect to K_3

V	$[1^6]$	$[21^4]$	$[2^21^2]$	$[222]$	$[31^3]$	$[321]$	$[33]$	$[41^2]$	$[42]$	$[51]$	$[6]$
$[E_6, 1]$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[E_6(a_1), 1]$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[E_6(a_3), +]$	$2, \bar{2}$	$1, \bar{3}$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[E_6(a_3), -]$	$1, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[A_1A_5, +]$	$1, \bar{2}$	$0, \bar{2}$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(71), 1]$	$1, \bar{1}$	$2, \bar{3}$	$1, \bar{2}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(71), \theta]$	$1, \bar{1}$	$2, \bar{3}$	$1, \bar{2}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(71), \theta^2]$	$1, \bar{2}$	$2, \bar{2}$	$1, \bar{2}$	$0, \bar{1}$	$1, \bar{2}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(a_1), 1]$	$3, \bar{5}$	$6, \bar{10}$	$7, \bar{10}$	$2, \bar{5}$	$4, \bar{6}$	$4, \bar{6}$	$1, \bar{1}$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(a_1), \epsilon]$	$0, \bar{0}$	$1, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(a_1), r]$	$2, \bar{1}$	$5, \bar{5}$	$4, \bar{4}$	$1, \bar{1}$	$4, \bar{6}$	$2, \bar{4}$	$0, \bar{0}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(53), 1]$	$2, \bar{2}$	$6, \bar{7}$	$5, \bar{6}$	$1, \bar{2}$	$4, \bar{6}$	$3, \bar{5}$	$0, \bar{0}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(53), \theta]$	$2, \bar{2}$	$6, \bar{7}$	$5, \bar{6}$	$1, \bar{2}$	$4, \bar{6}$	$3, \bar{5}$	$0, \bar{0}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(53), \theta^2]$	$3, \bar{3}$	$5, \bar{6}$	$5, \bar{6}$	$2, \bar{3}$	$5, \bar{7}$	$2, \bar{4}$	$1, \bar{1}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[3A_2, 1 \boxtimes 1]$	$0, \bar{1}$	$1, \bar{2}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[3A_2, 1 \boxtimes \theta]$	$1, \bar{2}$	$0, \bar{1}$	$1, \bar{2}$	$1, \bar{2}$	$1, \bar{1}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[3A_2, 1 \boxtimes \theta^2]$	$0, \bar{1}$	$1, \bar{2}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(3311), \mathbf{1} \boxtimes \mathbf{1}]$	$3, \bar{4}$	$13, \bar{15}$	$17, \bar{20}$	$7, \bar{9}$	$15, \bar{18}$	$20, \bar{24}$	$4, \bar{5}$	$9, \bar{10}$	$6, \bar{8}$	$2, \bar{3}$	$0, \bar{0}$
$[D_4(3311), \mathbf{1} \boxtimes \theta]$	$3, \bar{4}$	$13, \bar{15}$	$17, \bar{20}$	$7, \bar{9}$	$15, \bar{18}$	$20, \bar{24}$	$4, \bar{5}$	$9, \bar{10}$	$6, \bar{8}$	$2, \bar{3}$	$0, \bar{0}$
$[D_4(3311), \mathbf{1} \boxtimes \theta^2]$	$5, \bar{5}$	$11, \bar{14}$	$17, \bar{20}$	$9, \bar{11}$	$17, \bar{19}$	$18, \bar{22}$	$6, \bar{7}$	$9, \bar{11}$	$6, \bar{8}$	$2, \bar{2}$	$0, \bar{1}$
$[D_4(3311), \epsilon \boxtimes \mathbf{1}]$	$1, \bar{1}$	$6, \bar{5}$	$7, \bar{7}$	$3, \bar{3}$	$8, \bar{9}$	$9, \bar{10}$	$1, \bar{2}$	$5, \bar{7}$	$3, \bar{4}$	$1, \bar{2}$	$0, \bar{0}$
$[D_4(3311), \epsilon \boxtimes \theta]$	$1, \bar{1}$	$6, \bar{5}$	$7, \bar{7}$	$3, \bar{3}$	$8, \bar{9}$	$9, \bar{10}$	$1, \bar{2}$	$5, \bar{7}$	$3, \bar{4}$	$1, \bar{2}$	$0, \bar{0}$
$[D_4(3311), \epsilon \boxtimes \theta^2]$	$2, \bar{1}$	$5, \bar{5}$	$7, \bar{7}$	$4, \bar{3}$	$9, \bar{9}$	$8, \bar{10}$	$2, \bar{2}$	$5, \bar{7}$	$3, \bar{4}$	$1, \bar{2}$	$0, \bar{0}$

Table 20: Parahoric restriction of Iwahori-spherical $E_6(F)$-representations with respect to K_5

V	$[1^6]$	$[21^4]$	$[2^21^2]$	$[222]$	$[31^3]$	$[321]$	$[33]$	$[41^2]$	$[42]$	$[51]$	$[6]$
$[E_6, 1]$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[E_6(a_1), 1]$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[E_6(a_3), +]$	$2, \bar{2}$	$1, \bar{3}$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[E_6(a_3), -]$	$1, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[A_1A_5, +]$	$1, \bar{2}$	$0, \bar{2}$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(71), 1]$	$1, \bar{1}$	$2, \bar{3}$	$1, \bar{2}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(71), \theta]$	$1, \bar{2}$	$2, \bar{2}$	$1, \bar{2}$	$0, \bar{1}$	$1, \bar{2}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(71), \theta^2]$	$1, \bar{1}$	$2, \bar{3}$	$1, \bar{2}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(a_1), 1]$	$3, \bar{5}$	$6, \bar{10}$	$7, \bar{10}$	$2, \bar{5}$	$4, \bar{6}$	$4, \bar{6}$	$1, \bar{1}$	$1, \bar{1}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(a_1), \epsilon]$	$0, \bar{0}$	$1, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(a_1), r]$	$2, \bar{1}$	$5, \bar{5}$	$4, \bar{4}$	$1, \bar{1}$	$4, \bar{6}$	$2, \bar{4}$	$0, \bar{0}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(53), 1]$	$2, \bar{2}$	$6, \bar{7}$	$5, \bar{6}$	$1, \bar{2}$	$4, \bar{6}$	$3, \bar{5}$	$0, \bar{0}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(53), \theta]$	$3, \bar{3}$	$5, \bar{6}$	$5, \bar{6}$	$2, \bar{3}$	$5, \bar{7}$	$2, \bar{4}$	$1, \bar{1}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(53), \theta^2]$	$2, \bar{2}$	$6, \bar{7}$	$5, \bar{6}$	$1, \bar{2}$	$4, \bar{6}$	$3, \bar{5}$	$0, \bar{0}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$
$[3A_2, 1 \boxtimes 1]$	$0, \bar{1}$	$1, \bar{2}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[3A_2, 1 \boxtimes \theta]$	$0, \bar{1}$	$1, \bar{2}$	$1, \bar{2}$	$0, \bar{1}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[3A_2, 1 \boxtimes \theta^2]$	$1, \bar{2}$	$0, \bar{1}$	$1, \bar{2}$	$1, \bar{2}$	$1, \bar{1}$	$0, \bar{0}$	$1, \bar{1}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$	$0, \bar{0}$
$[D_4(3311), \mathbf{1} \boxtimes \mathbf{1}]$	$3, \bar{4}$	$13, \bar{15}$	$17, \bar{20}$	$7, \bar{9}$	$15, \bar{18}$	$20, \bar{24}$	$4, \bar{5}$	$9, \bar{10}$	$6, \bar{8}$	$2, \bar{3}$	$0, \bar{0}$
$[D_4(3311), \mathbf{1} \boxtimes \theta]$	$5, \bar{5}$	$11, \bar{14}$	$17, \bar{20}$	$9, \bar{11}$	$17, \bar{19}$	$18, \bar{22}$	$6, \bar{7}$	$9, \bar{11}$	$6, \bar{8}$	$2, \bar{2}$	$0, \bar{1}$
$[D_4(3311), \mathbf{1} \boxtimes \theta^2]$	$3, \bar{4}$	$13, \bar{15}$	$17, \bar{20}$	$7, \bar{9}$	$15, \bar{18}$	$20, \bar{24}$	$4, \bar{5}$	$9, \bar{10}$	$6, \bar{8}$	$2, \bar{3}$	$0, \bar{0}$
$[D_4(3311), \epsilon \boxtimes \mathbf{1}]$	$1, \bar{1}$	$6, \bar{5}$	$7, \bar{7}$	$3, \bar{3}$	$8, \bar{9}$	$9, \bar{10}$	$1, \bar{2}$	$5, \bar{7}$	$3, \bar{4}$	$1, \bar{2}$	$0, \bar{0}$
$[D_4(3311), \epsilon \boxtimes \theta]$	$2, \bar{1}$	$5, \bar{5}$	$7, \bar{7}$	$4, \bar{3}$	$9, \bar{9}$	$8, \bar{10}$	$2, \bar{2}$	$5, \bar{7}$	$3, \bar{4}$	$1, \bar{2}$	$0, \bar{0}$
$[D_4(3311), \epsilon \boxtimes \theta^2]$	$1, \bar{1}$	$6, \bar{5}$	$7, \bar{7}$	$3, \bar{3}$	$8, \bar{9}$	$9, \bar{10}$	$1, \bar{2}$	$5, \bar{7}$	$3, \bar{4}$	$1, \bar{2}$	$0, \bar{0}$

Unipotent $A_2 \times A_2 \times A_2$-types

Table 21: Parahoric restriction of Iwahori-spherical $E_6(F)$-representations with respect to K_4

V	$[\varepsilon\varepsilon\varepsilon]$	$[\varepsilon\varepsilon r]$	$[\varepsilon\varepsilon 1]$	$[\varepsilon rr]$	$[\varepsilon r1]$	$[\varepsilon 11]$	$[rrr]$	$[rr1]$	$[r11]$	$[111]$
$[E_6, 1]$	1	0	0	0	0	0	0	0	0	0
$[E_6(a_1), 1]$	1	1	0	0	0	0	0	0	0	0
$[E_6(a_3), +]$	4	3	1	2	0	0	1	0	0	0
$[E_6(a_3), -]$	0	1	1	1	0	0	0	0	0	0
$[A_1A_5, +]$	4	2	0	1	0	0	1	0	0	0
$[D_4(71), 1]$	3	3,4,3	1,0,1	3,4,3	1,0,1,0,1,1	0,1,0	2	0	0	0
$[D_4(71), \theta]$	3	4,3,3	0,1,1	3,3,4	0,1,0,1,1,1	0,0,1	2	0	0	0
$[D_4(71), \theta^2]$	3	3,3,4	1,1,0	4,3,3	1,1,1,1,0,0	1,0,0	2	0	0	0
$[D_4(a_1), 1]$	15	15	3	17	4	1	19	4	1	0
$[D_4(a_1), \epsilon]$	0	0	1	1	1	0	1	0	0	0
$[D_4(a_1), r]$	3	8	4	11	4	1	11	3	0	0
$[D_4(53), 1]$	7	10,11,10	4	13,14,13	5,4,4,4,4,5	1	14	3,4,3	0,1,0	0
$[D_4(53), \theta]$	7	10,10,11	4	14,13,13	4,4,5,5,4,4	1	14	3,3,4	1,0,0	0
$[D_4(53), \theta^2]$	7	11,10,10	4	13,13,14	4,5,4,4,5,4	1	14	4,3,3	0,0,1	0
$[3A_2, 1 \boxtimes 1]$	4	2,3,2	0	2,3,2	1,0,0,0,0,1	0	3	0,1,0	0,1,0	0
$[3A_2, 1 \boxtimes \theta]$	4	2,2,3	0	3,2,2	0,0,1,1,0,0	0	3	0,0,1	1,0,0	0
$[3A_2, 1 \boxtimes \theta^2]$	4	3,2,2	0	2,2,3	0,1,0,0,1,0	0	3	1,0,0	0,0,1	0
$[D_4(3311), \mathbf{1} \boxtimes \mathbf{1}]$	19	29,31,29	11	46,48,46	19,17,17,17,17,19	4	72	26,28,26	9,11,9	3
$[D_4(3311), \mathbf{1} \boxtimes \theta]$	19	31,29,29	11	46,46,48	17,19,17,17,19,17	4	72	28,26,26	9,9,11	3
$[D_4(3311), \mathbf{1} \boxtimes \theta^2]$	19	29,29,31	11	48,46,46	17,17,19,19,17,17	4	72	26,26,28	11,9,9	3
$[D_4(3311), \epsilon \boxtimes \mathbf{1}]$	5	11	6,7,6	20	10,9,10,9,10,10	4,3,4	32	13,14,13	4,5,4	1
$[D_4(3311), \epsilon \boxtimes \theta]$	5	11	7,6,6	20	9,10,9,10,10,10	4,4,3	32	14,13,13	4,4,5	1
$[D_4(3311), \epsilon \boxtimes \theta^2]$	5	11	6,6,7	20	10,10,10,10,9,9	3,4,4	32	13,13,14	5,4,4	1

Table 22: Parahoric restriction of the virtual characters $\Pi(u,s,h)$ with respect to K_4

Row Label	$[\varepsilon\varepsilon\varepsilon]$	$[\varepsilon\varepsilon r]$	$[\varepsilon\varepsilon 1]$	$[\varepsilon rr]$	$[\varepsilon r1]$	$[\varepsilon 11]$	$[rrr]$	$[rr1]$	$[r11]$	$[111]$
$\Pi(E_6,1,1)$	1	0	0	0	0	0	0	0	0	0
$\Pi(E_6(a_1),1,1)$	1	1	0	0	0	0	0	0	0	0
$\Pi(E_6(a_3),1,1)$	4	4	2	3	0	0	1	0	0	0
$\Pi(E_6(a_3),1,g_2)$	4	2	0	1	0	0	1	0	0	0
$\Pi(E_6(a_3),g_2,1)$	4	2	0	1	0	0	1	0	0	0
$\Pi(E_6(a_3),g_2,g_2)$	4	2	0	1	0	0	1	0	0	0
$\Pi(D_4,s_3,w_3)$	0	$\theta,1,\theta^2$	$-\theta,-1,-\theta^2$	$\theta^2,1,\theta$	$-\theta,-1,-\theta,-1,-\theta^2,-\theta^2$	$\theta^2,1,\theta$	0	0	0	0
$\Pi(D_4,s_3,w_3^{-1})$	0	$\theta^2,1,\theta$	$-\theta^2,-1,-\theta$	$\theta,1,\theta^2$	$-\theta^2,-1,-\theta^2,-1,-\theta,-\theta$	$\theta,1,\theta^2$	0	0	0	0
$\Pi(D_4(a_1),1,g_3)$	12	7	0	7	1	0	9	1	1	0
$\Pi(D_4(a_1),z,g_3)$	0	$\theta^2,1,\theta$	0	$\theta,1,\theta^2$	$1,\theta^2,\theta,\theta,\theta^2,1$	0	0	$\theta^2,1,\theta$	$\theta,1,\theta^2$	0
$\Pi(D_4(a_1),z^2,g_3)$	0	$\theta,1,\theta^2$	0	$\theta^2,1,\theta$	$1,\theta,\theta^2,\theta^2,\theta,1$	0	0	$\theta,1,\theta^2$	$\theta^2,1,\theta$	0
$\Pi(D_4(a_1),g_3,1)$	12	7	0	7	1	0	9	1	1	0
$\Pi(D_4(a_1),g_3,z)$	0	$\theta^2,1,\theta$	0	$\theta,1,\theta^2$	$1,\theta^2,\theta,\theta,\theta^2,1$	0	0	$\theta^2,1,\theta$	$\theta,1,\theta^2$	0
$\Pi(D_4(a_1),g_3,z^2)$	0	$\theta,1,\theta^2$	0	$\theta^2,1,\theta$	$1,\theta,\theta^2,\theta^2,\theta,1$	0	0	$\theta,1,\theta^2$	$\theta^2,1,\theta$	0
$\Pi(A_2,s_{3,3},w_{3,3})$	0	$2\theta,2,2\theta^2$	$\theta,1,\theta^2$	$2\theta^2,2,2\theta$	$2-\theta,2\theta-1,2\theta^2-\theta,2\theta^2-1,2\theta-\theta^2,2-\theta^2$	$-\theta^2,-1,-\theta$	0	$3\theta,3,3\theta^2$	$3\theta^2,3,3\theta$	0
$\Pi(A_2,s_{3,3},w_{3,3}^{-1})$	0	$2\theta^2,2,2\theta$	$\theta^2,1,\theta$	$2\theta,2,2\theta^2$	$2-\theta^2,2\theta^2-1,2\theta-\theta^2,2\theta-1,2\theta^2-\theta,2-\theta$	$-\theta,-1,-\theta^2$	0	$3\theta^2,3,3\theta$	$3\theta,3,3\theta^2$	0

V	$K_0 \to \mathrm{Sp}_6(\mathbb{F}_q)$	$K_1 \to \mathrm{Sp}_2(\mathbb{F}_q) \times \mathrm{Sp}_4(\mathbb{F}_q)$	$K_1 \to \mathrm{Sp}_4(\mathbb{F}_q) \times \mathrm{Sp}_2(\mathbb{F}_q)$	$K_3 \to \mathrm{Sp}_6(\mathbb{F}_q)$
$[(511), 1]$	$[\mathbf{1}, \mathbf{1^2}] + [-, 1^3]$	$[-,1] \boxtimes ([\mathbf{1},\mathbf{1}] + [-,1^2]) + [1,-] \boxtimes [-,1^2]$	$([\mathbf{1},\mathbf{1}] + [-,1^2]) \boxtimes [-,1] + [-,1^2] \boxtimes [1,-]$	$[\mathbf{1}, \mathbf{1^2}] + [-, 1^3]$
$[(511), \varepsilon]$	$[-, 21]$	$[-,1] \boxtimes ([-,2] + [-,1^2])$	$([-,2] + [-,1^2]) \boxtimes [-,1]$	$[-, 21]$
$[(5), 1]$	$[-, 21] + [-, 1^3]$	$[-,1] \boxtimes ([-,2] + [-,1^2]) + [1,-] \boxtimes [-,1^2]$	$([\mathbf{1},\mathbf{1}] + [-,1^2]) \boxtimes [-,1]$	$[\mathbf{1}, \mathbf{1^2}]$
$[(5), \varepsilon]$	$[\mathbf{1}, \mathbf{1^2}]$	$[-,1] \boxtimes ([\mathbf{1},\mathbf{1}] + [-,1^2])$	$([-,2] + [-,1^2]) \boxtimes [-,1] + [-,1^2] \boxtimes [1,-]$	$[-, 21] + [-, 1^3]$
$[(4), 1 \boxtimes 1]$	$[-, 1^3]$	$[1,-] \boxtimes [-,1^2]$	$[1^2,-] \boxtimes [-,1]$	$[1^3,-]$
$[(4), 1 \boxtimes -1]$	$[1^3,-]$	$[-,1] \boxtimes [1^2,-]$	$[-,1^2] \boxtimes [1,-]$	$[-, 1^3]$
$[(4), -1 \boxtimes 1]$	0	0	$\theta'_2 \boxtimes [-,1]$	θ'_3
$[(4), -1 \boxtimes -1]$	θ'_3	$[-,1] \boxtimes \theta'_2$	0	0
$[(331), 1]$	$[\mathbf{1}, \mathbf{1^2}] + [1^2, 1] + [-, 1^3]$	$[-,1] \boxtimes ([\mathbf{1},\mathbf{1}] + [1^2,-] + [-,1^2]) + [1,-] \boxtimes ([\mathbf{1},\mathbf{1}] + [-,1^2])$	$([\mathbf{1},\mathbf{1}] + [1^2,-] + [-,1^2]) \boxtimes [-,1] + ([\mathbf{1},\mathbf{1}] + [-,1^2]) \boxtimes [1,-]$	$[\mathbf{1}, \mathbf{1^2}] + [1^2, 1] + [-, 1^3]$
$[(331), \varepsilon]$	$[1^3,-]$	$[1,-] \boxtimes [1^2,-]$	$[1^2,-] \boxtimes [1,-]$	$[1^3,-]$
$[(31), 1]$	$[\mathbf{1}, \mathbf{1^2}] + [-, 1^3]$	$[-,1] \boxtimes [-,1^2] + [1,-] \boxtimes ([\mathbf{1},\mathbf{1}] + [-,1^2])$	$([\mathbf{1},\mathbf{1}] + [1^2,-]) \boxtimes [-,1] + [1^2,-] \boxtimes [1,-]$	$[1^3,-] + [1^2, 1]$
$[(31), \varepsilon]$	$[1^3,-] + [1^2, 1]$	$[-,1] \boxtimes ([\mathbf{1},\mathbf{1}] + [1^2,-]) + [1,-] \boxtimes [1^2,-]$	$[-,1^2] \boxtimes [-,1] + ([\mathbf{1},\mathbf{1}] + [-,1^2]) \boxtimes [1,-]$	$[\mathbf{1}, \mathbf{1^2}] + [-, 1^3]$
$[(2,2,2), 1 \boxtimes 1]$	$[-, 21] + [-, 1^3]$	$[-,1] \boxtimes [-,1^2] + [1,-] \boxtimes ([-,2] + [-,1^2])$	$([\mathbf{1},\mathbf{1}] + [1^2,-]) \boxtimes [-,1]$	$[1^2, 1]$
$[(2,2,2), 1 \boxtimes -1]$	$[1^2, 1]$	$[-,1] \boxtimes ([\mathbf{1},\mathbf{1}] + [1^2,-])$	$[-,1^2] \boxtimes [-,1] + ([-,2] + [-,1^2]) \boxtimes [1,-]$	$[-, 21] + [-, 1^3]$
$[(2,2,2), -1 \boxtimes 1]$	θ'_3	$[1,-] \boxtimes \theta'_2$	0	0
$[(2,2,2), -1 \boxtimes -1]$	0	0	$\theta'_2 \boxtimes [1,-]$	θ'_3

Table 23: Parahoric restrictions of unipotent representations for $Sp_6(\mathbb{Q}_p)$. Red indicates representations in non-trivial Lusztig families, with bold denoting special representations.

V	$K_0 \to \mathrm{Sp}_6(\mathbb{F}_q)$	$K_1 \to \mathrm{Sp}_2(\mathbb{F}_q) \times \mathrm{Sp}_4(\mathbb{F}_q)$	$K_1 \to \mathrm{Sp}_4(\mathbb{F}_q) \times \mathrm{Sp}_2(\mathbb{F}_q)$	$K_3 \to \mathrm{Sp}_6(\mathbb{F}_q)$
$\Pi((511), 1, \delta)$	$[\mathbf{1}, \mathbf{1^2}] - [-, 21] + [-, 1^3]$	$[-, 1] \boxtimes ([\mathbf{1}, \mathbf{1}] - [-, 2]) + [1, -] \boxtimes [-, 1^2]$	$([\mathbf{1}, \mathbf{1}] - [-, 2]) \boxtimes [-, 1] + [-, 1^2] \boxtimes [1, -]$	$[\mathbf{1}, \mathbf{1^2}] - [-, 21] + [-, 1^3]$
$\Pi((511), \delta, 1)$	$[1^3, -] + \theta'_3 + [-, 1^3]$	$[-, 1] \boxtimes ([1^2, -] + \theta'_2) + [1, -] \boxtimes [-, 1^2]$	$([1^2, -] + \theta'_2) \boxtimes [-, 1] + [-, 1^2] \boxtimes [1, -]$	$[1^3, -] + \theta'_3 + [-, 1^3]$
$\Pi((511), -1, \delta)$	$-[\mathbf{1}, \mathbf{1^2}] + [-, 21] + [-, 1^3]$	$[-, 1] \boxtimes (-[\mathbf{1}, \mathbf{1}] + [-, 2]) + [1, -] \boxtimes [-, 1^2]$	$([\mathbf{1}, \mathbf{1}] - [-, 2]) \boxtimes [-, 1] - [-, 1^2] \boxtimes [1, -]$	$[\mathbf{1}, \mathbf{1^2}] - [-, 21] - [-, 1^3]$
$\Pi((511), \delta, -1)$	$-[1^3, -] - \theta'_3 + [-, 1^3]$	$[-, 1] \boxtimes (-[1^2, -] - \theta'_2) + [1, -] \boxtimes [-, 1^2]$	$([1^2, -] + \theta'_2) \boxtimes [-, 1] - [-, 1^2] \boxtimes [1, -]$	$[1^3, -] + \theta'_3 - [-, 1^3]$
$\Pi((511), \delta, \delta)$	$[1^3, -] - \theta'_3 + [-, 1^3]$	$[-, 1] \boxtimes ([1^2, -] - \theta'_2) + [1, -] \boxtimes [-, 1^2]$	$([1^2, -] - \theta'_2) \boxtimes [-, 1] + [-, 1^2] \boxtimes [1, -]$	$[1^3, -] - \theta'_3 + [-, 1^3]$
$\Pi((511), \delta, -\delta)$	$-[1^3, -] + \theta'_3 + [-, 1^3]$	$[-, 1] \boxtimes (-[1^2, -] + \theta'_2) + [1, -] \boxtimes [-, 1^2]$	$([1^2, -] - \theta'_2) \boxtimes [-, 1] - [-, 1^2] \boxtimes [1, -]$	$-[1^3, -] - \theta'_3 - [-, 1^3]$
$\Pi((331), 1, \delta)$	$[\mathbf{1}, \mathbf{1^2}] - [1^3, -] + [1^2, 1] + [-, 1^3]$	$[-, 1] \boxtimes ([\mathbf{1}, \mathbf{1}] + [1^2, -] + [-, 1^2]) + [1, -] \boxtimes ([\mathbf{1}, \mathbf{1}] - [1^2, -] + [-, 1^2])$	$([\mathbf{1}, \mathbf{1}] + [1^2, -] + [-, 1^2]) \boxtimes [-, 1] + ([\mathbf{1}, \mathbf{1}] - [1^2, -] + [-, 1^2]) \boxtimes [1, -]$	$[\mathbf{1}, \mathbf{1^2}] - [1^3, -] + [1^2, 1] + [-, 1^3]$
$\Pi((331), \delta, 1)$	$[-, 21] + \theta'_3 + [1^2, 1] + [-, 1^3]$	$[-, 1] \boxtimes ([\mathbf{1}, \mathbf{1}] + [1^2, -] + [-, 1^2]) + [1, -] \boxtimes ([-, 2] + \theta'_2 + [-, 1^2])$	$([\mathbf{1}, \mathbf{1}] + [1^2, -] + [-, 1^2]) \boxtimes [-, 1] + ([-, 2] + \theta'_2 + [-, 1^2]) \boxtimes [1, -]$	$[-, 21] + \theta'_3 + [1^2, 1] + [-, 1^3]$
$\Pi((331), -1, \delta)$	$[\mathbf{1}, \mathbf{1^2}] - [1^3, -] - [1^2, 1] + [-, 1^3]$	$[-, 1] \boxtimes (-[\mathbf{1}, \mathbf{1}] - [1^2, -] + [-, 1^2]) + [1, -] \boxtimes ([\mathbf{1}, \mathbf{1}] - [1^2, -] + [-, 1^2])$	$([\mathbf{1}, \mathbf{1}] + [1^2, -] - [-, 1^2]) \boxtimes [-, 1] + (-[\mathbf{1}, \mathbf{1}] + [1^2, -] - [-, 1^2]) \boxtimes [1, -]$	$-[\mathbf{1}, \mathbf{1^2}] + [1^3, -] + [1^2, 1] - [-, 1^3]$
$\Pi((331), \delta, -1)$	$[-, 21] + \theta'_3 - [1^2, 1] + [-, 1^3]$	$[-, 1] \boxtimes (-[\mathbf{1}, \mathbf{1}] - [1^2, -] + [-, 1^2]) + [1, -] \boxtimes ([-, 2] + \theta'_2 + [-, 1^2])$	$([\mathbf{1}, \mathbf{1}] + [1^2, -] - [-, 1^2]) \boxtimes [-, 1] + (-[-, 2] - \theta'_2 - [-, 1^2]) \boxtimes [1, -]$	$-[-, 21] - \theta'_3 + [1^2, 1] - [-, 1^3]$
$\Pi((331), \delta, \delta)$	$[-, 21] - \theta'_3 + [1^2, 1] + [-, 1^3]$	$[-, 1] \boxtimes ([\mathbf{1}, \mathbf{1}] + [1^2, -] + [-, 1^2]) + [1, -] \boxtimes ([-, 2] - \theta'_2 + [-, 1^2])$	$([\mathbf{1}, \mathbf{1}] + [1^2, -] + [-, 1^2]) \boxtimes [-, 1] + ([-, 2] - \theta'_2 + [-, 1^2]) \boxtimes [1, -]$	$[-, 21] - \theta'_3 + [1^2, 1] - [-, 1^3]$
$\Pi((331), \delta, -\delta)$	$[-, 21] - \theta'_3 - [1^2, 1] + [-, 1^3]$	$[-, 1] \boxtimes (-[\mathbf{1}, \mathbf{1}] - [1^2, -] + [-, 1^2]) + [1, -] \boxtimes ([-, 2] - \theta'_2 + [-, 1^2])$	$([\mathbf{1}, \mathbf{1}] + [1^2, -] - [-, 1^2]) \boxtimes [-, 1] + (-[-, 2] + \theta'_2 - [-, 1^2]) \boxtimes [1, -]$	$-[-, 21] + \theta'_3 + [1^2, 1] - [-, 1^3]$

Table 24: Linear combinations of parahoric restrictions for unipotent representations of $\mathrm{Sp}_6(\mathbb{Q}_p)$. Red indicates representations in non-trivial Lusztig families, with bold denoting special representations.

	$J_0 \longrightarrow E_6$
$[[-,4],1]$	$\phi_{1,36}+\phi_{15,17}+\phi_{20,20}+\phi_{24,12}+\phi_{30,15}$ (90)
$[[-,4],\theta^k]$	$\phi_{6,25}+\phi_{20,20}+\phi_{64,13}$ (90)
$[[1,3],1]$	$\phi_{6,25}+\phi_{20,10}+\phi_{15,16}+\phi_{30,15}+\phi_{80,7}+2\phi_{64,13}+\phi_{81,10}$ (360)
$[[1,3],\theta^k]$	$\phi_{15,17}+\phi_{20,20}+\phi_{30,15}+\phi_{90,8}+\phi_{60,11}+\phi_{64,13}+\phi_{81,10}$ (360)
$[[11,2],1]$	$\phi_{15,5}+\phi_{15,17}+\phi_{24,6}+\phi_{24,12}+2\phi_{90,8}+\phi_{60,5}+\phi_{60,11}+\phi_{81,6}+\phi_{81,10}$ (540)
$[[11,2],\theta^k]$	$\phi_{20,10}+\phi_{60,8}+\phi_{80,7}+\phi_{90,8}+\phi_{64,4}+\phi_{64,13}+\phi_{81,6}+\phi_{81,10}$ (540)
$[[1,21],1]$	$\phi_{10,9}+\phi_{24,6}+\phi_{24,12}+\phi_{30,3}+\phi_{30,15}+\phi_{60,8}+\phi_{80,7}+2\phi_{90,8}+\phi_{60,5}+\phi_{60,11}+\phi_{81,6}+\phi_{81,10}$
$[[1,21],\varepsilon]$	$\phi_{60,8}+2\phi_{80,7}+\phi_{90,8}+\phi_{60,5}+\phi_{60,11}+\phi_{64,4}+\phi_{64,13}+\phi_{81,6}+\phi_{81,10}$
[3-dim reps,-]	$\phi_{15,16}+\phi_{20,20}+\phi_{24,12}+\phi_{30,15}+\phi_{60,8}+\phi_{80,7}+\phi_{90,8}+2\phi_{60,11}+2\phi_{64,13}+\phi_{81,6}+2\phi_{81,10}$

V	$[1^6]$	$[21^4]$	$[2^21^2]$	$[222]$	$[31^3]$	$[321]$	$[33]$	$[41^2]$	$[42]$	$[51]$	$[6]$
$[D_4(3311),1]$	$11,\bar{13}$	$37,\bar{44}$	$51,\bar{60}$	$23,\bar{29}$	$47,\bar{55}$	$58,\bar{70}$	$14,\bar{17}$	$27,\bar{31}$	$18,\bar{24}$	$6,\bar{8}$	$0,\bar{1}$
$[D_4(3311),\epsilon]$	$4,\bar{3}$	$17,\bar{15}$	$21,\bar{21}$	$10,\bar{9}$	$25,\bar{27}$	$26,\bar{30}$	$4,\bar{6}$	$15,\bar{21}$	$9,\bar{12}$	$3,\bar{6}$	$0,\bar{0}$
$[D_4(3311),\mathbf{1}\boxtimes\mathbf{1}]$	$5,\bar{5}$	$11,\bar{14}$	$17,\bar{20}$	$9,\bar{11}$	$17,\bar{19}$	$18,\bar{22}$	$6,\bar{7}$	$9,\bar{11}$	$6,\bar{8}$	$2,\bar{2}$	$0,\bar{1}$
$[D_4(3311),\mathbf{1}\boxtimes\theta^k]$	$3,\bar{4}$	$13,\bar{15}$	$17,\bar{20}$	$7,\bar{9}$	$15,\bar{18}$	$20,\bar{24}$	$4,\bar{5}$	$9,\bar{10}$	$6,\bar{8}$	$2,\bar{3}$	$0,\bar{0}$
$[D_4(3311),\epsilon\boxtimes\mathbf{1}]$	$2,\bar{1}$	$5,\bar{5}$	$7,\bar{7}$	$4,\bar{3}$	$9,\bar{9}$	$8,\bar{10}$	$2,\bar{2}$	$5,\bar{7}$	$3,\bar{4}$	$1,\bar{2}$	$0,\bar{0}$
$[D_4(3311),\epsilon\boxtimes\theta^k]$	$1,\bar{1}$	$6,\bar{5}$	$7,\bar{7}$	$3,\bar{3}$	$8,\bar{9}$	$9,\bar{10}$	$1,\bar{2}$	$5,\bar{7}$	$3,\bar{4}$	$1,\bar{2}$	$0,\bar{0}$
$[[-,4],1]$	$1,\bar{2}$	$2,\bar{2}$	$1,\bar{2}$	$0,\bar{1}$	$1,\bar{2}$	$0,\bar{0}$	$0,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$[[-,4],\theta^k]$	$1,\bar{1}$	$2,\bar{3}$	$1,\bar{2}$	$0,\bar{0}$	$1,\bar{1}$	$0,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$	$0,\bar{0}$
$[[1,3],1]$	$2,\bar{1}$	$3,\bar{4}$	$4,\bar{4}$	$2,\bar{2}$	$4,\bar{5}$	$2,\bar{4}$	$1,\bar{0}$	$1,\bar{2}$	$0,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$
$[[1,3],\theta^k]$	$1,\bar{1}$	$4,\bar{4}$	$4,\bar{4}$	$1,\bar{2}$	$3,\bar{5}$	$3,\bar{4}$	$0,\bar{0}$	$1,\bar{2}$	$0,\bar{1}$	$0,\bar{0}$	$0,\bar{0}$
$[[11,2],1]$	$0,\bar{0}$	$2,\bar{1}$	$3,\bar{3}$	$2,\bar{1}$	$5,\bar{4}$	$6,\bar{6}$	$1,\bar{2}$	$4,\bar{5}$	$3,\bar{3}$	$1,\bar{2}$	$0,\bar{0}$
$[[11,2],\theta^k]$	$0,\bar{0}$	$2,\bar{1}$	$3,\bar{3}$	$2,\bar{1}$	$5,\bar{4}$	$6,\bar{6}$	$1,\bar{2}$	$4,\bar{5}$	$3,\bar{3}$	$1,\bar{2}$	$0,\bar{0}$
$[[1,21],1]$	$1,\bar{0}$	$1,\bar{2}$	$5,\bar{4}$	$3,\bar{4}$	$5,\bar{5}$	$8,\bar{8}$	$4,\bar{3}$	$5,\bar{5}$	$4,\bar{5}$	$2,\bar{1}$	$0,\bar{1}$
$[[1,21],\theta^k]$	$0,\bar{0}$	$2,\bar{2}$	$5,\bar{4}$	$2,\bar{3}$	$4,\bar{5}$	$9,\bar{9}$	$3,\bar{2}$	$5,\bar{4}$	$4,\bar{5}$	$2,\bar{2}$	$0,\bar{0}$
[3-dim reps, $-$]	$1,\bar{2}$	$5,\bar{6}$	$7,\bar{10}$	$4,\bar{4}$	$7,\bar{7}$	$8,\bar{10}$	$1,\bar{3}$	$3,\bar{4}$	$2,\bar{2}$	$0,\bar{1}$	$0,\bar{0}$

References

[ACR25] Anne-Marie Aubert, Dan Ciubotaru, and Beth Romano, *A nonabelian Fourier transform for tempered unipotent representations*, Compos. Math. **161** (2025), no. 1, 13–73. MR4877237

[Alv05] Dean Alvis, *Induce/restrict matrices for exceptional weyl groups*, 2005. Preprint, arXiv:math/0506377 [math.RT].

[BK98] Colin J. Bushnell and Philip C. Kutzko, *Smooth representations of reductive p-adic groups: structure theory via types*, Proc. London Math. Soc. (3) **77** (1998), no. 3, 582–634. MR1643417

[Car85] Roger W. Carter, *Finite groups of lie type: Conjugacy classes and complex characters*, Wiley, Chichester, 1985. A Wiley-Interscience publication.

[Ciu20] Dan Ciubotaru, *The nonabelian fourier transform for elliptic unipotent representations of exceptional p-adic groups*, 2020.

[CMBO25] Dan Ciubotaru, Lucas Mason-Brown, and Emile Okada, *Wavefront sets of unipotent representations of reductive p-adic groups II*, J. Reine Angew. Math. **823** (2025), 191–253. MR4912263

[CO17] Dan Ciubotaru and Eric Opdam, *On the elliptic nonabelian Fourier transform for unipotent representations of p-adic groups*, Representation theory, number theory, and invariant theory, 2017, pp. 87–113. MR3753909

[Kat83] Shin-ichi Kato, *A realization of irreducible representations of affine weyl groups*, Indagationes Mathematicae (Proceedings) **86** (1983), no. 2, 193–201.

[KL87] David Kazhdan and George Lusztig, *Proof of the Deligne-Langlands conjecture for Hecke algebras*, Invent. Math. **87** (1987), no. 1, 153–215. MR862716

[Lus02] G. Lusztig, *Classification of unipotent representations of simple p-adic groups. II*, Represent. Theory **6** (2002), 243–289. MR1927955

[Lus84] George Lusztig, *Characters of reductive groups over a finite field*, Annals of Mathematics Studies, vol. 107, Princeton University Press, Princeton, NJ, 1984. MR742472

[Lus89] ———, *Affine Hecke algebras and their graded version*, J. Amer. Math. Soc. **2** (1989), no. 3, 599–635. MR991016

[Lus95a] ———, *Classification of unipotent representations of simple p-adic groups*, Internat. Math. Res. Notices **11** (1995), 517–589. MR1369407

[Lus95b] ———, *Cuspidal local systems and graded Hecke algebras. II*, Representations of groups (Banff, AB, 1994), 1995, pp. 217–275. With errata for Part I [Inst. Hautes Études Sci. Publ. Math. No. 67 (1988), 145–202; MR0972345 (90e:22029)]. MR1357201

[MW03] Colette Mœglin and Jean-Loup Waldspurger, *Paquets stables de représentations tempérées et de réduction unipotente pour* $\mathrm{SO}(2n+1)$, Invent. Math. **152** (2003), no. 3, 461–623. MR1988295

[Ree00] Mark Reeder, *Formal degrees and L-packets of unipotent discrete series representations of exceptional p-adic groups*, J. Reine Angew. Math. **520** (2000), 37–93. With an appendix by Frank Lübeck. MR1748271

[Ree02] ———, *Isogenies of Hecke algebras and a Langlands correspondence for ramified principal series representations*, Represent. Theory **6** (2002), 101–126. MR1915088

[Ree97] ———, *Nonstandard intertwining operators and the structure of unramified principal series representations*, Forum Math. **9** (1997), no. 4, 457–516. MR1457135

[Sol20] Maarten Solleveld, *Langlands parameters, functoriality and Hecke algebras*, Pacific J. Math. **304** (2020), no. 1, 209–302. MR4053201

[Sol21] ———, *Affine hecke algebras and their representations*, Indagationes Mathematicae **32** (2021), no. 5, 1005–1082.

[Sol23a] ———, *A local Langlands correspondence for unipotent representations*, Amer. J. Math. **145** (2023), no. 3, 673–719. MR4596175

[Sol23b] ———, *On unipotent representations of ramified p-adic groups*, Represent. Theory **27** (2023), 669–716. MR4620884

[Spr98] T. A. Springer, *Linear algebraic groups*, 2nd ed., Progress in Mathematics, vol. 9, Birkhäuser, Boston, MA, 1998.

[Wal18] Jean-Loup Waldspurger, *Représentations de réduction unipotente pour $SO(2n+1)$, I: une involution*, J. Lie Theory **28** (2018), no. 2, 381–426. MR3733069

8 Explicit parahoric restriction of $\pi(D_4(71), -)$ wrt K_4


For this explicit example, let $G = E_6(F)_{\mathrm{sc}}$ and $G^\vee = E_6(\mathbb{C})_{\mathrm{ad}}$ be the dual group. Fix a unipotent element $u \in G^\vee$ of type D_4. Under the natural map $\pi : G^\vee_{\mathrm{sc}} \to G^\vee$, the element u has a unique *unipotent* lift $\tilde{u}$, also of type D_4. With GAP, we see that $\Gamma_{\tilde{u}}$, the reductive part of $Z_{G^\vee_{\mathrm{sc}}}(\tilde{u})$, has type A_2. Since $Z(G^\vee_{\mathrm{sc}}) \subseteq Z(\Gamma_{\tilde{u}})$, it follows that $\Gamma_{\tilde{u}} \cong \mathrm{SL}_3(\mathbb{C})$. The reductive part Γ_u of $Z_{G^\vee}(u)$ is also of type A_2, and the since the kernel of the surjective map $\pi|_{\Gamma_{\tilde{u}}} : \Gamma_{\tilde{u}} \to \Gamma_u$ is $Z(\Gamma_{\tilde{u}})$, we have that $\Gamma_u \cong \mathrm{PGL}_3(\mathbb{C})$. In particular,

$$\Gamma_u \backslash \mathcal{Y}(\Gamma_u)_{\mathrm{ell}} = \{(s, w), (s, w^2)\}, \quad \text{where} \quad s = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \theta_3 & 0 \\ 0 & 0 & \theta_3^2 \end{pmatrix} \quad \text{and} \quad w = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$$

The aim is therefore to compute the parahoric restrictions with respect to maximal parahoric subgroups of the virtual representations

$$\Pi(u, s, w^i) = \sum_{\phi \in \widehat{A_{us}}} \phi(w)^i \pi(us, \phi), \ i = 1, 2.$$

To calculate the parahoric restrictions of $\pi(us, \phi)$ first we want to understand the semisimple element s inside $G^\vee$. It is an order 3 element commuting with a unipotent element of type D_4. The Kac coordinates or s are therefore either of



I believe there are a few different ways to show that s is in fact the first one. The easiest one I can think of uses the fact that $Z_{\Gamma_u}(s) = T_u \sqcup T_u w \sqcup T_u w^2 \cong T_u \rtimes C_3$, where T_u is the two dimensional split torus of $\Gamma_u \cong \mathrm{PGL}_3(\mathbb{C})$. The component group of $Z_{G^\vee}(su) = Z_{Z_{G^\vee}(u)}(s)$ is isomorphic to the component group of $Z_{\Gamma_u}(s)$, isomorphic to C_3 by the above observation. If the Kac coordinates of s were the latter option, then $Z_{G^\vee}(s)$ would be a connected reductive group of type D_5, and one can check with GAP that there is no unipotent element v inside that group such that the component group of $Z_{G^\vee(s)}(u)$ is C_3. Letting $x = su \in G^\vee$, we then have $A_x = C_3$ so the corresponding L-packet has size 3.

Therefore, we know that the Kac coordinates of s are the left ones, so $Z_{G^\vee}(s) = Z_{G^\vee}(s)^0 \rtimes C_3$, where $M^\vee := Z_{G^\vee}(s)^0$ is a connected reductive group of type D_4. In particular, the unipotent element u is regular in $M^\vee$, commonly labelled by the partition (71). Following Reeder's notation, we write $\pi(x, \phi) = \pi(D_4(71), \phi), \ \phi \in \{\mathbf{1}, \theta, \theta^2\}$ for the three representations inside the L-packet.

These three representations are all Iwahori-spherical, so to continue we use Reeder's reduction steps that show that for any $J \subsetneq S_{\text{aff}}$ (set of simple affine roots of $G^\vee$) and irreducible character χ of $\overline{K}_J$ (the reductive quotient), we have that

$$\langle \pi(us,\phi)^{U_J}, \chi \rangle_{\overline{K}_J} = \langle \pi(us,\phi)^{\mathcal{I}}_{q=1}, \chi^{\mathcal{I}}_{q=1} \rangle_{W_J}.$$

Moreover, Lusztig showed that $\pi(us,\phi)^{\mathcal{I}}_{q=1} \cong H(\mathcal{B}^x)^\phi \otimes \varepsilon$ as W_{aff}-modules, where $\mathcal{B}^x$ is the partial flag variety of Borel subgroups of $G^\vee$ containing x.

At this point, we use Kato's work to understand the structure of $H(\mathcal{B}^x)$ as a $W_{\text{aff}} \times A_x$-module. As a W_{aff}-module, it is straightforward:

$$H(\mathcal{B}^x) \cong \text{Ind}_{W_{M,\text{aff}}}^{W_{\text{aff}}} H(\mathcal{B}^u_s) \otimes s,$$

where $W_{M,\text{aff}} = W_M \ltimes \mathbb{Z}\Phi^\vee$, $\mathcal{B}^u_s$ is the partial flag variety of Borel subgroups of $M^\vee$ containing u and s is the character of $\mathbb{Z}\Phi^\vee$ given by $\alpha \mapsto \alpha(s), \alpha \in \Phi$. Since u is regular in $M^\vee$, $H(\mathcal{B}^u_s)$ is the trivial W_M-representation. Thus,

$$H(\mathcal{B}^x) \cong \text{Ind}_{W_{M,\text{aff}}}^{W_{\text{aff}}} s.$$

We now focus on the parahoric restriction of these representations with respect to the maximal parahoric subgroup K_4 of type $3A_2$ corresponding to $J_4 = S_{\text{aff}} \setminus \{\alpha_4\}$.

Using GAP, we can compute this directly, either following Reeder's method or by computing the action explicitly with matrices. Either way, and noting that $H(\mathcal{B}^x) = H(\mathcal{B}^x)^{\mathbf{1}} + H(\mathcal{B}^x)^\theta + H(\mathcal{B}^x)^{\theta^2}$, we obtain the following, where, for example $[\varepsilon\varepsilon r]$ represents the three representations obtained by permuting the labels:

V	$[\varepsilon\varepsilon\varepsilon]$	$[\varepsilon\varepsilon r]$	$[\varepsilon\varepsilon 1]$	$[\varepsilon r r]$	$[\varepsilon r 1]$	$[\varepsilon 11]$	$[rrr]$	$[rr1]$	$[r11]$	$[111]$
$\sum_{\phi \in \widehat{A_x}} \pi(x,\phi)^{\mathcal{I}}_{q=1}$	9	10	2	10	4	1	6	0	0	0

Table 25: Parahoric restriction of Iwahori-spherical $E_6(F)$-representations with respect to K_4

Since $\Pi(u,s,w) = \pi(x,\mathbf{1}) + \theta\pi(x,\theta) + \theta^2\pi(x,\theta^2)$ and $\Pi(u,s,w^2) = \pi(x,\mathbf{1}) + \theta^2\pi(x,\theta) + \theta\pi(x,\theta^2)$, we want to determine the parahoric restriction for each $\pi(x,\phi)$ individually.

We get a first hint by observing the following: while the affine Weyl group of $E_6(F)_{\text{ad}}$ is the extended affine Weyl group and thus contains the fundamental alcove stabilizers, the affine Weyl group of $E_6(F)_{\text{sc}}$ is the affine Weyl group, which acts simply transitively on the alcoves. In particular, the triality automorphism permuting the legs of the affine Dynkin diagram of E_6 is inner for $E_6(F)_{\text{ad}}$ but outer for $E_6(F)_{\text{sc}}$. This outer automorphism acts non-trivially on the unipotent representations of G, and this action is described by the work of Solleveld. In this particular case, if we pick an appropriate

fundamental alcove stabilizer $\omega \in \Omega$, we get that

$$\pi(x,\mathbf{1})^\omega = \pi(x,\theta), \quad \pi(x,\theta)^\omega = \pi(x,\theta^2), \quad \pi(x,\theta^2)^\omega = \pi(x,\mathbf{1}).$$

Since the twist by ω permutes the legs of the affine Dynkin diagram, we immediately get that the table looks roughly like

V	$[\varepsilon\varepsilon\varepsilon]$	$[\varepsilon\varepsilon r]$	$[\varepsilon\varepsilon 1]$	$[\varepsilon r r]$	$[\varepsilon r 1]$	$[\varepsilon 11]$	$[rrr]$	$[rr1]$	$[r11]$	$[111]$
$\pi(x,\mathbf{1})^{\mathcal{I}}_{q=1}$	3	a,b,c				0,1,0	2	0	0	0
$\pi(x,\mathbf{1})^{\mathcal{I}}_{q=1}$	3	b,c,a				0,0,1	2	0	0	0
$\pi(x,\mathbf{1})^{\mathcal{I}}_{q=1}$	3	c,a,b				1,0,0	2	0	0	0

Table 26: Parahoric restriction of Iwahori-spherical $E_6(F)$-representations with respect to K_4

where the three multiplicities correspond to the permutations of the labels, and the white columns look analogous.

Without further effort, this already highlights a problem since we obtain the multiplicities

V	$[\varepsilon 11]$	$[1\varepsilon 1]$	$[11\varepsilon]$
$\pi(x,\mathbf{1})^{\mathcal{I}}_{q=1}$	0	1	0
$\pi(x,\theta)^{\mathcal{I}}_{q=1}$	0	0	1
$\pi(x,\theta^2)^{\mathcal{I}}_{q=1}$	1	0	0
$\Pi(u,s,w)$	θ^2	1	θ
$\Pi(u,s,w^2)$	θ	1	θ^2

However, since the pair (w,s) is Γ_u-conjugate to (s,w^2) and Lusztig's nonabelian Fourier transform is trivial for the representations of finite groups of Lie type of type A_n, the conjecture predicts that $\Pi(u,s,w)$ and $\Pi(u,s,w^2)$ should have the same parahoric restrictions with respect to K_4. A similar problem arises with other columns.

Using Kato's results, it is also possible to program this explicitly using GAP. Kato shows that, as $W_{\mathrm{aff}} \times A_x$-modules,

$$H(\mathcal{B}^x) = \mathrm{Ind}_{W_{M,\mathrm{aff}} \rtimes A_x}^{W_{\mathrm{aff}} \times A_x} H(\mathcal{B}^u_s) \otimes s,$$

where (in this case) $a \in A_x$ (nontrivial) acts on $W_{M,\mathrm{aff}}$ by conjugation over a 3-cycle $j_0(a) \in N_W(W_M)$ under the canonical isomorphism $N_W(W_M) \cong W_M \rtimes S_3$ (here S_3 is the group of elements in W mapping the set $\{s_2,s_3,s_4,s_5\}$ to itself). Moreover, $W_{M,\mathrm{aff}} \rtimes A_x$ embeds in $W_{\mathrm{aff}} \times A_x$ via the map $w \rtimes a \mapsto (wj_0(a),a)$. I am not quite sure how does $A_x \subset W_{M,\mathrm{aff}} \rtimes A_x$ act on the one-dimensional $H(\mathcal{B}^u_s)$, but it does not really matter since the distinct options just permute the results cyclically. So let's assume that A_x acts trivially on $H(\mathcal{B}^u_s)$. In that case, using GAP to explicitly compute the induction, we get the complete table:

This is a general pattern that can be found for other sets of representations $\pi(us,\phi)$ when $Z_{G^\vee}(s)$ is not connected. Here is the relevant table:

Table 27: Parahoric restriction of $\pi(x,\phi)$ with respect to K_4

V	$[\varepsilon\varepsilon\varepsilon]$	$[\varepsilon\varepsilon r]$	$[\varepsilon\varepsilon 1]$	$[\varepsilon rr]$	$[\varepsilon r1]$	$[\varepsilon 11]$	$[rrr]$	$[rr1]$	$[r11]$	$[111]$
$\pi(x,\mathbf{1})^{\mathcal{I}}_{q=1}$	3	3,4,3	1,0,1	3,4,3	1,0,1,0,1,1	0,1,0	2	0	0	0
$\pi(x,\theta)^{\mathcal{I}}_{q=1}$	3	4,3,3	0,1,1	3,3,4	0,1,0,1,1,1	0,0,1	2	0	0	0
$\pi(x,\theta^2)^{\mathcal{I}}_{q=1}$	3	3,3,4	1,1,0	4,3,3	1,1,1,1,0,0	1,0,0	2	0	0	0
$\Pi(u,s,w)$	0	$\theta,1,\theta^2$	$-\theta,-1,-\theta^2$	$\theta^2,1,\theta$	$-\theta,-1,-\theta,-1,-\theta^2,-\theta^2$	$\theta^2,1,\theta$	0	0	0	0
$\Pi(u,s,w^2)$	0	$\theta^2,1,\theta$	$-\theta^2,-1,-\theta$	$\theta,1,\theta^2$	$-\theta^2,-1,-\theta^2,-1,-\theta,-\theta$	$\theta,1,\theta^2$	0	0	0	0

Table 28: Parahoric restriction of Iwahori-spherical $E_6(F)$-representations with respect to K_4

V	$[\varepsilon\varepsilon\varepsilon]$	$[\varepsilon\varepsilon r]$	$[\varepsilon\varepsilon 1]$	$[\varepsilon rr]$	$[\varepsilon r1]$	$[\varepsilon 11]$	$[rrr]$	$[rr1]$	$[r11]$	$[111]$
$[E_6,1]$	1	0	0	0	0	0	0	0	0	0
$[E_6(a_1),1]$	1	1	0	0	0	0	0	0	0	0
$[E_6(a_3),+]$	4	3	1	2	0	0	1	0	0	0
$[E_6(a_3),-]$	0	1	1	1	0	0	0	0	0	0
$[A_1A_5,+]$	4	2	0	1	0	0	1	0	0	0
$[D_4(71),1]$	3	3,4,3	1,0,1	3,4,3	1,0,1,0,1,1	0,1,0	2	0	0	0
$[D_4(71),\theta]$	3	4,3,3	0,1,1	3,3,4	0,1,0,1,1,1	0,0,1	2	0	0	0
$[D_4(71),\theta^2]$	3	3,3,4	1,1,0	4,3,3	1,1,1,1,0,0	1,0,0	2	0	0	0
$[D_4(a_1),1]$	15	15	3	17	4	1	19	4	1	0
$[D_4(a_1),\epsilon]$	0	0	1	1	1	0	1	0	0	0
$[D_4(a_1),r]$	3	8	4	11	4	1	11	3	0	0
$[D_4(53),1]$	7	10,11,10	4	13,14,13	5,4,4,4,4,5	1	14	3,4,3	0,1,0	0
$[D_4(53),\theta]$	7	10,10,11	4	14,13,13	4,4,5,5,4,4	1	14	3,3,4	1,0,0	0
$[D_4(53),\theta^2]$	7	11,10,10	4	13,13,14	4,5,4,4,5,4	1	14	4,3,3	0,0,1	0
$[3A_2,1\boxtimes 1]$	4	2,3,2	0	2,3,2	1,0,0,0,0,1	0	3	0,1,0	0,1,0	0
$[3A_2,1\boxtimes\theta]$	4	2,2,3	0	3,2,2	0,0,1,1,0,0	0	3	0,0,1	1,0,0	0
$[3A_2,1\boxtimes\theta^2]$	4	3,2,2	0	2,2,3	0,1,0,0,1,0	0	3	1,0,0	0,0,1	0
$[D_4(3311),1]$	9	29	21	80	53	33	216	140	89	57
$[D_4(3311),\epsilon]$	3	13	11	40	29	19	96	60	33	15

To be clear, this is the ONLY case where such a discrepancy with the conjecture arises. It seems to be a combination of the extra symmetry of the affine Dynkin diagram with the node 4 deleted and the fact that $Z_{G^\vee}(s)$ is disconnected in this case, thus allowing L-packets to be larger than the adjoint case.

The obvious fix, maybe quite unnatural, is to define both Fourier transforms to be $\mathbb{C}$-antilinear (or conjugate linear map) instead of $\mathbb{C}$-linear. With that definition, this issue is completely resolved, and it does not seem to break any of the other exceptional groups. I will next try to check whether it breaks $\mathrm{SL}_n(F)$ or not. Would that make sense theoretically?