

Local Langlands for GL_2

Yiannis Fam, Albert Lopez Bruch, Jakab Schrettner

June 6, 2024

1 Locally Profinite Groups and Smooth Representations

The aim of this first section is to motivate the abstract notions of a *locally profinite group* and of a *smooth representation*, which will be the main objects of study during the later subsections. Most of the motivation for studying such groups comes from the study of representations of reductive groups over non-Archimedean local fields. Therefore, we begin this section by briefly recalling some basic facts about these fields and linear groups associated to them. For the sake of brevity, we will omit proofs, and therefore we assume familiarity with the subject. For the unfamiliar reader, we encourage them to read **(insert here Gouvea reference, or others)**, where a detailed explanation is provided.

1.1 Local Fields and Locally Profinite Groups

We begin by recalling some basic objects from algebraic number theory. Given a field F , a **discrete valuation** on F is a surjective function $\nu : F \rightarrow \mathbb{Z} \cup \{\infty\}$ satisfying the three conditions

1. $\nu(xy) = \nu(x) + \nu(y)$ for any $x, y \in F$
2. $\nu(x + y) \geq \min\{\nu(x), \nu(y)\}$ for any $x, y \in F$.
3. $\nu(x) = \infty$ if and only if $x = 0$.

Any discrete valuation ν induces an absolute value given by the formula

$$|x| = c^{\nu(x)}$$

for any $c \in (0, 1)$, and therefore it also induces a topology which is independent of the choice of c . One easily checks that this absolute value satisfies $|x + y| \leq \max\{|x|, |y|\}$ for any $x, y \in K$. Absolute values with this property are denoted as *non-Archimedean*.

A field F with an absolute value $|\cdot|$ induced by a discrete valuation ν is the fraction field of the **valuation ring**

$$R := \{x \in F : \nu(x) \geq 0\} = \{x \in K : |x| \leq 1\},$$

which contains a unique maximal ideal

$$\mathfrak{p} := \{x \in F : v(x) > 0\} = \{x \in K : |x| < 1\},$$

denoted as the **valuation ideal** or the **ring of integers of F** . The valuation ideal is principal, and it is generated by any $\varpi \in K$ with $v(\varpi) = 1$. Such an element is called a uniformizer of F . Finally, the residue field κ of F is the quotient R/\mathfrak{p} . This motivates the following important definition.

Definition 1.1. A field F is a (non-Archimedean) *local field* if it is complete with respect to a topology induced by a discrete valuation and with finite residue field.

Remark 1.2. When the residue field is finite, it is conventional to write $|x| = q^{-v(x)}$, where $q = |\kappa|$. From here onwards, we will follow this convention.

Remark 1.3. Local fields are ubiquitous in number theory. They arise as completions of number fields at non-archimedean places, if they have 0 characteristic, or as completions of finite extensions of $\mathbb{F}_p(t)$ at non-archimedean places, if the characteristic is positive.

Let us now discuss important aspects of the topology on F and R induced by the discrete valuation v . We have already seen that R is a local ring with maximal ideal \mathfrak{p} and therefore $U_F := R \setminus \mathfrak{p}$ is the set of units of R . The ideals

$$\mathfrak{p}^n = \{x \in F : v(x) \geq n\} = \{x \in F : |x| \leq q^{-n}\} = \varpi^n R, \quad n \in \mathbb{Z}$$

are a complete set of fractional ideals of R and, since the valuation is assumed to be discrete, they are also open subsets of F . Therefore, they are a fundamental system of neighbourhoods of the identity. A direct consequence of this fact implies that F (and therefore R) are totally disconnected topological rings.

Furthermore, the ring R is a closed subring of F , which is assumed to be complete. Hence, R is also complete, and a standard topological argument shows that R is in fact compact. This proves that R (and therefore any \mathfrak{p}^n) is in fact a profinite group, and we have a topological isomorphism

$$R \longrightarrow \varprojlim_{n \geq 1} R/\mathfrak{p}^n \quad x \mapsto (x \pmod{\mathfrak{p}^n})_{n \geq 1}$$

where the maps implicit in the right hand side are the obvious ones.

However, F itself is clearly not compact, and therefore it is not profinite either. Nevertheless, F has the important property that any open neighbourhood of the identity contains an open compact (and therefore profinite) subgroup - some \mathfrak{p}^n for a sufficiently large n .

We are now ready to give the main definition of this section, which encapsulates this last property in greater generality.

Definition 1.4. A topological group G (which we always assume to be Hausdorff) is a *locally profinite group* if every open neighbourhood of the identity contains a compact open subgroup.

In this document we will be interested in studying the representation theory of many important groups and rings related to the local field F . The notion of a locally profinite group is an abstract one, but it has the great advantage of accomodating many important groups and rings associated to non-Archimedean local fields and their representation theory.

Examples 1.5. (1) Trivially, any group equipped with the discrete topology is profinite, where $\{e\}$ is the fundamental neighbourhood.

- (2) In the preceding discussion, we have shown that F is a locally profinite group, where \mathfrak{p}^n for $n \geq 1$ is a fundamental system of open compact subgroups. We remark that F satisfies the rather special property of being the union of its open compact subgroups.
- (3) The multiplicative group F^\times is also a locally profinite group, where the congruence unit groups $U_F^n = 1 + \mathfrak{p}^n$ for $n \geq 1$ is a fundamental system of open compact subgroups. We remark that unlike F , the group F^\times is not the union of its open compact subgroups.
- (4) Given $m \geq 1$ an integer, the additive group $F^m = F \times \cdots \times F$ is also a locally profinite group endowed with the product topology. A fundamental system of open compact subgroups is given by $\mathfrak{p}^n \times \cdots \times \mathfrak{p}^n$ for $n \geq 1$. More generally, any product of locally profinite groups is locally profinite.
- (5) The matrix ring $M_m(F)$ is also locally profinite since it is isomorphic to F^{m^2} as additive groups. The open compact subgroups $\mathfrak{p}^n M_m(R)$ are a fundamental system of neighbourhood of the identity.
- (6) The group $\mathrm{GL}_m(F)$ of invertible matrices is an open subset of $M_m(F)$ since $\det : M_m(F) \rightarrow F$ is continuous and F^\times is an open subset of F . Furthermore, mutiplication by a matrix $A \in M_m(F)$ and inversion of matrices are continous maps in $M_m(F)$ and therefore $\mathrm{GL}_m(F)$ is also a topological group and the subgroups

$$K = \mathrm{GL}_m(R), \quad K_n = 1 + \mathfrak{p}^n M_m(R), \quad n \geq 1,$$

are compact open, and a fundamental neighbourhood of the identity.

- (7) Let G be a locally profinite group and $H \leq G$ be a closed subgroup. Then H is also a locally profinite group and if in addition H is assumed to be normal in G , then G/H is also locally profinite.

We give some further insight into the terminology used. It is an easy exercise to prove that a profinite group is compact and locally profinite. Rather strickingly, the converse also holds. That is, if K is a compact locally profinite group, then

$$K \longrightarrow \varprojlim_N K/N$$

is a topological isomorphism where N ranges over the normal open subgroups, and the implicit maps are the obvious ones. Since K is compact and N is open, K/N must be finite and discrete, showing that K is profinite.

1.2 Abstract Representations of Groups

Before discussing the representation theory of locally profinite groups, we first review some general results and constructions of representations of arbitrary groups G . We begin by recalling the notion of a representation.

Definition 1.6. A representation of a group G over a field k is a pair (π, V) where V is a k -vector space and $\pi : G \rightarrow \mathrm{GL}(V)$ is a group homomorphism. We say that $\dim V$ is the dimension of the representation.

Equivalently, a representation of G is a k -vector space V equipped with a k -linear G action. Whenever the representation is clear from the context, we will omit π from the notation and write $g \cdot v$ for $\pi(g)v$.

Throughout this document, we will only be interested in complex representations, so for now on we will assume that $k = \mathbb{C}$ unless otherwise stated, and hence we will omit the underlying field from the notation.

We say that $U \leq V$ is a G -subspace if U is closed under the G -action; i.e. if $g \cdot U \subseteq U$ for every $g \in G$. When this happens, both U and V/U are naturally G -representations. Importantly, we say that a representation (π, V) is irreducible (or simple) if V has no non-trivial G -subspaces. These are the *building blocks* of more complicated representations, and thus we are often interested in classifying them.

This also motivates the following definition.

Definition 1.7. A representation (π, V) of a group G is semisimple if it is the direct sum of simple subrepresentations.

Remark 1.8. If G is a finite group, Maschke's Theorem shows that all finite dimensional complex representations of G are semisimple. As a consequence, one can show that any complex irreducible representation of G is finite dimensional, appearing as a subrepresentation of the canonical representation $\mathbb{C}G$.

As we shall visit later in this chapter, continuous finite dimensional representations over profinite groups also share these properties. However, it is easy to construct representations of locally profinite groups which are continuous yet not semisimple. For example,

$$\begin{aligned} \phi : \mathbb{Z} &\longrightarrow \mathrm{GL}_2(\mathbb{C}) \\ n &\mapsto \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix} \end{aligned}$$

has a single one-dimensional invariant subspace. One can also construct irreducible representations that are infinite dimensional, but these are harder to construct. **Reference to a later section when we consider the reps of the Mirabolic subgroup or the Steinberg reps of $\mathrm{GL}_2(F)$.**

Naturally, we also define the notion of a homomorphism between representations.

Definition 1.9. A morphism between two complex representations (π, V) , (σ, W) of a group G is a linear map $\phi : V \rightarrow W$ compatible with the G action. That is,

$$\phi(\pi(g)v) = \sigma(g)\phi(v) \text{ for all } g \in G, v \in V.$$

Therefore, the set of complex representations of G form a category denoted by Rep_G , which importantly is an **abelian category**.

We finish this section by introducing important constructions and functors between these categories that allow us to obtain new representations from old ones, and which we will use heavily later on.

Definition 1.10. Given $(\pi, V) \in \text{Rep}_G$, define the dual space $V^* = \text{Hom}(V, \mathbb{C})$, and denote by

$$\begin{aligned} V^* \times V &\longrightarrow \mathbb{C}, \\ (v^*, v) &\longmapsto \langle v^*, v \rangle, \end{aligned}$$

the canonical evaluation homomorphism. Then V^* carries a natural representation

$$\langle \pi^*(g)v^*, v \rangle = \langle v^*, \pi(g^{-1})v \rangle,$$

denoted as the dual representation, and the functor

$$\begin{aligned} (-)^* : \text{Rep}_G &\longrightarrow \text{Rep}_G \\ (\pi, V) &\longrightarrow (\pi^*, V^*) \end{aligned}$$

is an additive and exact contravariant functor.

One can also consider the composition of this functor with itself to obtain the **double dual** (π^{**}, V^{**}) . Then there is a canonical G -homomorphism $\delta : V \rightarrow V^{**}$ such that

$$\langle \delta(v), v^* \rangle_{V^{**}} = \langle v^*, v \rangle_V.$$

When V is finite dimensional, δ is a G -isomorphism. For general representations of locally profinite groups, this is not always the case, but under additional assumptions it is possible to give a precise criterion that determines when δ is bijective ([BH06, 2.8 Corollary, 2.9 Proposition]).

Definition 1.11. Let $H \leq G$ be groups and let (π, V) and (σ, W) be representations of G and H respectively. The restriction of π to H gives a **restriction** functor

$$\begin{aligned} \text{Res}_H^G : \text{Rep}_G &\longrightarrow \text{Rep}_H \\ (\pi, V) &\longmapsto (\pi|_H, V) \end{aligned}$$

On the other hand, given $(\sigma, W) \in \text{Rep}_H$ one can construct the vector space

$$X = \{f : G \rightarrow W : f(hg) = \sigma(h)f(g) \text{ for all } h \in H, g \in G\},$$

equipped with the G -action $\Sigma : G \longrightarrow \text{Aut}_{\mathbb{C}}(X)$ such that

$$\Sigma(g)f : x \longmapsto f(xg), \quad x, g \in G.$$

This gives the **induction** functor

$$\begin{aligned} \text{Ind}_H^G : \text{Rep}_H &\longrightarrow \text{Rep}_G \\ (\pi, V) &\longmapsto (\Sigma, X). \end{aligned}$$

As with the dual functor, both the restriction and induction functors are additive and exact covariant functors. To simplify notation, we will write $\text{Ind}_H^G \sigma$ instead of $\text{Ind}_H^G(\sigma, W)$, which is the usual convention in the literature.

We remark that one can construct the following canonical H -homomorphisms

$$\begin{aligned} a_\sigma : \text{Ind}_H^G \sigma &\longrightarrow W \\ f &\longmapsto f(1) \end{aligned}$$

and

$$\begin{aligned} a_\sigma^c : W &\longrightarrow \text{Ind}_H^G \sigma \\ w &\longmapsto f_w \end{aligned}$$

where f_w is supported in H and $f_w(h) = \sigma(h)w$ for $h \in H$. The choice of notation will be understood later. These, in turn, induce the maps

$$\begin{aligned} \Psi : \text{Hom}_G(\pi, \text{Ind}_H^G \sigma) &\longrightarrow \text{Hom}_H(\text{Res}_H^G \pi, \sigma), \\ \phi &\longmapsto a_\sigma \circ \phi, \end{aligned}$$

and

$$\begin{aligned} \Psi^c : \text{Hom}_G(\text{Ind}_H^G \sigma, \pi) &\longrightarrow \text{Hom}_H(\sigma, \text{Res}_H^G \pi), \\ f &\longmapsto f \circ a_\sigma^c. \end{aligned}$$

When G is a finite group, we have the following result.

Theorem 1.12 (Frobenius reciprocity). *Let G be a finite group. Then the maps Ψ and Ψ^c are bijections that are functorial in both variables σ and π . In categorical terms, we have the adjunctions*

$$\text{Ind}_H^G \dashv \text{Res}_H^G \dashv \text{Ind}_H^G.$$

This theorem fails for general representations of locally profinite groups G and subgroups $H \leq G$. However, the theorem does hold under certain additional assumptions on the topology of H inside G , a smoothness condition on the representations and an adequate modification of the set X above. This will all be discussed in Section **Insert here reference to the section in which this is discussed**.

1.3 Characters of Local Fields

Now we turn our attention to the representation theory of locally profinite groups. In studying their representations, it turns out that the entire set is too big, so we need to restrict our attention to those representations satisfying a certain “smoothness” condition. To motivate this condition, we will first describe the simplest (yet very relevant!) case: one-dimensional representations of a local field F : that is, group homomorphisms $\phi : F \rightarrow \mathbb{C}^\times$. Later in this section we will also study the one-dimensional representations of F^\times .

As we have discussed in the previous section, locally profinite groups carry a certain topology, so a natural condition to impose is **continuity** with respect to the usual topologies in \mathbb{C}^\times and G . A continuous homomorphism $\psi : G \rightarrow \mathbb{C}^\times$ will be denoted as a *character* of G .

Characters of a locally profinite group G are a group under multiplication, denoted by \hat{G} . It turns out that for one-dimensional representations, continuity coincides with the smoothness condition we require which will be introduced later.

When G is a finite group with discrete topology, then any one-dimensional representation is a character, and we have the following simple description.

Proposition 1.13. *If G is a finite group with the discrete topology, then $\hat{G} \cong G^{ab}$. In particular, if G is abelian then $\hat{G} \cong G$.*

Proof. **Insert reference here** □

For general locally profinite results, we have this rather surprising result.

Lemma 1.14. *Let G be a locally profinite group and $\psi : G \rightarrow \mathbb{C}^\times$ a homomorphism. Then ψ is continuous if and only if $\ker \psi$ is open in G . Furthermore, if G is the union of its compact open subgroups, then¹*

$$\psi(G) \subseteq \{z \in \mathbb{C}^\times : |z| = 1\} = S^1.$$

Proof. If $\ker \psi = \psi^{-1}(1)$ is open in G , then for any $z \in \text{Im } \psi$, then $\psi^{-1}(z) = g \ker \psi$ is also open, where $\psi(g) = z$. So in fact, **for any** $U \subseteq \mathbb{C}^\times$,

$$\psi^{-1}(U) = \bigcup_{z \in U \cap \text{Im } \psi} \psi^{-1}(z),$$

and so in particular it is continuous. Conversely, if ψ is continuous, then for any open neighbourhood \mathcal{N} of 1, $\psi^{-1}(\mathcal{N})$ contains an open compact subgroup K of G . But \mathcal{N} can be chosen sufficiently small so that it does not contain any non-trivial subgroup of \mathbb{C}^\times . Hence, $\psi(K) = 1$ so $K \subseteq \ker \psi$, and since K is open, so is $\ker \psi$. The last assertion is a direct consequence of the fact that the continuous image of a compact set is compact and S^1 is the unique maximal compact subgroup of \mathbb{C}^\times . □

Since F is the union of its open compact subgroups, all characters of F are unitary. However, this is not the case for F^\times . Indeed, the map $x \mapsto |x|$ is a character of F^\times , yet it is clearly not unitary.

Before stating the classification theorem for characters of F , we need one last definition.

Definition 1.15. Let ψ be a non-trivial character of F (resp. of F^\times). The **level** of ψ is defined as be the least $d \geq 0$ such that $\mathfrak{p}^d \subseteq \ker \psi$ (resp. $U_F^{d+1} \subseteq \ker \psi$).

Lemma 1.16. *Let $\psi \in \hat{F}$ be a character of level d and let $a \in F$. Then the map $a\psi : x \mapsto \psi(ax)$ is a character of F and if $a \neq 0$ then $a\psi$ has level $d - \nu_F(a)$.*

¹Characters satisfying this property are called *unitary*

Proof. It is clear that $a\psi$ is a character since if $x \in \mathfrak{p}^{d-\nu_F(a)}$, then $ax \in \mathfrak{p}^d$ so $a\psi(x) = 1$ so $\mathfrak{p}^{d-\nu_F(a)} \subseteq \ker(a\psi)$ and the kernel is open. Furthermore, there is some $y \in \mathfrak{p}^{d-1}$ such that $\psi(y) \neq 1$ so $a\psi(a^{-1}y) \neq 1$. Since $a^{-1}y \in \mathfrak{p}^{d-1-\nu_F(a)}$, this indeed shows that the level of $a\psi$ is $d - \nu_F(a)$. \square

We are now ready to give the classification theorem for \hat{F} .

Theorem 1.17 (Additive duality). *Let $\psi \in \hat{F}$ be character with level d . The map $a \mapsto a\psi$ induces an isomorphism $F \cong \hat{F}$.*

The proof of surjectivity of the theorem requires an inductive step, which heavily relies on the following results.

Lemma 1.18. *Let $\psi \in \hat{F}$ be a character of level d and let $u, u' \in U_F$ be two units of F . Then $u\psi$ coincides with $u'\psi$ on \mathfrak{p}^{d-n} if and only if $u'u^{-1} \in U_F^n$.*

Proof. Let $\alpha = \nu_F(u - u')$. A simple definition chase shows that $u\psi$ and $u'\psi$ agree on \mathfrak{p}^{d-n} if and only if $\mathfrak{p}^{d-n+\alpha} = (u - u')\mathfrak{p}^{d-n} \subseteq \ker \psi$. By definition of level, this is the case if and only if $\alpha \geq n$; that is, if $u \equiv u' \pmod{\mathfrak{p}^n}$ or $u'u^{-1} \in U_F^n$. \square

Lemma 1.19. *Let $\theta : \mathfrak{p}^n \rightarrow \mathbb{C}^\times$ be a character. Then there are exactly q characters Θ of \mathfrak{p}^{n-1} such that $\Theta|_{\mathfrak{p}^n} = \theta$.*

Proof. Since $\hat{\kappa} \cong \kappa$, it is enough to construct a bijection between $\mathcal{A} := \{\Theta \in \widehat{\mathfrak{p}^{n-1}} : \Theta|_{\mathfrak{p}^n} = \theta\}$ and $\hat{\kappa}$. Let $\phi = \theta^{-1}$ and let Φ be **any** lift of ϕ as a character of \mathfrak{p}^{n-1} . Now given $\Theta \in \mathcal{A}$, the character $\Theta \cdot \Phi$ is trivial on \mathfrak{p}^n and thus it descends to a map

$$\overline{\Theta \cdot \Phi} : \kappa \cong \mathfrak{p}^{n-1}/\mathfrak{p}^n \longrightarrow \mathbb{C}^\times.$$

To construct an inverse to the map $\Theta \mapsto \overline{\Theta \cdot \Phi}$, choose some $\chi \in \hat{\kappa}$, view it as a character of $\mathfrak{p}^{n-1}/\mathfrak{p}^n$ and consider the map $\tilde{\chi} : \mathfrak{p}^{n-1} \rightarrow \mathbb{C}^\times$ given by $\tilde{\chi}(u) = \chi(u + \mathfrak{p}^n)$. Then the map $\chi \mapsto \Phi^{-1} \cdot \tilde{\chi}$ is the required inverse map. \square

We are now ready for the proof of Additive duality.

Proof of Theorem 1.17. The map $a \mapsto a\psi$ is clearly a homomorphism. To prove injectivity, suppose that $a \neq b$ but $a\psi = b\psi$. Then it follows that $x(a - b) \in \ker \psi$ for all $x \in F$. But since $a - b \neq 0$, then $\ker \psi = F$, a contradiction.

Let $\theta \in \hat{F}$ be any non-trivial character (if θ were trivial, then $0\psi = \theta$), and let l be the level of θ . By replacing θ with $\varpi^{l-d}\theta$, which has level d , we may assume without loss of generality that θ and ψ have the same level d , and therefore they both agree on \mathfrak{p}^d . To show there is some $u \in F$ (in fact, $u \in U_F$ necessarily) such that $u\psi = \theta$, we construct a sequence $\{u_n\}_{n \geq 0}$ inductively such that $u_n\psi|_{\mathfrak{p}^{d-n}} = \theta|_{\mathfrak{p}^{d-n}}$ and $u_{n+1} \equiv u_n \pmod{\mathfrak{p}^n}$. Such a sequence is clearly Cauchy, and since F is complete, it converges to some $u \in U_F$ such that $u \equiv u_n \pmod{\mathfrak{p}^n}$ for all $n \geq 1$ and thus $u\psi$ agrees with θ on $\cup_{n \in \mathbb{Z}} \mathfrak{p}^n = F$, which concludes the proof.

Thus, it remains to construct the sequence above. To construct u_1 we note that by Lemma 1.19, there are exactly $q-1$ non-trivial characters on \mathfrak{p}^{d-1} that are trivial on \mathfrak{p}^d . In addition, by Lemma 1.18, as u ranges over the cosets of U_F/U_F^1 , the characters $u\psi|_{\mathfrak{p}^{d-1}}$ are distinct. Since $|U_F/U_F^1| = |\kappa^\times| = q-1$, there is some $u_1 \in U_F$ such that $u_1\psi$ agrees with θ on \mathfrak{p}^{d-1} .

Assuming now we have constructed u_1, \dots, u_n in U_F with the desired conditions, we note that by Lemma 1.19, there are exactly q characters of \mathfrak{p}^{d-n-1} that coincide with $\theta|_{\mathfrak{p}^{d-n}}$ when they are restricted. Again by Lemma 1.18, as α ranges over the cosets of U_F^n/U_F^{n+1} the characters $\alpha u_n\psi$ are distinct on \mathfrak{p}^{d-n-1} but they all coincide on \mathfrak{p}^{d-n} . Since $|U_F^n/U_F^{n+1}| = |\kappa| = q$, there is some α_n such that $\alpha_n u_n\psi$ coincides with θ on \mathfrak{p}^{d-n-1} . Since $\alpha_n \in U_F^n$, $\alpha_n u_n \equiv u_n \pmod{\mathfrak{p}^n}$. Hence $u_{n+1} := \alpha_n u_n$ has the required properties. \square

1.4 Smooth Representations of Locally Profinite Groups

We now turn our attention representations of locally profinite groups of arbitrary dimension. For one-dimensional representations, we imposed a natural continuity condition, and Lemma 1.14 showed that characters have open kernel. This is a remarkable result, since this means that the homomorphism is continuous with respect to **any** topology on \mathbb{C}^\times , not just the usual one.

If V is a finite dimensional representation of a locally profinite group G , the group $\text{Aut}_{\mathbb{C}}(V)$ has a natural topology as an open subspace of $M_n(\mathbb{C}) \cong \mathbb{C}^{n^2}$. It turns out that, analogously to \mathbb{C}^\times , small neighbourhoods of the identity of $\text{Aut}_{\mathbb{C}}(V)$ do not contain any non-trivial subgroups. The same reasoning as in Lemma 1.14 shows that continuous finite dimensional representations have open kernel too. That is, the homomorphism is continuous with respect to any topology on $\text{Aut}_{\mathbb{C}}(V)$.

However, for infinite dimensional representations V , equipping $\text{Aut}_{\mathbb{C}}(V)$ with a topology is not as straightforward and the requirement of having an open kernel is too restrictive. The natural idea behind a *smooth representation* is the requirement that any finite dimensional subrepresentation V (if any) should have an open kernel. To give the precise definition, we first introduce the module of invariants and coinvariants.

Definition 1.20. Let $H \leq G$ be groups and (π, V) a representation of G . We define the H -invariants to be

$$V^H := \{v \in V : \pi(h)v = v \text{ for all } h \in H\},$$

and the H -coinvariants to be

$$V_H := V/V(H) \text{ where } V(H) = \text{Span}_{\mathbb{C}}\{v - \pi(h)v : v \in V, h \in H\}.$$

That is, V^H (resp. V_H) is the largest subspace (resp. quotient) on which H acts trivially.

We are now ready to define a *smooth representation* of a locally profinite group G .

Definition 1.21. A representation V of G is **smooth** if for $v \in V$ there exists a compact-open subgroup $K \subseteq G$ such that $v \in V^K$. In other words,

$$V = \bigcup_K V^K.$$

Moreover, we say V is **admissible** if V^K is finite-dimensional for all compact-open K .

Smooth representations of G are a full, abelian subcategory of Rep_G , and this category is denoted by Smo_G .

Remark 1.22. If (π, V) is a finite dimensional smooth representation and $\{v_1, \dots, v_n\}$ is a \mathbb{C} -basis such that $v_i \in V^{K_i}$ for some open compact subgroups K_i , then

$$K := \bigcap_{i=1}^n K_i \subseteq \ker \pi$$

and it is open and compact too, so the kernel is also open. Conversely, if $\ker \pi$ is open, then there is some open compact subgroup K fixing all of V , so in this case smooth and continuous coincide.

As we hinted in Remark 1.8, smooth representations of locally profinite groups have remarkable algebraic structures, and they share many properties with representations of finite groups, particularly if the group is compact (and thus profinite), which we briefly recall. A direct application (yet technical) of Zorn's Lemma provides the following useful criterion to determine whether a representation is semisimple.

Proposition 1.23. *Let (π, V) be a smooth representation of a locally profinite group G . The following are equivalent:*

1. *V is the sum of its irreducible G -subspaces.*
2. *V is the direct sum of a family of irreducible G -subspaces (i.e. V is semisimple)*
3. *any G -subspace of V has a G -complement in V .*

Proof. [BH06, 2.2 Lemma] □

Using this Proposition, we can now prove that smooth representations of profinite groups behave in a similar way to those of finite groups. We note that any open compact subgroup K of a locally profinite group G is profinite, and that any smooth G -representation is naturally a smooth K -representation by restriction. Therefore, the following results apply for any open compact subgroup of G .

Proposition 1.24. *Let (π, V) be a representation of a profinite group K . If V is irreducible then it is finite dimensional. Conversely, if V is finite dimensional, then it is semisimple.*

Proof. The first statement is a matter of following the definitions. Fix any non-zero $v \in V$, and suppose $v \in V^{K_0}$. Then the subspace

$$U = \text{Span}\{\pi(k)v : k \in K\} = \text{Span}\{\pi(k)v : k \in K/K_0\}$$

is clearly a K -subspace and it is also finite dimensional since K_0 is open and K is compact, so $(K : K_0)$ is finite.

To prove the second statement, let v and K_0 be as above. By replacing K_0 by $\bigcap_{g \in K/K_0} gK_0g^{-1}$ if needed, we may assume that K_0 is normal. As above, the subspace

$$W = \text{Span}\{\pi(k)v : k \in K\}$$

is finite dimensional and K_0 acts trivially on it. Thus W is effectively a finite dimensional representation of the finite group K/K_0 and thus by Maschke's Theorem, W is the sum of its irreducible K subspaces. Since v was arbitrary this shows that condition 1. of Proposition 1.23 is satisfied, so V is semisimple.

□

This proposition has important structure results. Let \hat{K} denote the set of equivalence classes of irreducible smooth representations of K . As we shall see, this notation is consistent with \hat{F} since all irreducible smooth representations of F are one-dimensional.

Let (π, V) be a smooth representation of a locally profinite group G and let K be an open compact subgroup. For each $\rho \in \hat{K}$, let V^ρ be the sum of all irreducible K -subspaces of V isomorphic to K , which is denoted as the ρ -isotypic component of V . Note also that $V^{1_K} = V^K$.

Proposition 1.25. *Let G be a locally profinite group and K a compact open subgroup of G . Let $(\tau, U), (\pi, V), (\sigma, W) \in \text{Smo}_G$ and $a : U \rightarrow V$ and $b : V \rightarrow W$ be G -homomorphisms.*

1. *The space V is the sum of the K -isotypic components:*

$$V = \bigoplus_{\rho \in \hat{K}} V^\rho.$$

2. *The following holds:*

$$W^\rho \cap b(V) = b(V^\rho).$$

3. *The sequence*

$$U \xrightarrow{a} V \xrightarrow{b} W$$

is exact if and only if

$$U^K \xrightarrow{a} V^K \xrightarrow{b} W^K$$

is exact for every compact open subgroup K of G .

4. *If $V(K)$ is the span of the elements $v - \pi(k)v$ for $v \in V, k \in K$, then*

$$V(K) = \bigoplus_{\substack{\rho \in \hat{K} \\ \rho \neq 1}} V^\rho \text{ and thus } V = V^K \oplus V(K)$$

and $V(K)$ is the unique K -compement of V^K in V .

Proof. [BH06, 2.3 Proposition and Corollary 1,2]

□

As promised in §1.2, we now discuss the dual, restriction and induction functors in the context of smooth representations of locally profinite groups. From our previous discussion, two major problems arise in this context. Firstly, given a locally profinite group G and a subgroup H , there is no guarantee that H is locally profinite and thus Smo_H may not be well-defined. Secondly, when we perform some construction on a smooth representation (constructing its dual, inducing to a bigger group,...) there is no guarantee that the resulting representation is smooth. Thankfully, both of this problems can be resolved in a straightforward way.

To ensure that H is locally profinite, we must add a condition on the topology on H . Based on Example 1.5(7), we just need to assume that H is a closed subgroup of G . In some cases, we will need to assume that H

is also open, which is a more restrictive condition. To ensure that the functors give smooth representations of G , we simply compose them with the **smoothness functor**

$$\begin{aligned} (-)^\infty : \text{Rep}_G &\longrightarrow \text{Smo}_G, \\ (\pi, V) &\longmapsto (\pi^\infty, V^\infty) \end{aligned}$$

where

$$V^\infty = \bigcup_K V^K \text{ and } \pi^\infty(g) = \pi(g)|_{V^\infty} \text{ for each } g \in G,$$

and K ranges over the compact open subgroups of G . By chasing definitions, one can show that $(-)^\infty$ is a well-defined left-exact functor such that

$$\text{Hom}_G(V, W) = \text{Hom}_G(V, W^\infty) \text{ for all } V \in \text{Smo}_G, W \in \text{Rep}_G.$$

Using this constructions, we can define the smooth dual, restriction and induction functors. We remark that as long as $H \leq G$ is closed, the usual restriction sends smooth representations of G to smooth representations of H . This is not the case for the dual and induction functors, so we apply to construction above.

Definition 1.26. If G is a locally profinite group, define the smooth dual functor

$$\begin{aligned} (\check{-}) : \text{Smo}_G &\longrightarrow \text{Smo}_G, \\ (\pi, V) &\longmapsto (\check{\pi}, \check{V}) \end{aligned}$$

where $(\check{\pi}, \check{V}) = (\pi^*, V^*)^\infty$.

The smooth dual satisfies an important property: if V is a smooth representation of G and $v \in V, v \neq 0$, then there is some $\check{v} \in \check{V}$ such that $\langle \check{v}, v \rangle \neq 0$. Consequently, the map $\delta : V \rightarrow \check{\check{V}}$ is injective, and the following proposition gives a criterion for surjectivity.

Proposition 1.27. *With the notation as above, the canonical map $\delta : V \rightarrow \check{\check{V}}$ is an isomorphism if and only if (π, V) is admissible.*

Proof. [BH06, 2.9 Proposition] □

We also define the smooth induction functor as the composition of the induction and smoothness functor.

Definition 1.28. Let G be a locally profinite group and $H \leq G$ a closed subgroup. Define the smooth induction functor

$$\begin{aligned} \text{Ind}_H^G : \text{Smo}_H &\longrightarrow \text{Smo}_G, \\ (\sigma, W) &\longmapsto (\Sigma, X) \end{aligned}$$

where X is the space of functions $f : G \rightarrow W$ satisfying

1. $f(hg) = \sigma(h)f(g)$ for all $h \in H, g \in G$.

2. There is a compact open subgroup $K \subseteq G$ (depending on f) such that $f(gk) = f(g)$ for all $g \in G, k \in K$.

Since the action Σ on X is given by $\Sigma(g)f : x \mapsto f(xg)$, condition 2. is precisely the smoothness condition that $f \in X^K$ for some open compact subgroup K . As above, we will denote this representation of G as $\text{Ind}_H^G \sigma$. Under these conditions, the first half of Frobenius Reciprocity holds:

Theorem 1.29 (Frobenius reciprocity). *Let (π, V) be a smooth representation of G , and (σ, W) a smooth representation of a closed subgroup H . Then the map*

$$\begin{aligned} \Psi : \text{Hom}_G(\pi, \text{Ind}_H^G \sigma) &\longrightarrow \text{Hom}_H(\text{Res}_H^G \pi, \sigma), \\ \varphi &\longmapsto \alpha_\sigma \circ \varphi, \end{aligned}$$

is a bijection that is functorial in both variables π, σ , and where $\alpha_\sigma : \text{Ind}_H^G \sigma \rightarrow W$ is the canonical map $\alpha_\sigma(f) = f(1)$. In categorical terms,

$$\text{Res}_H^G \dashv \text{Ind}_H^G.$$

However, in this context, it is not the case that Ind_H^G is left adjoint to Res_H^G . With a small modification we can however obtain an analogous result. Firstly, we note that to ensure that a_σ^c is a H -homomorphism, we need the stronger assumption that H is open in G . Secondly, we observe that given representations (π, V) and (σ, W) , of G and H respectively, $a_\sigma^c(w)$ is supported only in H for any $w \in W$. Hence, one should not consider the entire representation $\text{Ind}_H^G \sigma$, but rather a subrepresentation of it. Here is the precise construction.

Definition 1.30. Let G be a locally profinite group, H a closed subgroup, and (σ, W) a smooth representation of H . Consider the functor

$$\begin{aligned} c - \text{Ind}_H^G : \text{Smo}_H &\longrightarrow \text{Smo}_G, \\ (\sigma, W) &\longmapsto (\Sigma, X_c) \end{aligned}$$

where

$$X_c = \{f \in X : \text{supp } f \subseteq H \backslash G \text{ is compact}\}.$$

We denote functions satisfying the later condition as compactly supported modulo H , and this condition is equivalent to $\text{supp } f \subseteq HC$ for some compact set C . The action by Σ is closed in Σ_c , so the functor is well-defined.

This construction is mainly of interest in the case when H is open in G , in which case a_σ^c is a H -homomorphism. This construction satisfies the second half of Frobenius Reciprocity.

Theorem 1.31. *Let (π, V) be a smooth representation of G , and (σ, W) a smooth representation of an open subgroup H . Then the map*

$$\begin{aligned} \Psi^c : \text{Hom}_G(c - \text{Ind}_H^G \sigma, \pi) &\longrightarrow \text{Hom}_H(\sigma, \text{Res}_H^G \pi) \\ \varphi &\longmapsto \varphi \circ \alpha_\sigma^c \end{aligned}$$

*is a bijection that is functorial in both variables π, σ , and **where** $\alpha_\sigma^c : W \rightarrow c - \text{Ind}_H^G \sigma$ **is the map** $w \mapsto f_w$ **where** f_w **is supported in** H **and satisfies** $f_w(h) = hw$.*

In categorical terms, we have the important property

$$c - \text{Ind}_H^G \dashv \text{Res}_H^G \dashv \text{Ind}_H^G.$$

1.5 Schur's Lemma

We end this section by discussing a version of Schur's Lemma for smooth representations of locally profinite groups. We recall Schur's Lemma for finite groups.

Theorem 1.32. *Let G be a finite group and let (π, V) be a complex irreducible representation of G . Then for any $\phi \in \text{End}_G(V)$, there is some $\lambda \in \mathbb{C}$ such that $\phi(v) = \lambda v$ for all $v \in V$. In other words, $\text{End}_G(V) \cong \mathbb{C}$.*

As it turns out, Schur's Lemma does not hold for complex smooth irreducible representations of a locally profinite group G . However, it is true under a mild hypothesis that is satisfied by all locally profinite groups that will be of interest for us.

Hypothesis. For any compact open subgroup K of G , the set $K \backslash G$ is countable.

This is in fact a hypothesis that can be tested easily, as if it is true for one open subgroup K , then it is true for all. Under this hypothesis, the following result holds.

Lemma 1.33. *Let (π, V) be an irreducible smooth representation of G . Then the dimension $\dim_{\mathbb{C}} V$ is countable.*

Proof. Let $v \in V$, $v \neq 0$ and let K open compact subgroup such that $v \in V^K$. The set $\{\pi(g)v : g \in G\} = \{\pi(g)v : g \in K \backslash G\}$ spans V and it is countable. \square

We are now ready to state and prove Schur's Lemma in our context.

Theorem 1.34. *If (π, V) is a smooth irreducible representation of G , then $\text{End}_{\mathbb{C}} V \cong \mathbb{C}$.*

This results has two important corollaries worth recalling. For the first one, we note that given a locally profinite group G , its centre Z is a closed subgroup of G and therefore a locally profinite group too.

Corollary 1.35. *Let (π, V) be an irreducible smooth representation of G . The centre Z of G acts on V via a character $\omega_{\pi} : Z \rightarrow \mathbb{C}^{\times}$. In other words, $\pi(z)v = \omega_{\pi}(z)v$ for all $v \in V$ and $z \in Z$.*

Proof. [BH06, 2.6 Schur's Lemma] \square

Proof. For any $z \in Z$, the automorphism $\pi(z) : V \rightarrow V$ lies in $\text{End}_G(V) \cong \mathbb{C}$. Hence, the desired map $\omega_{\pi} : Z \rightarrow \mathbb{C}^{\times}$ does indeed exist, and it is a group homomorphism. To prove smoothness, we note that if K is an open compact subgroup such that $V^K \neq 0$, then ω_{π} is trivial on the open compact subgroup $K \cap Z$ of Z . So ω_{π} is indeed a character of Z . \square

The character ω_{π} is denoted as the **central character** of (π, V) .

Corollary 1.36. *If G is abelian, any irreducible smooth representation of G is one-dimensional.*

This justifies the notation \hat{K} for the set of equivalence classes of irreducible smooth representation of a locally profinite group K , since this notation can now be seen to coincide with the set of characters \hat{F} of F .

2 Measure and the Duality Theorem

So far, we have introduced the central objects that we will study throughout: locally profinite groups and smooth representations. In addition, we have given a complete classification of the equivalence classes of irreducible smooth representations of a local field F . These are all 1-dimensional by Schur's Lemma and therefore $\hat{F} \cong F$ by Additive Duality.

However, the analogous task for other locally profinite groups is considerably harder. Even the structure of the group of characters of F^\times is a lot more subtle. To describe the local Langlands correspondence for GL_2 we will need a classification theorem of some family of irreducible smooth representations of $\mathrm{GL}_2(F)$ denoted as principal series representations. To advance in this task, however, we need to introduce the notion of Haar measure and develop some measure theory on locally profinite groups. This is precisely the aim of this chapter, which follows a similar development to [BH06, Chapter 3].

We finish this section by studying the relationship between induction and duality, which is encapsulated by the Duality Theorem.

2.1 The Space $C_c^\infty(G)$ and the Haar Measure

Let G be a locally profinite group. Then we define $C_c^\infty(G)$ to be the space of functions $f : G \rightarrow \mathbb{C}$ which are locally constant and that have compact support. Such a function f can be equivalently characterized as a finite linear combination of characteristic functions of double cosets KgK for some open compact subgroup K of G . Hence, the space $C_c^\infty(G)$ is a complex vector space and admits two natural actions by G by left and right translation:

$$\lambda_g f : x \mapsto f(g^{-1}x), \quad \text{and} \quad \rho_g f : x \mapsto f(xg),$$

for $x, g \in G$ and $f \in C_c^\infty(G)$. These actions define smooth representation since characteristic functions of double cosets of K are invariant under translation by K . We are now ready to define the notion of a *Haar integral* and *Haar measure*.

Definition 2.1. A *left Haar integral* on G is a non-zero linear functional

$$I : C_c^\infty(G) \longrightarrow \mathbb{C}$$

such that

- (1) $I(\lambda_g f) = I(f)$, $g \in G$, $f \in C_c^\infty(G)$, and
- (2) $I(f) \geq 0$ for any $f \in C_c^\infty(G)$ such that $f \geq 0$.

A *right Haar integral* is defined analogously by replacing λ_g by ρ_g .

The usefulness of the Haar integral relies on the fact that locally profinite groups possess essentially one unique left Haar integral.

Proposition 2.2. *There exists a left Haar integral $I : C_c^\infty(G) \rightarrow \mathbb{C}$. Moreover, a linear functional $I' : C_c^\infty(G) \rightarrow \mathbb{C}$ is a left Haar integral if and only if $I' = cI$ for some constant $c > 0$.*

Proof. [BH06, 3.1 Proposition] □

Whenever we have a *left Haar integral* I , we can define the associated *left Haar measure* as follows. Let $S \subset G$ and let Γ_S be its characteristic function. Then $\Gamma_S \in C_c^\infty(G)$ if and only if S is open and compact. In that case, we define

$$\mu_G(S) = I(\Gamma_S)$$

to be the Haar measure of S . We note that $\mu_G(S) > 0$ and by left invariance, $\mu_G(gS) = \mu_G(S)$ for any $g \in G$. The relationship is commonly expressed by using the usual integral notation

$$I(f) = \int_G f(g) d\mu_G(g), \quad f \in C_c^\infty(G).$$

This choice of notation is motivated by the fact that since f is locally constant and has constant support, the integral $I(f)$ is effectively a finite weighted sum of $\mu_G(S_i)$ where S_i are open, compact and f is constant on them.

Example 2.3. The notion of a left Haar measure is only determined up to a constant. In practice, to uniquely determine the measure, we associate a particular open compact subset with a value. For example if $G = F$ is a local field, one commonly chooses μ_F so that $\mu_F(R) = 1$. Under this choice, we calculate that $\mu_F(\mathfrak{p}^n) = q^{-n}$.

Left Haar measures behave predictably under usual group constructions. For example, if G_1, G_2 are profinite groups, then $G = G_1 \times G_2$ is also profinite group, and we have an isomorphism

$$C_c^\infty(G_1) \otimes C_c^\infty(G_2) \longrightarrow C_c^\infty(G)$$

$$\sum_{i=1}^r f_i^1 \otimes f_i^2 \longmapsto \left((g_1, g_2) \mapsto \sum_{i=1}^r f_i^1(g_1) f_i^2(g_2) \right).$$

If μ_i is a left Haar measure in G_i for $i = 1, 2$, then there is a unique left Haar measure μ_G on G such that

$$\int_G f_1 \otimes f_2(g) d\mu_G(g) = \int_{G_1} f_1(g_1) d\mu_1(g_1) \int_{G_2} f_2(g_2) d\mu_2(g_2),$$

usually denoted as $\mu_G = \mu_1 \otimes \mu_2$.

2.2 The Module of a Group

Of course, the discussion from the previous subsection holds if we replace ‘right’ by ‘left’ throughout. At this point it is therefore natural to ask whether a left Haar integral I on G is also a right Haar integral. This important consideration motivates the following definition.

Definition 2.4. The group G is *unimodular* if any left Haar integral on G is also a right Haar integral.

As a first observation, we note that if the group G is abelian, then $\lambda_g f = \rho_{g^{-1}} f$ and therefore G is unimodular. However, for general groups this is not the case.

To investigate this, choose some left Haar measure μ_G on G , and consider the functional

$$\begin{aligned} I_g : C_c^\infty(G) &\longrightarrow \mathbb{C} \\ f &\longmapsto \int_G f(xg) d\mu_G(x). \end{aligned}$$

In other words, if I is the associated left Haar integral of μ_G , then $I_g(f) = I(\rho_g f)$. Since the actions of G on $C_c^\infty(G)$ by left and right translation commute,

$$I_g(\lambda_h f) = I(\rho_g \lambda_h f) = I(\lambda_h \rho_g f) = I(\rho_g f) = I_g(f)$$

and so I_g is also a left Haar integral. Therefore, there is a unique $\delta_G(g) \in \mathbb{R}_+^\times$ such that $\delta_G(g) I_g(f) = I(f)$ for all $f \in C_c^\infty(G)$. In the integral notation, this means that

$$\delta_G(g) \int_G f(xg) d\mu_G(x) = \int_G f(x) d\mu_G(x)$$

for all $f \in C_c^\infty(G)$. Moreover, the map δ_G also interacts predictably with the left Haar measure. If S is an open compact subset of G and $f = \Gamma_S$ is its characteristic function then one obtains that

$$\delta_G(g) \mu_G(Sg) = \mu_G(S),$$

which also uniquely identifies $\delta_G(g)$.

Lemma 2.5. *The map $\delta_G : G \rightarrow \mathbb{R}_+^\times$ is a homomorphism independent of the choice of left Haar integral I and it is trivial on any open compact subgroup of G . In particular, δ_G is a character of G .*

Proof. By above, we have that

$$\delta_G(gh) I(\rho_{gh} f) = I(f) = \delta_G(g) I(\rho_g f) = \delta_G(g) \delta_G(h) I(\rho_h \rho_g f)$$

for any $g, h \in G$ and $f \in C_c^\infty(G)$. By uniqueness of δ_G and the fact that $\rho_{gh} = \rho_g \rho_h$, it follows that δ_G is a homomorphism. The fact that it is independent of the left Haar measure follows immediately from its definition and Proposition 2.2. If K is an open compact subgroup of G and $k \in K$, then by choosing $f = \Gamma_K$ to be the characteristic function of K , it follows that $\rho_k f = f$ and therefore $\delta_G(k) = 1$. \square

The character $\delta_G : G \rightarrow \mathbb{C}$ is denoted as the *module* of G , and its importance relies on the following result.

Lemma 2.6. *Let G be a locally profinite group and $\delta_G : G \rightarrow \mathbb{C}$ be its module. Then G is unimodular if and only if δ_G is trivial.*

Proof. Let I be a left Haar integral on G . Then G is unimodular if and only if I is a right Haar integral. This is equivalent to $I(f) = I(\rho_g f) = I_g(f) = \delta(g)^{-1} I(f)$ for every $g \in G$. But this is clearly equivalent to δ_G being trivial. \square

Finally, when the group G is not unimodular, the module δ_G gives a canonical relationship between left and right Haar integrals.

Lemma 2.7. *Let I be a left Haar integral on G with associated left Haar measure μ_G . If δ_G is the module of G , then the functional*

$$J : C_c^\infty(G) \longrightarrow \mathbb{C}$$

$$f \longmapsto \int_G \delta_G(x)^{-1} f(x) d\mu_G(x)$$

is a right Haar integral for G .

Proof. The functional J can also be expressed as $J(f) = I(\delta_G^{-1}f)$. We note that $\delta_G^{-1}\rho_g(f) = \delta_G(g)\rho_g\delta_G^{-1}f$ as elements of $C_c^\infty(G)$ for all $g \in G$ and $f \in C_c^\infty(G)$. Hence,

$$J(\rho_g f) = I(\delta_G^{-1}\rho_g f) = \delta_G(g)I(\rho_g\delta_G^{-1}f) = I(\delta_G^{-1}f) = J(f)$$

for every $g \in G$ and $f \in C_c^\infty(G)$, as desired. \square

2.3 Positive Semi-invariant Measures and the Duality Theorem

To classify the principal series representations of $\mathrm{GL}_2(F)$ in the following section, one needs to understand the interaction between the induction and the duality functor for smooth representations of locally profinite groups and their closed groups. To this aim, we need to develop one last bit of machinery from measure theory called *positive semi-invariant measures*, which generalise the notion of Haar measures.

Let G be a locally profinite group and let H be a closed subgroup. Fix some character θ of H and consider the space of functions $f : G \rightarrow \mathbb{C}$ that are G -smooth under right translation, are compactly supported modulo H and satisfy

$$f(hg) = \theta(h)f(g), \quad h \in H, g \in G.$$

This space is in fact the compact induction $c\text{-Ind}_H^G \theta$, but in analogy to $C_c^\infty(G) = c\text{-Ind}_e^G \mathbb{1}$ we denote it as $C_c^\infty(H \backslash G, \theta)$. At this point it is natural to ask whether there exists some non-zero linear functional $I_\theta : C_c^\infty(H \backslash G, \theta) \rightarrow \mathbb{C}$ such that $I_\theta(\rho_g f) = I_\theta(f)$ for all $g \in G$. As it turns out, this is not possible and there is a simple criterion to determine when it is possible.

Proposition 2.8. *Let $\theta : H \rightarrow \mathbb{C}^\times$ be a character of H . Then there exists a non-zero linear functional $I_\theta : C_c^\infty(H \backslash G, \theta) \rightarrow \mathbb{C}$ such that $I_\theta(\rho_g f) = I_\theta(f)$ for all $g \in G$ and $f \in C_c^\infty(H \backslash G, \theta)$ if and only if $\theta\delta_H = \delta_G|_H$.*

Furthermore, when this holds, the functional I_θ is uniquely determined up to a constant.

Proof. [BH06, 3.4 Proposition] \square

We remark that this is a generalisation of Proposition 2.2; indeed, by setting $H = \{e\}$ one recovers the usual right Haar integral on G . Similarly to the above case, when $\theta = \delta_H^{-1}\delta_G|_H$, one commonly expresses the functional I_θ with the integral notation

$$I_\theta(f) = \int_{H \backslash G} f(g) d\mu_{H \backslash G}(g), \quad f \in C_c^\infty(H \backslash G, \theta),$$

where $\mu_{H \setminus G}$ is called a *positive semi-invariant measure* on $H \setminus G$. Also, since such a θ for which Proposition 2.8 holds is uniquely defined, it is common to write $\delta_{H \setminus G}$ for $\delta_H^{-1} \delta_G|_H$. We now have all the required machinery to describe the Duality Theorem.

Theorem 2.9. *Let H be a closed subgroup of a locally profinite group G and let μ be a positive semi-invariant measure on $H \setminus G$. Let (σ, W) be a smooth representation of H . Then there is a natural isomorphism*

$$\left(c\text{-Ind}_H^G \sigma \right)^\vee \cong \text{Ind}_H^G (\delta_{H \setminus G} \otimes \check{\sigma}),$$

which only depends on the choice of μ .

Proof. We sketch a proof to motivate why one would expect $\delta_{H \setminus G}$ to appear. For a detailed proof, check [BH06,] Throughout, we view the action of $\delta_{H \setminus G} \otimes \check{\sigma}$ naturally on \check{W} (where $\check{\sigma}$ acts). For $\phi \in c\text{-Ind}_H^G \sigma$ and $\Phi \in \text{Ind}_H^G \delta_{H \setminus G} \otimes \check{\sigma}$, we have that $\phi(g) \in W$ and $\Phi(g) \in \check{W}$ for any $g \in G$. We can then consider the function

$$f : g \longmapsto \langle \Phi(g), \phi(g) \rangle, \quad g \in G$$

where $\langle \cdot, \cdot \rangle$ is the standard evaluation pairing on $\check{W} \times W$. This function satisfies

$$f(hg) = \langle \Phi(hg), \phi(hg) \rangle = \delta_{H \setminus G}(h) \langle \check{\sigma}(h) \Phi(g), \sigma(h) \phi(g) \rangle = \delta_{H \setminus G}(h) \langle \Phi(g), \phi(g) \rangle = \delta_{H \setminus G}(h) f(g) \quad h \in H, g \in G.$$

Therefore, there is a well-defined pairing

$$\begin{aligned} \Psi : \text{Ind}_H^G (\delta_{H \setminus G} \otimes \check{\sigma}) \times c\text{-Ind}_H^G \sigma &\longrightarrow \mathbb{C}, \\ (\Phi, \phi) &\longmapsto \int_{H \setminus G} \langle \Phi(x), \phi(x) \rangle d\mu(x). \end{aligned}$$

Crucially, this pairing is G -invariant. Indeed,

$$\Psi(\rho_g \Phi, \rho_g \phi) = \int_{H \setminus G} \langle \Phi(xg), \phi(xg) \rangle d\mu(x) = \int_{H \setminus G} \langle \Phi(x), \phi(x) \rangle d\mu(x) = \Psi(\Phi, \phi)$$

by right translation invariance of the positive semi-invariant measure on H/G . This induces a G -homomorphism $\text{Ind}_H^G (\delta_{H \setminus G} \otimes \check{\sigma}) \rightarrow \left(c\text{-Ind}_H^G \sigma \right)^\vee$. The remaining of the proof consists on proving that this is an isomorphism.

The main idea is that the G -homomorphism above is an isomorphism if and only if the following holds.

Lemma 2.10. *The above pairing identifies $\left(\text{Ind}_H^G (\delta_{H \setminus G} \otimes \check{\sigma}) \right)^K$ bijectively with the linear dual of $\left(c\text{-Ind}_H^G \sigma \right)^K$.*

Proof. We omit the proof of this result. The advantage is that one can explicitly describe a canonical basis for each space, which are canonically identified by the pairing. For a complete description, check [BH06, 3.5 Lemma 2]. □

This concludes the proof of the Duality Theorem. □

3 Hecke Algebras

[Possibly put this whole section at the end with modular forms. I think we only need the computation of δ_B from here.]

In this section, we define the Hecke algebra $\mathcal{H}(G)$ associated to a locally profinite (unimodular) group G and explain how to switch between smooth representations of G and smooth modules of $\mathcal{H}(G)$. Under certain conditions on G we consider a particular subalgebra of $\mathcal{H}(G)$; the unramified Hecke algebra $\mathcal{H}(G, K)$, which turns out to be commutative by the Satake isomorphism. We use as reference Chapter 4 of [BH06] and Chapter 5 of [GH24].

If G is a finite group, representations of G are the same as $\mathbb{C}[G]$ -modules. We want to extend this notion to smooth representations of locally profinite groups, where we need to correctly interpret the group algebra.

Let G be a locally profinite unimodular group and K an open compact subgroup of G . Let $C_c^\infty(G)$ be the space of locally constant compactly supported functions $G \rightarrow \mathbb{C}$ and $C_c^\infty(G//K)$ the K bi-invariant subspace.

These are naturally \mathbb{C} -vector spaces and we endow them with an associative (not necessarily unital) ring structure coming from convolution

$$f * h(g) := \int_G f(x)h(x^{-1}g)dx$$

where we fix a Haar measure $\mu = dx$ on G .

When G is discrete this is the usual product on $\mathbb{C}[G]$.

Definition 3.1. Let $\mathcal{H}(G)$ and $\mathcal{H}(G, K)$ denote $C_c^\infty(G)$ and $C_c^\infty(G//K)$ with the algebra structure specified above. We call $\mathcal{H}(G)$ the Hecke algebra of G .

We study these algebras in more detail:

The element $e_K = \mu(K)^{-1}\mathbb{1}_K \in \mathcal{H}(G)$ is idempotent and we have the property that

$$e_K * f = f \Leftrightarrow f \text{ is } K \text{ left invariant}.$$

Thus $\mathcal{H}(G, K) = e_K * \mathcal{H}(G) * e_K$, and this subalgebra now has a unit e_K . The compactness of K ensures $e_K \in C_c^\infty(G)$.

By Lemma 5.2.1 of [GH24], $\mathcal{H}(G)$ is spanned by indicator functions of K' -double cosets, where K' ranges over all compact open subgroups of G . If we normalise these indicator functions by defining

$$[K\alpha K] = \mu(K)^{-1}\mathbb{1}_{K\alpha K},$$

then we have the formula

$$[K\alpha K] * [K\beta K] = \sum_{i,j} [K\alpha_i\beta_j K]$$

where $K\alpha K = \sqcup K\alpha_i$ and $K\beta K = \sqcup \beta_j K$. This determines multiplication in the Hecke algebra.

3.1 Smooth representations and $\mathcal{H}(G)$ -modules

We now explain how the concepts of smooth representations of G and smooth modules over $\mathcal{H}(G)$ are interchangeable. To define these smooth modules, we note that the Hecke algebra $\mathcal{H}(G)$ does not in general have a unit. Consequently, not every $\mathcal{H}(G)$ -module M satisfies $\mathcal{H}(G)M = M$.

Definition 3.2. We say that a $\mathcal{H}(G)$ -module M is smooth if $\mathcal{H}(G)M = M$.

Definition 3.3. From a representation V of G we define the action of $\mathcal{H}(G)$ on V via

$$f \cdot v := \int_G f(g)g \cdot v dg.$$

This can be viewed as a weighted average of the action of G on v , where the weighting is described by $f \in C_c^\infty(G)$. The integral defines an element of V when $f \in C_c^\infty(G)$ as the integral reduces to a finite sum.

Lemma 3.4. *Under this action, $e_K \in \mathcal{H}(G)$, for $K \leq G$ a compact open, is the projection $V \rightarrow V^K$ onto the K -invariants of V . In particular, e_K is an idempotent element of $\mathcal{H}(G)$, it is the identity element of $\mathcal{H}(G, K)$, and V^K is a $\mathcal{H}(G, K)$ -module.*

Proof. Let $V(K) \leq V$ be the subspace spanned by vectors of the form $k \cdot v - v$. Since e_K is invariant under K -translation, it is zero on $V(K)$. The normalisation of e_K is such that e_K is the identity on V^K , and this implies the result. \square

Proposition 3.5. *A representation V of G is smooth if and only if it is a smooth $\mathcal{H}(G)$ -module.*

Proof. If V is a smooth representation, then any $v \in V$ is K -invariant for some compact open K , and so $v = e_K \cdot v$. This implies that V is a smooth $\mathcal{H}(G)$ -module. Conversely, $\mathcal{H}(G)$ is the union of $e_K * \mathcal{H}(G) * e_K = \mathcal{H}(G, K)$ over all compact open K , and so if V is a smooth $\mathcal{H}(G)$ -module then any $v \in V$ is of the form $e_K * f * e_K \cdot v'$ for some K, f, v' . Then $e_K \cdot v = v$ and so $v \in V^K$. \square

So we can view smooth representations of G as smooth $\mathcal{H}(G)$ -module. In the other direction, given M a smooth $\mathcal{H}(G)$ -module, we have

$$\mathcal{H}(G) \otimes_{\mathcal{H}(G)} M = M$$

by smoothness. We can then view M as a smooth G representation by letting G act on the first factor by left translation. Concretely, if $m \in M$ there exists K such that $m \in \mathcal{H}(G, K)M = e_K * \mathcal{H}(G) * e_K M$. Then $m = e_K \cdot m'$ for some $m' \in M$ and therefore $e_K \cdot m = m$ because e_K is idempotent in $\mathcal{H}(G)$. Then define

$$g \cdot m := \mu(K)^{-1} \mathbb{1}_{gK} \cdot m,$$

where this is independent of K due to the normalisation factor $\mu(K)^{-1}$.

3.2 Information in the K -invariants V^K

For a smooth representation V of G it is often easier to study the K -invariants V^K for compact open subgroups K of G .

Lemma 3.6. *A smooth representation V of G is irreducible if and only if each V^K is either 0 or a simple $\mathcal{H}(G, K)$ -module for all compact open $K \leq G$.*

Proof. Suppose V is irreducible. If we had $0 \neq M \subset V^K$ a $\mathcal{H}(G, K)$ -module, then $0 \neq \mathcal{H}(G)M \subset V$ as smooth $\mathcal{H}(G)$ -modules. Since smooth $\mathcal{H}(G)$ -modules are the same as smooth G -representations, and V is irreducible, we deduce $\mathcal{H}(G)M = V$. So then

$$V^K = e_K V = e_K * \mathcal{H}(G)M = e_K * \mathcal{H}(G) * e_K M = \mathcal{H}(G, K)M = M$$

which implies the result.

If V is not irreducible, and $W \neq 0$ is a proper subrepresentation, pick $v \in V - W$. By smoothness, there exists K such that $v \in V^K$, and also $W^K \neq 0$, but then $v \notin W^K$ so that V^K is not 0 or simple. \square

The next result tells us that for any K , any smooth representation V of G is determined by V^K with its structure as a $\mathcal{H}(G, K)$ -module, provided $V^K \neq 0$.

Proposition 3.7. *The map $V \mapsto V^K$ induces a bijection between*

- *equivalence classes of irreducible smooth representations V of G with $V^K \neq 0$;*
- *isomorphism classes of simple (by definition nonzero) $\mathcal{H}(G, K)$ -modules.*

Proof. Proposition 4.3 of [BH06]. \square

3.3 Unramified representations of G

It is interesting to study the smooth representations V with $V^K \neq 0$ as above. For example, in an automorphic representation, Flath's theorem ([GH24] Section 5.7) allows us to decompose into local factors, and furthermore tells us that almost all such local representations are unramified in the following sense:

Definition 3.8. We consider the case $G = \mathrm{GL}_2(F)$. We say that an irreducible smooth representation V of G is unramified if $V^K \neq 0$ for $K = \mathrm{GL}_2(\mathcal{O}_F)$. See Section 5.5 of [GH24] for a more general definition for reductive groups.

For the remainder of this subsection we work in the context of $G = \mathrm{GL}_2(F)$ and $K = \mathrm{GL}_2(\mathcal{O}_F)$ for simplicity. The results generalise to reductive groups G as in Sections 5.5 and 7.1 of [GH24].

Definition 3.9. For K as above, $\mathcal{H}(G, K)$ is called the unramified Hecke algebra of G .

An application of the Satake isomorphism ([GH24] Theorem 5.5.1) tells us that in this unramified case, the unramified Hecke algebra $\mathcal{H}(G, K)$ is commutative. It follows that if V is K -unramified (in particular irreducible) then V^K is 1-dimensional by Lemma 3.6. Thus $\mathcal{H}(G, K)$ acts on V^K via scaling, called the Hecke character of V .

Definition 3.10. The Hecke character (with respect to K) of a smooth representation (π, V) of G is the \mathbb{C} -linear map

$$\begin{aligned}\mathcal{H}(G, K) &\rightarrow \mathbb{C} \\ f &\mapsto \pi(f)\end{aligned}$$

defined by $f \cdot v =: \pi(f)v$ for any $v \in V^K$.

We give an alternative proof of Proposition 3.7.

Proposition 3.11. *Let $K \leq G$ be a compact open subgroup. If V_1, V_2 are irreducible smooth representations of G such that V_1^K and V_2^K are nonzero and isomorphic as $\mathcal{H}(G, K)$ -modules, then $V_1 \cong V_2$. In particular, unramified representations are determined by their Hecke characters.*

Proof. This is Proposition 7.1.1 of [GH24]. The idea is to extend an isomorphism

$$I : V_1^K \rightarrow V_2^K$$

to a G -intertwining map $V_1 \rightarrow V_2$ of $\mathcal{H}(G)$ -modules. By irreducibility, $V_i = \mathcal{H}(G)V_i^K$. Take an element $\pi_1(f) \cdot \phi \in V_1$, with $f \in \mathcal{H}(G), \phi \in V_1^K$, then the obvious choice is to map this to $\pi_2(f) \cdot I(\phi)$. Provided this is well defined, this is a nonzero homomorphism of $\mathcal{H}(G)$ -modules, so irreducibility of V_1, V_2 implies this is an isomorphism $V_1 \cong V_2$.

We now check that this is well defined. Suppose $\pi_1(f_1)\phi_1 = \pi_1(f_2)\phi_2$. Since V_1^K is irreducible, there exists $f_3 \in \mathcal{H}(G, K)$ such that $\pi_1(f_3)\phi_1 = \phi_2$, and so also $\pi_2(f_3)I(\phi_1) = I(\phi_2)$ since I is $\mathcal{H}(G, K)$ -intertwining. Then $\pi_1(f_1)\phi_1 = \pi_1(f_2 * f_3)\phi_1$. Thus it suffices to show that if $\pi_1(f)\phi = 0$ then $\pi_2(f)I(\phi) = 0$. We exploit the $\mathcal{H}(G, K)$ -intertwining of I (for the second implication below). For all $f_1 \in \mathcal{H}(G)$ we have:

$$\pi_1(f)\phi = 0 \Rightarrow \pi_1(e_K * f_1 * f * e_K)\phi = 0 \Rightarrow \pi_2(e_K * f_1 * f * e_K)I(\phi) = \pi_2(e_K * f_1 * f)I(\phi) = 0.$$

By Lemma 3.4, e_K acts on V_2 by projection to V_2^K . If $\pi_2(f)I(\phi) \neq 0$, then $\pi_2(f_1)\pi_2(f)I(\phi)$, over all $f_1 \in \mathcal{H}(G)$, generates V_2 by irreducibility. The image under $\pi_2(e_K)$ is the exactly V^K , which is nonzero, contradicting the implication above. \square

3.4 Example computation of Hecke operators for $\mathrm{GL}_2(F)$

[I haven't checked this subsection. Some parts might be more suitable for a section on modular forms. The computation of the modular character of B will be needed in the main text. And the last proposition naturally goes with the unramified representations above.]

Let $G = \mathrm{GL}_2(F)$ and $K = \mathrm{GL}_2(\mathcal{O})$ for F a nonarchimedean local field with uniformiser ϖ . We have the Cartan decomposition

$$G = \bigsqcup_{a \geq b \in \mathbb{Z}} K \begin{pmatrix} \varpi^a & \\ & \varpi^b \end{pmatrix} K.$$

Let $S = K \begin{pmatrix} \varpi & \\ & \varpi \end{pmatrix} K$ and $T = K \begin{pmatrix} \varpi & \\ & 1 \end{pmatrix} K$, viewed as elements of $\mathcal{H}(G, K)$ via their indicator functions.

Lemma 3.12. *The unramified Hecke algebra is $\mathcal{H}(G, K) \cong \mathbb{C}[S, S^{-1}, T]$. In particular, this is commutative.*

Proof. This is some induction argument using the formula for convolutions of these indicator functions. \square

Remark 3.13. This fits into a general phenomenon - if G is unramified and K is a hyperspecial subgroup then the Satake isomorphism implies that the unramified Hecke algebra $\mathcal{H}(G, K)$ is always commutative.

Later we will be interested in principal series representations, which are representations of G coming from parabolic induction. So let $\chi = \begin{pmatrix} \chi_1 & \\ & \chi_2 \end{pmatrix}$ be a character of the torus T , and consider the normalised induced representation

$$I(\chi) = \mathrm{Ind}_B^G \left(\chi \otimes \delta_B^{-1/2} \right)$$

where we recall that this is the space of functions $G \rightarrow \mathbb{C}$ with $f(bg) = \chi(b)\delta_B^{-1/2}(b)f(g)$ for $b \in B$.

We briefly discuss the module character δ_B . Although G is unimodular (see Bushnell-Henniart Section 7.5), the Borel subgroup is not. We have $B = NT$ with $N \cong F$, $T \cong F^\times \times F^\times$ and N normal in B . The failure of B to be unimodular is a consequence of T and N not commuting. We can then define a linear function I on $C_c^\infty(B) = C_c^\infty(T) \otimes C_c^\infty(N)$ by

$$I(\Phi) = \int_T \int_N \Phi(tn) dt dn$$

using Haar measures on T and N .

Proposition 3.14. *I is a left Haar integral on B .*

Proof. Let $b = sm \in TN$. By left invariance of dt we have

$$\int_T \int_N \Phi(smtn) dt dn = \int_T \int_N \Phi(mtn) dt dn = \int_T \int_N \Phi(tt^{-1}mntn) dt dn.$$

Since we integrate N first, we are integrating over fixed values of t so that $t^{-1}mt \in N$ is just constant, so left invariance of dn let's us pull out the $t^{-1}mt$ factor. \square

Proposition 3.15. *The modular character δ_B of the group B is*

$$\delta_B : tn \mapsto |t_2/t_1|, \quad n \in N, t = \begin{pmatrix} t_1 & 0 \\ 0 & t_2 \end{pmatrix} \in T$$

Proof. By a similar argument as above, we have

$$\int_T \int_N \Phi(tnsm) dt dn = \int_T \int_N \Phi(tss^{-1}nsm) dt dn = \int_T \int_N \Phi(ts^{-1}ns) dt dn.$$

Identifying $N \cong F$ this is

$$\int_T \int_N \Phi\left(t \cdot \begin{pmatrix} 1 & s_1^{-1}xs_2 \\ 0 & 1 \end{pmatrix}\right) d\mu_F(x) = |s_1/s_2| \int_T \int_N \Phi(tn) dt dn$$

so by definition of the module character we have $\delta_B(sm) = |s_2/s_1|$. \square

Going back to our principal series representation, the following proposition computes the action of the unramified Hecke algebra on the K -invariant subspace:

Proposition 3.16. *Let $\chi : T \rightarrow \mathbb{C}^\times$ be an unramified character of the torus (meaning trivial on $\begin{pmatrix} \mathcal{O}^\times & \\ & \mathcal{O}^\times \end{pmatrix}$) and consider the normalised parabolic induction*

$$I(\chi) = \text{Ind}_B^G(\chi \otimes \delta_B^{-1/2}).$$

For $K = \text{GL}_2(\mathcal{O})$ as usual, the space $I(\chi)^K$ is 1-dimensional. As a $\mathcal{H}(G, K)$ -module this is determined by the actions of S and T . Since χ is unramified we know $\chi_1(z) = \alpha^{v_F(z)}$ and $\chi_2(z) = \beta^{v_F(z)}$ for some $\alpha, \beta \in \mathbb{C}^\times$. Then S acts on $I(\chi)^K$ by scaling by $\alpha\beta$ and T acts by scaling by $q^{1/2}(\alpha + \beta)$.

Proof. We have the Iwasawa decomposition $G = BK$ so that the functions $f \in I(\chi)^K$ satisfy

$$f(bk) = f(b) = \chi(b)\delta_B^{-1/2}(b) \cdot f(1)$$

with $f(1) \in \mathbb{C}$, so the space is 1-dimensional spanned by $\hat{f}(bk) = \chi(b)\delta_B^{-1/2}(b)$.

The action of S is given by:

$$\begin{aligned} S \cdot f &= \mu(K)^{-1} \int_G \mathbb{1}_{K \begin{pmatrix} \varpi & \\ & \varpi \end{pmatrix} K}(g) g \cdot f dg \\ &= \mu(K)^{-1} \int_K \begin{pmatrix} \varpi & \\ & \varpi \end{pmatrix} k \cdot f dk \\ &= \begin{pmatrix} \varpi & \\ & \varpi \end{pmatrix} \cdot f \\ &= \chi \left(\begin{pmatrix} \varpi & \\ & \varpi \end{pmatrix} \right) \delta_B^{-1/2} \left(\begin{pmatrix} \varpi & \\ & \varpi \end{pmatrix} \right) f \\ &= \alpha\beta f \end{aligned}$$

because $K \begin{pmatrix} \varpi & \\ & \varpi \end{pmatrix} K = \begin{pmatrix} \varpi & \\ & \varpi \end{pmatrix} K$.

And for T we pick coset representatives for $K \begin{pmatrix} \varpi & a \\ & 1 \end{pmatrix} K / K$ given by $\begin{pmatrix} \varpi & a \\ & 1 \end{pmatrix}$ and $\begin{pmatrix} 1 & \\ & \varpi \end{pmatrix}$, where a ranges over representatives of \mathcal{O}/ϖ . Writing down the integral for the action of T we decompose this into a sum over these left cosets and we deduce that T acts by

$$\chi_2(\varpi)|\varpi|^{-1/2}f + \sum_{a \in \mathcal{O}/\varpi} \chi_1(\varpi)|\varpi|^{1/2}f = q^{1/2}(\alpha + \beta)$$

since, for example, $\chi \left(\begin{pmatrix} \varpi & a \\ & 1 \end{pmatrix} \right) = \chi_1(\varpi) = \alpha$ and $\delta_B^{-1/2} \left(\begin{pmatrix} \varpi & a \\ & 1 \end{pmatrix} \right) = |\varpi|^{1/2}$. □

Remark 3.17. If we know the action of S, T on $I(\chi)^K$ for some unramified character χ of the torus T , then we can recover $\alpha, \beta \in \mathbb{C}^\times$ from the roots of the Satake polynomial $X^2 - q^{-1/2}TX + S \in \mathcal{H}(G, K)[X]$.

4 Principal series representations of GL_2

Let F be a nonarchimedean local field, $G = \text{GL}_2(F)$, and $B = \left\{ \begin{pmatrix} a & b \\ & d \end{pmatrix} \mid a, d \in F^\times, b \in F \right\}$ the Borel subgroup of upper triangular matrices, so that $B = N \rtimes T$ for $T = \left\{ \begin{pmatrix} a & 0 \\ & d \end{pmatrix} \mid a, d \in F^\times \right\} \cong F^\times \times F^\times$ and $N = \left\{ \begin{pmatrix} 1 & b \\ & 1 \end{pmatrix} \mid b \in F \right\}$.

$b \in F\} \cong F$. Between N and B we also have the mirabolic subgroup $M = \{ \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} \mid a \in F^\times, b \in F \}$ with $M/N \cong F^\times$.

In studying the local Langlands correspondence, we want to understand all the irreducible smooth representations of G . One method for producing representations of G is by induction from a subgroup of G . Typically one takes this subgroup to be ‘parabolic’; in our case there is one nontrivial parabolic, namely B . From our decomposition $B = N \rtimes T$ (more generally we have a so-called Levi decomposition) we see that we can produce representations of B by inflating representations of the torus T . Since $T \cong F^\times \times F^\times$, the irreducible representations of T are products of characters of F^\times , which are relatively easy to get a handle on.

Definition 4.1. For $\chi : T \rightarrow \mathbb{C}^\times$ a character of the torus, we say that the representation $\text{Ind}_B^G \chi$ is a parabolically induced representation. A principal series representation is an irreducible subrepresentation of a parabolically induced representation.

In this section, we will only concern ourselves with classifying the principal series representations of G . This means that we must understand how $\text{Ind}_B^G \chi$ decomposes into irreducible representations of G , and also study the morphisms between them using Frobenius reciprocity.

To understand these decompositions, we want to study how they decompose into irreducibles over a less unwieldy subgroup of G , such as B . Note that restricting $\text{Ind}_B^G \chi$ to B is analogous to applying Mackey theory in the finite group context. It turns out that the $\text{Ind}_B^G \chi$ do not decompose any further over M than over B . On the other hand, the representation theory of M is very easy to classify - the combination of these two observations is what makes the mirabolic subgroup so ‘miraculous’. To get representations of M we can induce from characters of N , or inflate from $M/N \cong F^\times$. There are many characters of $N \cong F$, in fact these are in bijection with F [REFER TO SECTION 1]. The key property of M is that conjugation by M acts transitively on these characters ψ , which greatly simplifies the representation theory of M coming via induction from N . The mirabolic M is also small enough that this induction, together with the characters of F^\times , give all irreducible representations of M .

In this section, we begin by studying the representations of N and introducing the Jacquet functor, before discussing representations of M . From there we determine that parabolically induced representations of G decompose over M with length at most 3. Theorem 4.21 gives the decomposition of $\text{Ind}_B^G \chi$ into irreducible representations of G , and then Theorem 4.29 lists the isomorphism classes of principal series representations. The presentation follows sections 8 and 9 of [BH06].

4.1 Representations of N

We first study the representation theory of $N \cong F$. This is an abelian group so, by Schur’s lemma, all irreducible representations are characters (Corollary 2.6.2 [BH06]). For finite abelian groups, any representation V decomposes into a direct sum of characters. This is no longer true when $N \cong F$ is infinite, but it is still true that any vector in V is nonzero in some quotient on which N acts via a character. To formalise this, we define

Notation 4.2. Let V be a smooth representation of N and θ a character of N . Let $V(\theta) \leq V$ be the subspace

spanned by $\{n \cdot v - \theta(n)v \mid n \in N, v \in V\}$. Set $V_\theta = V/V(\theta)$ so that N acts on V_θ by θ . When θ is trivial we write $V(N)$ and V_N respectively.

The following is a useful equivalent definition of $V(\theta)$:

Lemma 4.3. *The vector $v \in V$ lies in $V(\theta)$ if and only if*

$$\int_{N_0} \theta(n)^{-1} n \cdot v dn = 0$$

for some compact open subgroup N_0 of N .

In the lemma we restrict to compact opens for the integral to be well defined.

Proof. [BH06, Lemma 8.1]. □

Corollary 4.4. *The functor $V \mapsto V_\theta$ from smooth representations of N to complex vector spaces is exact.*

Proof. One checks formally that the functor is right exact. For left exactness we need to show that if $f : V \hookrightarrow V'$ is injective then $V_\theta \hookrightarrow V'_\theta$ is injective. If $v \in V$ with $f(v) \in V'(\theta)$, then

$$\int_{N_0} \theta(n)^{-1} n \cdot f(v) dn = 0$$

for some N_0 by the above lemma. Since f is compatible with the action of N , we can pull f out of the integral so that the injectivity of f implies

$$\int_{N_0} \theta(n)^{-1} n \cdot v dn = 0.$$

We deduce that $v \in V(\theta)$ by the above lemma. □

Proposition 4.5. *Let V be a smooth representation of N . For any $v \neq 0$ in V , there exists a character θ of N such that $v \notin V(\theta)$.*

Proof. [BH06, Proposition 8.1]. □

Corollary 4.6. *If V is a smooth representation of N such that $V_\theta = 0$ for all θ then $V = 0$.*

4.2 Representations of M

Now we consider V an irreducible smooth representation of M .

Lemma 4.7. *The subspace $V(N) \leq V$ is a representation of M , and so V_N is as well. Moreover, $S = \{ \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \mid a \in F^\times \}$ permutes the subspaces $V(\theta)$ with $\theta \neq 1$ transitively, and hence the V_θ are isomorphic as vector spaces.*

Proof. The first claim comes from the computation

$$mn \cdot v - m \cdot v = n' m \cdot v - m \cdot v$$

for some $n' \in N$, using the fact that $N \triangleleft M$. For the second claim we have the computation

$$s(nv - \theta(n)v) = sns^{-1} \cdot sv - \theta(s^{-1}(sns^{-1})s)sv = n' \cdot sv - \theta(s^{-1}n's)sv$$

where $n' = sns^{-1} \in N$. Hence $sV(\theta) = V(\theta')$ where $\theta'(n) := \theta(s^{-1}ns)$. Now the computation

$$\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a^{-1} & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & ax \\ 0 & 1 \end{pmatrix}$$

together with [ADDITIVE DUALITY] implies the claim. \square

Theorem 4.8. *Let (π, V) be an irreducible smooth representation of M . Either*

- $\dim V = 1$ and π is the inflation of a character of $M/N \cong F^\times$, or
- $\dim V = \infty$ and $\pi \cong c\text{-Ind}_N^M \theta$, for any nontrivial character θ of N .

This itself follows from the following theorems. To compare V and $c\text{-Ind}_N^M \theta$, it is more natural to compare V and $\text{Ind}_N^M V_\theta$. By Frobenius reciprocity,

$$\text{Hom}_N(V, V_\theta) \cong \text{Hom}_M(V, \text{Ind}_N^M V_\theta).$$

Let $q_* : V \rightarrow \text{Ind}_N^M(V_\theta)$ be the image of the quotient map $q : V \rightarrow V_\theta$.

Theorem 4.9. *The M -homomorphism $q_* : V \rightarrow \text{Ind}_N^M V_\theta$ induces an isomorphism $V(N) \cong c\text{-Ind}_N^M V_\theta$.*

Proof. [BH06, Theorem 8.3]. \square

Theorem 4.10. *For any nontrivial character θ of N , the smooth representation $c\text{-Ind}_N^M \theta$ of M is irreducible.*

Proof. [BH06, Corollary 8.2] \square

Proof of Theorem 4.8. If V is an irreducible smooth representation of M , then either $V(N) = 0$ or $V(N) = V$. In the former case N acts trivially on V , so the action of M factors through $M/N \cong F^\times$. Schur's lemma implies that V is a character of M factoring through M/N .

In the latter case, $V_N = 0$, so we must have $V_\theta \neq 0$ for all nontrivial characters of N by Lemma 4.7 and Corollary 4.6. Thus the M -representation V must have infinite dimension, since there are infinitely many characters θ . Theorem 4.9 implies that $V = V(N)$ is isomorphic to $c\text{-Ind}_N^M V_\theta$, which is a direct sum of copies of $c\text{-Ind}_N^M \theta$. Since $c\text{-Ind}_N^M \theta$ is irreducible by Theorem 4.10, we must have $V \cong c\text{-Ind}_N^M \theta$. \square

4.3 Irreducible principal series representations

Let V be a smooth representation of G . In the preceding subsections, we defined the quotient $V_N = V/V(N)$, called the N -coinvariants of V . As in Lemma 4.7, this is a representation of B (as $N \triangleleft B$). As N acts trivially on V_N , V_N inherits the structure of a representation of $T = B/N$.

Definition 4.11. Let V be a smooth representation of G (or B). The Jacquet module of V at N is the space of N -coinvariants V_N viewed as a representation of T . The Jacquet functor is the functor sending the G -representation (π, V) to the T -representation (π_N, V_N) .

By Corollary 4.4, the Jacquet functor is exact.

If V is a representation of G , and χ is a character of T , then we have by Frobenius Reciprocity that

$$\mathrm{Hom}_G(V, \mathrm{Ind}_B^G \chi) \cong \mathrm{Hom}_B(V, \chi)$$

But since χ as a character of B has trivial N -action, maps $V \rightarrow \chi$ factor through V_N , and we obtain a version of Frobenius reciprocity for the Jacquet module:

$$\mathrm{Hom}_G(V, \mathrm{Ind}_B^G \chi) \cong \mathrm{Hom}_T(V_N, \chi)$$

i.e. the Jacquet module is left adjoint to parabolic induction.

In the classical setting of representations of $\mathbf{G} = \mathrm{GL}_2(k)$ for a finite field k , we have the following dichotomy (where $\mathbf{B}, \mathbf{T}, \mathbf{N}$ are the appropriate subgroups of \mathbf{G}):

Lemma 4.12. *Let (π, V) be an irreducible representation of \mathbf{G} . The following are equivalent:*

1. π contains the trivial character of \mathbf{N}
2. π is isomorphic to a \mathbf{G} -subrepresentation of $\mathrm{Ind}_{\mathbf{B}}^{\mathbf{G}} \chi$ for some character χ of \mathbf{T} inflated to \mathbf{B} .

Proof. [BH06, Lemma 6.3]. □

Returning to $G = \mathrm{GL}_2(F)$, if (π, V) is a smooth representation, the restriction to N is no longer necessarily semisimple because F is of infinite order. We instead replace the condition that $\pi|_N$ contains the trivial character of N with the condition that N acts trivially on some nonzero quotient of V (which is an equivalent condition in the finite field case). This is measured by the Jacquet module V_N . There is the analogous dichotomy which tells us that principal series representations can be identified as the irreducible smooth representations of G with nonzero Jacquet module:

Proposition 4.13. *Let (π, V) be an irreducible smooth representation of G . The following are equivalent:*

1. $V_N \neq 0$
2. π is isomorphic to a G -subrepresentation of $\mathrm{Ind}_B^G \chi$ for some character χ of T inflated to B .

Proof sketch. (2) implies (1) is a consequence of Frobenius reciprocity:

$$\mathrm{Hom}_G(\pi, \mathrm{Ind}_B^G \chi) = \mathrm{Hom}_T(\pi_N, \chi)$$

Given (1), one shows by a technical argument that V_N is finitely generated as a representation of T . An application of Zorn's lemma allows us to construct a maximal T -subspace U of V_N , so that V_N/U is a nonzero irreducible T -representation, and is thus a character χ by Schur's lemma. The above Frobenius reciprocity implies (2). □

Remark 4.14. The same proof holds for the finite field case, where we bypass the technical details in showing (1) implies (2) because any representation of the finite group T admits an irreducible quotient.

Remark 4.15. We ask for a nonzero Jacquet module V_N rather than a trivial N -subrepresentation of V because of the following fact:

Lemma 4.16. *Let (π, V) be an irreducible smooth representation of G with a nonzero vector $v \in V$ fixed by N . Then $\pi = \phi \circ \det$, for some character ϕ of F^\times . In particular, π is one dimensional.*

Proof sketch. The vector v is fixed by N , but also by a compact open subgroup K of G by smoothness. As we are working with F a nonarchimedean local field (as opposed to a finite field), this implies K contains a unipotent lower triangular matrix, and one shows that v is fixed by $\mathrm{SL}_2(F)$. Thus π factors through \det . \square

Once again, let χ be a character of T and let (Σ, X) denote $\mathrm{Ind}_B^G \chi$. We want to study how X decomposes into irreducible G -representations. As mentioned earlier, we will begin by studying their decompositions over B or even M .

To begin with, X will never be irreducible over B because we always have the canonical B -homomorphism $\Sigma \rightarrow \chi$, given by sending $f \mapsto f(1) \in \mathbb{C}$. So we have an exact sequence of B -representations

$$0 \longrightarrow V \longrightarrow X \longrightarrow \mathbb{C} \longrightarrow 0,$$

where $V = \{f \in X \mid f(1) = 0\}$, and B acts on \mathbb{C} via χ . Now we want to understand how V decomposes over B . We have another exact sequence of B -representations,

$$0 \longrightarrow V(N) \longrightarrow V \longrightarrow V_N \longrightarrow 0,$$

so we reduce to studying $V(N)$ and V_N . We will show that $V(N)$ is irreducible over B (and even over M), while V_N will be determined by the Restriction-Induction lemma.

The following lemma makes the structure of V more apparent.

Lemma 4.17. *Let $V = \{f \in X : f(1) = 0\}$. The map*

$$\begin{aligned} V &\rightarrow C_c^\infty(N) \\ f(-) &\mapsto f(w-) \end{aligned}$$

is an N -isomorphism (with N acting by right translation on either side), where $w = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.

Proof. We have the Bruhat decomposition $G = B \sqcup BwN$. Since $f(1) = 0$, and f is induced from B , we must have that f is supported on BwN . G -smoothness of f implies that f is also zero on some compact open $K \leq G$. This will contain $\begin{pmatrix} 1 & 0 \\ \varpi^n \mathcal{O} & 1 \end{pmatrix}$ for some n , so that f vanishes on

$$\begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix} \in Bw \begin{pmatrix} 1 & x^{-1} \\ 0 & 1 \end{pmatrix}$$

for all $x \in \varpi^n \mathcal{O}$. Thus $f(w-)$ is supported on $\begin{pmatrix} 1 & y \\ 0 & 1 \end{pmatrix} \in N$ with $v(y) > -n$ and so is compactly supported. G -smoothness of f also implies that $f(w-)$ is N -smooth. Since f is induced from B and is supported on BwN , the map is injective. Conversely, any $g \in C_c^\infty(N)$ determines $f \in \mathrm{Ind}_B^G \chi$ such that $f(w-) = g$ and $f(B) = 0$. \square

Proposition 4.18. *For V as above, $V(N)$ is irreducible over M (and hence over B). Moreover, $V(N)$ is infinite dimensional.*

Proof. The idea will be to use Theorem 4.9, which tells us $V(N) \cong c\text{-Ind}_N^M V_\theta$. This is irreducible over M (and infinite dimensional) if we can show that V_θ is one dimensional, by Theorem 4.10.

By the above lemma we can identify $V \cong C_c^\infty(N)$ as N -representations. But M also acts via right translation on V (since $BwB = BwN = BwM$), which gives the structure of a M -representation on $C_c^\infty(N)$. We can calculate it explicitly (but we won't need it), where

$$f \left(bw \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \right) = f \left(b \begin{pmatrix} 1 & 0 \\ 0 & a \end{pmatrix} w \begin{pmatrix} 1 & a^{-1}x \\ 0 & 1 \end{pmatrix} \right)$$

tells us that the corresponding $M = F^\times N$ action on $C_c^\infty(N)$ is the composite of right translation by N with the action

$$a \cdot \phi \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} = \chi_2(a) \phi \begin{pmatrix} 1 & a^{-1}x \\ 0 & 1 \end{pmatrix}$$

of $a \in F^\times$.

So now we may consider $V = C_c^\infty(N)$. The benefit is that for this representation, the spaces of coinvariants of characters θ of N are very simple. In particular, the map $f \mapsto \theta f$ is a linear automorphism of $C_c^\infty(N)$ taking $V(N)$ to $V(\theta)$, since

$$n \cdot f - f \mapsto \theta(n \cdot f) - \theta f = \theta(n)^{-1} n \cdot (\theta f) - \theta f \in V(\theta).$$

Hence all the V_θ have the same dimension as $V_N = V/V(N)$, which has dimension 1 (we can see this from the characterisation of $V(N)$ as the zeros of some integral (Lemma 4.3), or from the Restriction-Induction lemma to follow). The result follows from Theorem 4.9 and Theorem 4.10. \square

We turn our attention to the Jacquet module V_N . Recall V fits in the exact sequence

$$0 \longrightarrow V \longrightarrow X = \text{Ind}_B^G \chi \xrightarrow{f \mapsto f(1)} \mathbb{C} \longrightarrow 0$$

of smooth representations of B , where B acts via χ on \mathbb{C} . Since the Jacquet functor is exact, we get the exact sequence

$$0 \longrightarrow V_N \longrightarrow X_N \longrightarrow \mathbb{C} \longrightarrow 0$$

of T -representations. The following lemma determines the structure of V_N as a T -representation. This can be stated in more generality:

Lemma 4.19 (Restriction-Induction lemma). *Let (σ, U) be a smooth representation of T and $(\Sigma, X) = \text{Ind}_B^G \sigma$. Then there is an exact sequence of smooth T representations:*

$$0 \longrightarrow \sigma^w \otimes \delta_B^{-1} \longrightarrow \Sigma_N \longrightarrow \sigma \longrightarrow 0.$$

Here, $\sigma^w(t) := \sigma(wtw)$ for $w = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, so that if σ is the character $\chi_1 \otimes \chi_2$ of T , then $\sigma^w = \chi_2 \otimes \chi_1$.

Proof. The proof of Lemma 4.17 generalises to show that the vector space $V = \{f \in X \mid f(1) = 0\}$ is isomorphic, as N -representations, to the space \mathcal{S} of smooth compactly supported functions $N \rightarrow U$, by identifying f with $f(w-)$.

We can define a map $\mathcal{S} \rightarrow U$ by

$$g = f(w-) \mapsto \int_N f(wn)dn,$$

where this integral is finite since g is compactly supported. By Lemma 4.3, this induces an isomorphism $\mathcal{S}_N \cong U$.

Now V also carries the structure of a B -representation as well, since $BwB = BwN$. We can repeat the same calculation as in the previous proposition, replacing F^\times with $T \cong F^\times \times F^\times$, to compute the action of $B = TN$ on \mathcal{S} . As usual, N acts via right translation. If $t = \begin{pmatrix} t_1 & 0 \\ 0 & t_2 \end{pmatrix} \in T$, then for $\phi \in \mathcal{S}$,

$$t \cdot \phi \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} = \sigma^w(t) \phi \begin{pmatrix} 1 & \frac{t_2}{t_1}x \\ 0 & 1 \end{pmatrix}.$$

Thus the T -representation structure on $U \cong \mathcal{S}_N \cong V_N$ is given by

$$t \cdot \int_N f(wn)dn = \sigma^w(t) \left| \frac{t_1}{t_2} \right| \int_N f(wn)dn,$$

which is $\sigma^w \otimes \delta_B^{-1}$. □

Corollary 4.20. *As a representation of B or M , $\text{Ind}_B^G \chi$ has composition length 3. Two of the factors have dimension 1, and the other is infinite dimensional.*

Proof. This follows from the exact sequences

$$0 \longrightarrow V \longrightarrow \text{Ind}_B^G \mathbb{C} \longrightarrow \chi \longrightarrow 0$$

and

$$0 \longrightarrow V(N) \longrightarrow V \longrightarrow V_N \longrightarrow 0$$

where we saw that $V(N)$ is irreducible and infinite dimensional, and $V_N \cong \chi^w \otimes \delta_B^{-1}$. □

So we understand how $\text{Ind}_B^G \chi$ decomposes into irreducible B -representations, and we want to understand its decomposition into G -representations. Our goal is to prove the following:

Theorem 4.21 (Irreducibility Criterion). *Let $\chi = \chi_1 \otimes \chi_2$ be a character of T and let $X = \text{Ind}_B^G \chi$.*

1. *The representation X of G is irreducible if and only if $\chi_1 \chi_2^{-1}$ is either the trivial character of F^\times , or the character $x \mapsto |x|^2$ of F^\times .*
2. *Suppose X is reducible. Then*

- *the G -composition length of X is 2*
- *one factor has dimension 1, the other is infinite dimensional*
- *X has a 1-dimensional G -subspace exactly when $\chi_1 \chi_2^{-1} = 1$*

- X has a 1-dimensional G -quotient exactly when $\chi_1\chi_2^{-1}(x) = |x|^2$.

We make some comments in preparation for the proof. Suppose X is a reducible representation of G , and X_0 a nonzero proper subrepresentation. If X_0 is finite-dimensional, then its composition factors over B can only consist of the 1-dimensional composition factors of X over B described in Corollary 4.20. If X_0 is infinite dimensional, then it contains the infinite-dimensional B -composition factor of Corollary 4.20, and so the quotient X/X_0 can only consist of the 1-dimensional factors. In all, if X is reducible then it has a finite dimensional (dimension 1 or 2) G -subspace or G -quotient. By taking duals we can assume we are in the first case. In the Irreducibility Criterion, we want to show that this implies $\chi_1 = \chi_2$ and that X has a 1-dimensional G -subspace.

Definition 4.22. Let π be a smooth representation of G and ϕ a character of F^\times . The twist of π by ϕ is the representation $\phi\pi$ of G defined by

$$\phi\pi(g) = \phi(\det g)\pi(g).$$

In this way, for a character $\chi = \chi_1 \otimes \chi_2$ of T , we have $\phi\chi = \phi\chi_1 \otimes \phi\chi_2$.

Lemma 4.23. For χ a character of T and ϕ a character of F^\times , we have $\text{Ind}_B^G(\phi\chi) = \phi\text{Ind}_B^G\chi$.

Proof. Since $\phi\chi(b) = \phi \circ \det(b)\chi(b)$ for any $b \in B$, where χ is inflated from T , we see that

$$(\phi \circ \det)(bg)f(bg) = \phi\chi(b)(\phi \circ \det)(g)f(g)$$

for any $f \in \text{Ind}_B^G\chi$. Thus the map $f \mapsto (\phi \circ \det)f$ from $\text{Ind}_B^G\chi \rightarrow \text{Ind}_B^G(\phi\chi)$ is well defined on the underlying vector spaces. This induces an isomorphism of representations of G , $\phi\text{Ind}_B^G\chi \cong \text{Ind}_B^G(\phi\chi)$. \square

Proposition 4.24. The following are equivalent:

1. $\chi_1 = \chi_2$
2. X has a 1-dimensional N -subspace.

If this holds then this subspace is also a G -subspace of X not contained in V .

Proof. (1) implies (2): since induction commutes with twisting we may assume $\chi_1 = \chi_2 = 1$. Then any nonzero constant function spans a 1-dimensional G -subspace (not just N -subspace) of $X = \text{Ind}_B^G 1$.

(2) implies (1): suppose this subspace is spanned by f . The group N acts as a character on this subspace via right translation. We cannot have $f \in V$ (meaning $f(1) = 0$) because we saw earlier that f would then have support in some BwN_0 for $N_0 \leq N$ open compact, and this is not closed under multiplication by N .

So $f \notin V$ and therefore its image spans $X/V \cong \mathbb{C}$, on which B acts via χ . On this quotient, N acts trivially because χ was inflated from $B/N = T$. Thus f is in fact fixed by N under right translation. But f is also fixed under right translation by some compact open of G , so for sufficiently large $|x|$ we have

$$\begin{aligned} f(w) &= f\left(w\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}\right) = f\left(\begin{pmatrix} 1 & x^{-1} \\ 0 & 1 \end{pmatrix}\begin{pmatrix} -x^{-1} & 0 \\ 0 & x \end{pmatrix}\begin{pmatrix} 1 & 0 \\ x^{-1} & 1 \end{pmatrix}\right) \\ &= f\left(\begin{pmatrix} 1 & x^{-1} \\ 0 & 1 \end{pmatrix}\begin{pmatrix} -x^{-1} & 0 \\ 0 & x \end{pmatrix}\right) \\ &= \chi_1(-1)(\chi_1^{-1}\chi_2(x))f(1). \end{aligned}$$

The first equality comes from f being fixed by N . The third equality comes from f being fixed by a compact open subgroup of G .

This tells us that $\chi_1^{-1}\chi_2(x)$ is constant for $|x|$ sufficiently large. In particular, for large $|x|$ we have $\chi_1^{-1}\chi_2(x) = \chi_1^{-1}\chi_2(x^2) = (\chi_1^{-1}\chi_2(x))^2$. We deduce that $\chi_1(x) = \chi_2(x)$ for $|x|$ sufficiently large. Now for any $y \in F^\times$, we can pick $|x|$ large enough so that $\chi_1(x) = \chi_2(x)$ and $\chi_1(xy) = \chi_2(xy)$, from which we deduce that $\chi_1(y) = \chi_2(y)$. \square

Proof of Irreducibility Criterion. Assume that X is reducible and we are in the case that X has a finite dimensional G -subspace. It has a 1-dimensional N -subspace L because N is abelian. Then L is also a G -subspace by the above proposition. Since G must act via a character on L , it factors as $\phi \circ \det$, where $\chi_1 = \phi = \chi_2$.

Let Y be the G -representation X/L . Since L spans the vector space X/V , the B -homomorphism $V \hookrightarrow X \rightarrow X/L$ is surjective. It is injective since $L \cap V = 0$. Thus $Y \cong V$ as B -representations.

We need to show that X has G -length 2. By the Corollary 4.20 it has length at most 3. We know that V has B -length 2 with a 1-dimensional quotient V_N . If Y had G -length 2, then the B -factors of V are also G -factors, so that G must act on V_N , necessarily by a character $\phi' \circ \det$. But this is impossible because $B \leq G$ acts on V_N by $\phi\delta_B^{-1}$ by Restriction-Induction, and this does not factor through \det on B . So we must have that Y is irreducible over G and so X has G -length 2.

In the other case we have a finite dimensional G -quotient. The smooth dual X^\vee is then in the first case, where the Duality Theorem ([BH06, Theorem 3.5]) tells us that $X^\vee \cong \text{Ind}_B^G \delta_B^{-1} \chi^\vee$. If we write $\delta_B^{-1} \chi^\vee = \psi_1 \otimes \psi_2$ then we must have $\psi_1 = \psi_2$. Computing $\psi_1(x) = |x|^{-1} \chi_1(x)$ and $\psi_2(x) = |x| \chi_2(x)$ gives $\chi_1 \chi_2^{-1} = |\cdot|^2$.

The converse direction to (1) follows from the previous proposition. \square

4.4 Classification of principal series representations

Now that we've seen how parabolically induced representations decompose into irreducibles, we want to classify the isomorphism classes.

Proposition 4.25. *Let χ, ξ be characters of T . The space $\text{Hom}_G(\text{Ind}_B^G \chi, \text{Ind}_B^G \xi)$ is 1-dimensional if $\xi = \chi$ or $\chi^w \delta_B^{-1}$ and 0 otherwise.*

Proof. Frobenius reciprocity tells us

$$\text{Hom}_G(\text{Ind}_B^G \chi, \text{Ind}_B^G \xi) \cong \text{Hom}_T((\text{Ind} \chi)_N, \xi).$$

From the Restriction-Induction lemma we have the exact sequence of T -modules

$$0 \longrightarrow \chi^w \delta_B^{-1} \longrightarrow (\text{Ind} \chi)_N \longrightarrow \chi \longrightarrow 0.$$

By taking duals of these finite dimensional T -modules, we see that both χ and $\chi^w \delta_B^{-1}$ are subrepresentations of $(\text{Ind} \chi)_N$. In the case $\chi \neq \chi^w \delta_B^{-1}$ we must have $(\text{Ind} \chi)_N = \chi \oplus \chi^w \delta_B^{-1}$ and the result follows. If $\chi = \chi^w \delta_B^{-1}$ then $\chi_1 \chi_2^{-1}(x) = |x|$ so $\text{Ind} \chi$ is irreducible and the result still follows from Schur's lemma. \square

Remark 4.26. In the case that $\text{Ind}\chi$ is irreducible, we deduce that $\text{Ind}\chi \cong \text{Ind}\chi^w \delta_B^{-1}$. And in the case $\text{Ind}\chi$ is reducible, it is not semisimple, else $\text{Hom}_G(\text{Ind}_B^G \chi, \text{Ind}_B^G \chi)$ would have dimension strictly greater than 1.

We can be more explicit in the reducible case. One can check that the conditions for reducibility in the Irreducibility Criterion are equivalent to χ being of the form $\chi = \phi 1_T$ or $\chi = \phi \delta_B^{-1}$ for ϕ a character of F^\times . Untwisting, we may as well assume $\phi = 1$ in what follows.

Definition 4.27. The Steinberg representation of G is defined by the exact sequence

$$0 \longrightarrow 1_G \longrightarrow \text{Ind}_B^G 1_T \longrightarrow \text{St}_G \longrightarrow 0,$$

and is an infinite-dimensional irreducible smooth representation. By Restriction-Induction, the Jacquet module is $(\text{St}_G)_N \cong \delta_B^{-1}$. The representations ϕSt_G are called ‘twists of Steinberg’ or ‘special representations’.

The case $\chi = \delta_B^{-1}$ can be dealt with by taking smooth duals (which is exact by [BH06, Lemma 2.10]) to get

$$0 \longrightarrow \text{St}_G^\vee \longrightarrow \text{Ind}_B^G \delta_B^{-1} \longrightarrow 1_G \longrightarrow 0,$$

where we use the Duality Theorem, [BH06, Theorem 3.5]. The Irreducibility Criterion implies that St_G^\vee is also irreducible, and in fact the previous proposition applied to $\chi = 1, \xi = \delta_B^{-1}$ implies that

$$\text{St}_G \cong \text{St}_G^\vee.$$

Notation 4.28. Define normalised induction by

$$\iota_B^G \sigma = \text{Ind}_B^G (\delta_B^{-1/2} \otimes \sigma).$$

This has the benefit that $(\iota_B^G \sigma)^\vee \cong \iota_B^G \sigma^\vee$ ([BH06, Theorem 3.5]).

Theorem 4.29 (Classification Theorem). *The following are all the isomorphism classes of principal series representations of G :*

- the irreducible induced representations $\iota_B^G \chi$ when $\chi \neq \phi \delta_B^{\pm 1/2}$ for a character ϕ of F^\times .
- the one-dimensional representations $\phi \circ \det$ for ϕ a character of F^\times .
- the twists of Steinberg (special representations) ϕSt_G for ϕ a character of F^\times .

These are all distinct isomorphism classes except in the first case where $\iota_B^G \chi \cong \iota_B^G \chi^w$.

5 Functional equation for GL_2

In the previous section, we classified the principal series representations of $G = \text{GL}_2(F)$ over a nonarchimedean local field F . For characters χ of $\text{GL}_1(F)$, Tate’s thesis [Tat67] associates a space $\mathcal{Z}(\chi)$ of zeta functions in a complex variable s . This space will, in a sense to be made precise, be generated by a single element, the L -function $L(\chi, s)$. The zeta functions will also satisfy a functional equation depending on the ‘local constant’

$\epsilon(\chi, s, \psi)$. Here $\psi : F \rightarrow \mathbb{C}^\times$ is a character whose purpose is to fix a form of Fourier transform on F . These definitions and results in Tate's thesis are intended to mimic the classical theory of L -functions due largely to Hecke, which encompass the Riemann zeta function. The L -function and local constant of a character $\chi : F^\times \rightarrow \mathbb{C}^\times$ will turn out to carry the essential information of χ . In the classical setting see, for example, the converse theorem of Weil reproduced in [Bum97, Theorem 1.5.1].

In the setting of irreducible smooth representations π of G , in particular the principal series representations π , we want to again associate a space $\mathcal{Z}(\pi)$ of zeta functions, an L -function $L(\pi, s)$ and a local constant $\epsilon(\pi, s, \psi)$ determining a functional equation.

We begin this section with a brief review of harmonic and Fourier analysis and the role it plays in representation theory. For more details, see [Bum97, Chapter 3.1]. Following the presentation in [GH24], we define the L -functions and local constants of characters of F^\times . We explain how this theory generalises to irreducible smooth representations π of G , culminating in the Theorems 5.44 and 5.46, which determine the functional equations satisfied by the zeta functions associated to π . Propositions 5.33 and 5.41 prove these in the case where $\pi = \iota_B^G \chi$ is a principal series representation. The case where $\iota_B^G \chi$ is reducible, so that π is only a subquotient, requires slightly more work. The results are summarised in Table 1. Finally, we prove a converse theorem for principal series representations of G .

5.1 Review of harmonic analysis

Take as motivation the representation theory of a finite group H . Every irreducible representation of H appears as a direct summand of the regular representation $\mathbb{C}[H]$, with some multiplicity. For a locally compact topological group \mathbb{G} with Haar measure dg , the correct generalisation of $\mathbb{C}[H]$ is the space $L^2(\mathbb{G})$ of measurable functions $f : \mathbb{G} \rightarrow \mathbb{C}$ for which

$$\int_{\mathbb{G}} |f(g)|^2 dg < \infty.$$

The action of \mathbb{G} is by right translation. If \mathbb{G} is additionally abelian, the group $\hat{\mathbb{G}}$ of (unitary) characters of \mathbb{G} is also a locally compact abelian group, the Pontryagin dual of \mathbb{G} .

Example 5.1. The Pontryagin duals of $\mathbb{G} = \mathbb{R}, \mathbb{Z}, \mathbb{R}/\mathbb{Z}$ are $\mathbb{R}, \mathbb{R}/\mathbb{Z}, \mathbb{Z}$ respectively. The characters of \mathbb{R} are of the form $x \mapsto e^{-2\pi i xy}$ for $y \in \mathbb{R}$. The characters of \mathbb{Z} are of the form $n \mapsto e^{-2\pi i nx}$ for $x \in \mathbb{R}/\mathbb{Z} \cong S^1$. The characters of \mathbb{R}/\mathbb{Z} are of the form $x \mapsto e^{-2\pi i nx}$ for $n \in \mathbb{Z}$. In particular, \mathbb{R} is self-dual.

On a suitable dense subset of $L^2(\mathbb{G})$ (the Schwartz space), one can define the Fourier transform $\hat{f} \in L^2(\hat{\mathbb{G}})$ of f by

$$\hat{f}(\chi) = \int_{\mathbb{G}} f(g) \chi(g) dg.$$

The Fourier transform uniquely extends to a map $L^2(\mathbb{G}) \rightarrow L^2(\hat{\mathbb{G}})$. For suitable choices of Haar measures there is then a Fourier inversion formula

$$\hat{\hat{f}}(g) = f(-g),$$

so that the above map is a bijection.

Example 5.2. For $\mathbb{G} = \mathbb{R}$, the Fourier transform of f is

$$\hat{f}(x) = \int_{\mathbb{R}} f(y) e^{-2\pi i x y} dy$$

which is the classical Fourier transform. Identifying $\hat{\mathbb{R}} = \mathbb{R}$, the Fourier transform gives an invertible map $L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$, so that any element of $L^2(\mathbb{R})$ can be expressed as an integral of elements of $\hat{\mathbb{R}}$.

Inside the representation $L^2(\mathbb{R})$ of \mathbb{R} we therefore see this ‘continuous spectrum’ of the irreducible unitary representations (characters) of \mathbb{R} , parametrised by \mathbb{R} . Note, however, that each such character can not be realised as a subrepresentation of $L^2(\mathbb{R})$; for $y \in \mathbb{R}$ the character $x \mapsto e^{-2\pi i x y}$ is realised as the Fourier transform of a function on \mathbb{R} supported only at y , but such a function is not in $L^2(\mathbb{R})$.

Example 5.3. For $\mathbb{G} = \mathbb{Z}$, the Fourier transform of f is

$$\hat{f}(x) = \sum_{\mathbb{Z}} f(n) e^{-2\pi i n x}.$$

So any element of $L^2(\mathbb{R}/\mathbb{Z})$ can be expressed as a sum of unitary characters of \mathbb{Z} ; we have a ‘discrete spectrum’.

Remark 5.4. The terminology of discrete and continuous spectra comes from the analogy with the spectral theory of the Laplacian. Over \mathbb{R} , the Laplacian is $\Delta = \frac{\partial^2}{\partial x^2}$, and the characters $x \mapsto e^{-2\pi i x y}$ are eigenfunctions.

The dichotomy in the above examples is reflected in the compactness of S^1 and non compactness of \mathbb{R} . More generally,

Theorem 5.5 (Peter-Weyl). *Let K be a compact Hausdorff topological group. Any unitary representation of K decomposes into a completed Hilbert space direct sum of irreducible unitary subrepresentations. There is a unitary equivalence*

$$L^2(K) \cong \widehat{\bigoplus_{\pi \in \hat{K}} \text{End}(V_{\pi})}$$

of representations of $K \times K$, where (π, V_{π}) ranges over the set \hat{K} of equivalence classes of irreducible representations of K , and $\hat{\oplus}$ denotes the completed Hilbert space direct sum.

Proof. [DE09, Theorem 7.3.2] and [DE09, Theorem 7.2.3]. □

Even more generally, for so-called Type I groups one can decompose unitary representations through a combination of integrals and Hilbert space direct sums. See [GH24, Section 3.10] for further details.

Returning to $G = \text{GL}_2(F)$, as this is not compact we would expect the regular representation $L^2(G)$ to decompose according to both a continuous spectra and a discrete spectra. This continuous spectra is provided by the parabolically induced representations $\iota_B^G \chi$, where χ ranges over the characters of $T \cong F^{\times} \times F^{\times}$.

In order to compare representations of G and Galois representations through the local Langlands correspondence, we would like to classify them according to some common language. This is provided by the zeta functions, L -functions and functional equations discussed in this section.

The prototypical example of an L -function is the Riemann zeta function $\zeta(s) = \sum_{n \geq 1} n^{-s}$.

Proposition 5.6. *The function $\zeta(s) = \sum_{n \geq 1} n^{-s}$ satisfies the following properties:*

- (Analytic continuation) The Riemann zeta function converges absolutely to a holomorphic function on $\text{Re}(s) > 1$. It has a unique analytic continuation to the complex plane, except the point $s = 1$ where $\zeta(s)$ has a simple pole.
- (Euler product) We have the identity

$$\sum_{n=1}^{\infty} n^{-s} = \prod_{p \text{ prime}} \frac{1}{1 - p^{-s}},$$

convergent for $\text{Re}(s) > 1$.

- (Functional equation) There is an explicit function $\gamma(s)$ such that $\zeta(1 - s) = \gamma(s)\zeta(s)$.

The approach of Tate in his thesis was to view the Riemann (And Dedekind) zeta functions from an adelic perspective. There the Euler product formulation is immediate and we only need to study the zeta functions locally. Attached to any character $\chi : F^\times \rightarrow \mathbb{C}^\times$ there is an associated space $\mathcal{Z}(\chi)$ of zeta functions $\zeta(\Phi, \chi, s)$, where $\Phi \in C_c^\infty(F)$. The factor at the prime p of the Riemann zeta function corresponds to the trivial character of \mathbb{Q}_p^\times and the function $\mathbb{1}_{\mathbb{Z}_p} \in C_c^\infty(\mathbb{Q}_p)$. A key ingredient in the proof of the functional equation of the Riemann zeta function is the Fourier transform over \mathbb{C} . In general, the functional equation associated to χ relates zeta functions $\zeta(\hat{\Phi}, \chi^{-1}, 1 - s)$ and $\zeta(\Phi, \chi, s)$, where $\hat{\Phi}$ is the Fourier transform of Φ in $C_c^\infty(F)$.

5.2 Functional equation for GL_1

Let F be a nonarchimedean local field, ϖ be a uniformiser and q be the size of the residue field. We will later define L -functions attached to an irreducible smooth representation of $\text{GL}_2(F)$ and determine a functional equation they satisfy. First we explain this in the context of irreducible smooth representations χ of $\text{GL}_1(F)$, necessarily a character $\chi : F^\times \rightarrow \mathbb{C}^\times$.

Taking from the classical study of the Riemann zeta function and its functional equation, we want to introduce an analogue of the Fourier transform over F . We replace the additive character $e^{2\pi i \cdot} : \mathbb{R} \rightarrow \mathbb{C}^\times$ with any choice of additive character $\psi : F \rightarrow \mathbb{C}^\times$ with $\psi \neq 1$. In this way, all characters of F are of the form $\psi(-y)$ for $y \in F$, by Additive Duality. The functions we will apply the Fourier transform to will be the algebra $C_c^\infty(F)$ of locally constant compactly supported functions $F \rightarrow \mathbb{C}$. For any choice of Haar measure μ on F , we now define the Fourier transform.

Definition 5.7. Let $\Phi \in C_c^\infty(F)$, $\psi : F \rightarrow \mathbb{C}^\times$ be an additive character of F , and μ be a Haar measure on F . The Fourier transform of Φ (with respect to ψ and μ) is

$$\hat{\Phi}(x) := \int_F \Phi(y) \psi(xy) d\mu(y).$$

To match the classical definition over \mathbb{R} , we would like the Fourier transform to preserve $C_c^\infty(F)$, and to have a Fourier inversion formula. Indeed:

Proposition 5.8. • For any $\Phi \in C_c^\infty(F)$, we have $\hat{\hat{\Phi}} \in C_c^\infty(F)$.

- For any $\psi : F \rightarrow \mathbb{C}^\times$ with $\psi \neq 1$, there is a unique Haar measure μ_ψ on F such that for the associated Fourier transform we have

$$\hat{\Phi}(x) = \Phi(-x)$$

for any $\Phi \in C_c^\infty(F)$ and $x \in F$.

Proof. [BH06, Proposition 23.1] □

Notation 5.9. For the remainder of this subsection, $\psi \neq 1$ will be an additive character of F , and $\mu = \mu_\psi$ will denote the associated self-dual Haar measure on F .

Now let $\chi : F^\times \rightarrow \mathbb{C}^\times$ be a smooth character of F^\times . We want to attach to this character an L -function $L(\chi, s)$ in the formal variable s . This is defined to be $(1 - \chi(\varpi)q^{-s})^{-1}$ when χ is unramified, and 1 otherwise. In order to generalise to GL_2 it would be preferable to have a more intrinsic definition.

Definition 5.10. For $\Phi \in C_c^\infty(F)$ and $\chi : F^\times \rightarrow \mathbb{C}^\times$, define the zeta function $\zeta(\Phi, \chi, s)$ to be

$$\zeta(\Phi, \chi, s) := \int_{F^\times} \Phi(x) \chi(x) |x|^s d^*x,$$

in the formal variable s , where $d\mu^*(x) = d^*x$ denotes any choice of Haar measure on F^\times .

Equivalently, we have

$$\zeta(\Phi, \chi, s) = \sum_{m \in \mathbb{Z}} z_m q^{-ms}$$

for

$$z_m = \int_{\varpi^m \mathcal{O}_F^\times} \Phi(x) \chi(x) d^*x.$$

In this way it is clear that $\zeta(\Phi, \chi, s) \in \mathbb{C}((q^{-s}))$. The $z_m = z_m(\Phi, \chi)$ vanish for $m < 0$ because Φ is compactly supported on F .

The zeta function $\zeta(\Phi, \chi, s)$ only depends on d^*x up to scaling. To remove this dependence we define the following notation.

Notation 5.11. Let

$$\mathcal{Z}(\chi) = \{\zeta(\Phi, \chi, s) \mid \Phi \in C_c^\infty(F)\}.$$

Notation 5.12. For $a \in F^\times$ and $\Phi \in C_c^\infty(F)$, denote by $a\Phi$ the function $x \mapsto \Phi(a^{-1}x)$.

Lemma 5.13. The space $\mathcal{Z}(\chi)$ is a $\mathbb{C}[q^{-s}, q^s]$ -module, containing $\mathbb{C}[q^{-s}, q^s]$.

Proof. Let $a \in F^\times$ of valuation $v_F(a)$. Then

$$\zeta(a\Phi, \chi, s) = \chi(a) q^{-v_F(a)s} \zeta(\Phi, \chi, s),$$

giving the desired module structure. To establish the containment we show that $\mathcal{Z}(\chi)$ contains a nonzero constant. Let d be such that $\chi|_{U_F^{d+1}} = 1$. Taking $\Phi = \mathbb{1}_{U_F^{d+1}}$, we see that

$$\zeta(\Phi, \chi, s) = \mu^*(U_F^{d+1}) \neq 0.$$

□

Proposition 5.14. *Let $\chi : F^\times \rightarrow \mathbb{C}^\times$. There exists a unique polynomial $P_\chi \in \mathbb{C}[X]$ with $P_\chi(0) = 1$ such that*

$$\mathcal{Z}(\chi) = P_\chi(q^{-s})^{-1} \cdot \mathbb{C}[q^{-s}, q^s].$$

Moreover, we have

$$P_\chi(X) = \begin{cases} 1 - \chi(\varpi)X & \text{if } \chi \text{ is unramified} \\ 1 & \text{otherwise} \end{cases}$$

Proof. Suppose $\Phi(0) = 0$. Then $\Phi|_{F^\times} \in C_c^\infty(F^\times)$, and so Φ is identically zero on $\varpi^m \mathcal{O}_F^\times$ for $|m| \gg 0$. Thus only finitely many of the coefficients z_m are nonzero, so that $\Phi \in \mathbb{C}[q^{-s}, q^s]$.

The space $C_c^\infty(F)$ is spanned by $C_c^\infty(F^\times)$ and $\mathbb{1}_{\mathcal{O}_F}$. We compute

$$\zeta(\mathbb{1}_{\mathcal{O}_F}, \chi, s) = \sum_{m \geq 0} \chi(\varpi^m) q^{-ms} \int_{\mathcal{O}_F^\times} \chi(x) d^*x.$$

If χ is unramified (trivial on \mathcal{O}_F^\times), this gives us

$$\sum_{m \geq 0} \chi(\varpi)^m q^{-ms} \mu^*(\mathcal{O}_F^\times) = (1 - \chi(\varpi) q^{-s})^{-1} \mu^*(\mathcal{O}_F^\times).$$

When χ is ramified the integral is zero. Indeed, translation invariance of d^*x implies

$$\int_{\mathcal{O}_F^\times} \chi(x) d^*x = \int_{\mathcal{O}_F^\times} \chi(xy) d^*x = \chi(y) \int_{\mathcal{O}_F^\times} \chi(x) d^*x$$

for any $y \in \mathcal{O}_F^\times$, so that this is zero if there is some y with $\chi(y) \neq 1$. This computation, together with the previous lemma, establish the result. \square

Remark 5.15. The computation in the proof above shows, in the case $\chi = 1$, that $\zeta(\mathbb{1}_{\mathcal{O}_F}, 1, s) = (1 - q^{-s})^{-1}$, provided we normalise d^*x appropriately. If $F = K_v$ is the completion of a number field K at a nonarchimedean place v , we recover the Euler factor of the Dedekind zeta function $\zeta_K(s)$ at the place v . This explains the naming of our zeta functions.

Remark 5.16. The computations of Proposition 5.14 show that each $\zeta(\Phi, \chi, s)$ converges absolutely and uniformly in vertical strips in some right half plane, and admit analytic continuation to a rational function in q^{-s} .

Definition 5.17. Define the L -function attached to χ to be $L(\chi, s) = P_\chi(q^{-s})^{-1}$.

As with the Riemann zeta function, we have functional equations for the zeta functions.

Theorem 5.18. *Let $\chi : F^\times \rightarrow \mathbb{C}^\times$. There is a unique $\gamma(\chi, s, \psi) \in \mathbb{C}(q^{-s})$ such that*

$$\zeta(\hat{\Phi}, \check{\chi}, 1 - s) = \gamma(\chi, s, \psi) \zeta(\Phi, \chi, s)$$

for all $\Phi \in C_c^\infty(F)$, where $\check{\chi} = 1/\chi : F^\times \rightarrow \mathbb{C}^\times$.

Proof. [BH06, Theorem 23.3]. \square

Since $\mathcal{Z}(\chi) = L(\chi, s) \cdot \mathbb{C}[q^{-s}, q^s]$, it is natural to consider the terms $\frac{\zeta(\Phi, \chi, s)}{L(\chi, s)} \in \mathbb{C}[q^{-s}, q^s]$. This allows us to treat the case of χ ramified and unramified evenly.

Definition 5.19. Let

$$\epsilon(\chi, s, \psi) := \gamma(\chi, s, \psi) \cdot \frac{L(\chi, s)}{L(\tilde{\chi}, s)}.$$

This is known as Tate's local constant.

The functional equation for ζ can be rewritten as

$$\frac{\zeta(\hat{\Phi}, \tilde{\chi}, 1-s)}{L(\tilde{\chi}, 1-s)} = \epsilon(\chi, s, \psi) \frac{\zeta(\Phi, \chi, s)}{L(\chi, s)}.$$

Corollary 5.20. *The local constant satisfies the functional equation*

$$\epsilon(\chi, s, \psi) \epsilon(\tilde{\chi}, 1-s, \psi) = \chi(-1).$$

The local constant is of the form

$$\epsilon(\chi, s, \psi) = a q^{bs}$$

for some $a \in \mathbb{C}^\times$, $b \in \mathbb{Z}$.

Proof. The first statement comes from the Fourier inversion formula, where the $\chi(-1)$ term comes from the minus sign in $\hat{\Phi}(x) = \Phi(-x)$. The functional equation implies that ϵ is a unit in $\mathbb{C}[q^{-s}, q^s]$, and the units are precisely the elements of the form $a q^{bs}$ for $b \in \mathbb{Z}$. \square

5.3 Functional equation for GL_2

We turn now to smooth representations π of $G = \text{GL}_2(F)$ and define the L -functions and local constants in an analogous manner to the characters $\chi : F^\times \rightarrow \mathbb{C}^\times$.

In this context, we need an additive character of $A = M_2(F)$, which we will take to be $\psi_A = \psi \circ \text{tr}$ for $\psi : F \rightarrow \mathbb{C}^\times$ any nontrivial additive character of F . We will apply the Fourier transform to the F -algebra $\Phi \in C_c^\infty(A)$ of locally constant compactly supported functions on $M_2(F)$.

Definition 5.21. With respect to a Haar measure μ in A , and $\psi_A = \psi \circ \text{tr}$ an additive character of A , define for any $\Phi \in C_c^\infty(A)$

$$\hat{\Phi}(x) = \int_A \Phi(y) \psi_A(xy) d\mu(y).$$

Proposition 5.22. • For any $\Phi \in C_c^\infty(A)$, we have $\hat{\Phi} \in C_c^\infty(A)$.

- For any $\psi : F \rightarrow \mathbb{C}^\times$ with $\psi \neq 1$, there is a unique Haar measure μ_{ψ_A} on A such that for the associated Fourier transform we have

$$\hat{\hat{\Phi}}(x) = \Phi(-x)$$

for any $\Phi \in C_c^\infty(A)$ and $x \in A$.

Notation 5.23. For the remainder of this subsection, $\psi \neq 1$ will be an additive character of F , $\psi_A = \psi \circ \text{tr}$, and $\mu = \mu_{\psi_A}$ will denote the associated self-dual Haar measure on A .

For $\chi : F^\times \rightarrow \mathbb{C}^\times$ we defined for any $\Phi \in C_c^\infty(F)$ a zeta function

$$\zeta(\Phi, \chi, s) = \int_{F^\times} \Phi(x) \chi(x) |x|^s d^*x.$$

To replicate this with $\pi : G \rightarrow \mathrm{GL}(V)$ we need to extract scalar values from $\pi(g) \in \mathrm{GL}(V)$. These will come from matrix coefficients.

Definition 5.24. Let (π, V) be a smooth representation of G with smooth dual \check{V} . For vectors $v \in V, \check{v} \in \check{V}$, define the smooth function $\gamma_{v \otimes \check{v}} : G \rightarrow \mathbb{C}$ by

$$\gamma_{v \otimes \check{v}} : g \mapsto \langle \check{v}, \pi(g)v \rangle$$

where \langle, \rangle denotes the natural pairing $\check{V} \otimes V \rightarrow \mathbb{C}$. Let $\mathcal{C}(\pi)$ be the vector space spanned by the $\gamma_{v \otimes \check{v}}$. Elements of $\mathcal{C}(\pi)$ are called the matrix coefficients of π .

Remark 5.25. If $\pi = \chi : F^\times \rightarrow \mathbb{C}^\times$ is a character, any matrix coefficient (defined in the analogous way for F^\times) of χ is some scalar multiple of χ .

If V is the tautological representation of G with basis e_1, e_2 , then $\gamma_{e_i \otimes e_j}(g)$ is precisely the (i, j) -th entry of g as a matrix with respect to the basis e_1, e_2 .

Definition 5.26. Let (π, V) be an irreducible smooth representation of G . The centre Z of G acts on V via the central character $\omega_\pi : Z \rightarrow \mathbb{C}^\times$.

Lemma 5.27. For any $f \in \mathcal{C}(\pi), z \in Z, g \in G$ we have $f(zg) = \omega_\pi(z)f(g)$.

Fix a smooth representation π of G . We may now define zeta functions for any $f \in \mathcal{C}(\pi)$.

Definition 5.28. For $\Phi \in C_c^\infty(A)$ and $f \in \mathcal{C}(\pi)$, define the zeta function $\zeta(\Phi, f, s)$ to be

$$\zeta(\Phi, f, s) := \int_G \Phi(x) f(x) |\det x|^s d^*x,$$

in the formal variable s , where $d\mu^*(x) = d^*x$ denotes any choice of Haar measure on G .

Lemma 5.29. For any $\Phi \in C_c^\infty(A)$ and $f \in \mathcal{C}(\pi)$ we have $\zeta(\Phi, f, s) \in \mathbb{C}((q^{-s}))$ in the formal variable s .

Proof. This follows from [BH06, Lemma 24.4.1]. □

Notation 5.30. Let

$$\mathcal{Z}(\pi) = \{\zeta(\Phi, f, s + \frac{1}{2}) \mid \Phi \in C_c^\infty(A), f \in \mathcal{C}(\pi)\}.$$

Remark 5.31. The addition of $\frac{1}{2}$ will be explained in the case of principal series representations by the appearance of the modular character δ_B .

Lemma 5.32. The space $\mathcal{Z}(\pi)$ is a $\mathbb{C}[q^{-s}, q^s]$ -module, containing $\mathbb{C}[q^{-s}, q^s]$.

Proof. [BH06, Lemma 24.4.2]. □

Consider now the situation where $\pi = \iota_B^G \chi$ is a parabolically induced representation, where $\chi = \chi_1 \otimes \chi_2$ is a character of T . We want to study the space $\mathcal{Z}(\pi)$ and prove an analogous result to Proposition 5.14.

Proposition 5.33. *Let $\chi = \chi_1 \otimes \chi_2$ be a character of T and let $(\pi, V) = \iota_B^G \chi$. Then, formally, we have*

$$\mathcal{Z}(\pi) = \mathcal{Z}(\chi_1)\mathcal{Z}(\chi_2) \subset \mathbb{C}((q^{-s})).$$

In particular, there is a unique polynomial $P_\pi \in \mathbb{C}[X]$ with $P_\pi(0) = 1$ such that

$$\mathcal{Z}(\pi) = P_\pi(q^{-s})^{-1} \cdot \mathbb{C}[q^{-s}, q^s].$$

Moreover, $P_\pi(X) = P_{\chi_1}(X)P_{\chi_2}(X)$.

We make some comments in preparation for the proof. The Proposition concerns the zeta integrals

$$\zeta(\Phi, f, s + \frac{1}{2}) = \int_G \Phi(x) f(x) |\det x|^{s + \frac{1}{2}} d^* x.$$

The matrix coefficients $\mathcal{C}(\pi)$ are spanned by

$$\gamma_{\tau \otimes \theta} : g \mapsto \langle \tau, \pi(g)\theta \rangle$$

over $\theta \in V, \tau \in \check{V}$. Here $\theta \in \iota_B^G \chi$ is viewed as a smooth function $\theta : G \rightarrow \mathbb{C}$ satisfying

$$\theta(ntg) = \delta_B^{-1/2}(t)\chi(t)\theta(g)$$

for any $t \in T, n \in N, g \in G$. The Duality Theorem [ADD REFERENCE] identifies $\check{V} \cong \iota_B^G \check{\chi}$. In this way we view τ as a smooth function $\tau : G \rightarrow \mathbb{C}$ satisfying

$$\tau(ntg) = \delta_B^{-1/2}(t)\chi(t)^{-1}\tau(g)$$

for any $t \in T, n \in N, g \in G$. The proof of the Duality Theorem shows that the pairing between V and \check{V} gives

$$f(g) = \langle \tau, \pi(g)\theta \rangle = \int_{B \backslash G} \tau(x)\theta(xg) d\dot{x}$$

for a positive semi-invariant measure $d\dot{x}$ on $B \backslash G$. Let $K = \mathrm{GL}_2(\mathcal{O}_F)$. Since we have a bijection $B \backslash G \leftrightarrow K \cap B \backslash K$ and $\delta_B(tn) = \delta_B(t) = |t_2/t_1|$ (Proposition 3.15) is trivial on $K \cap B$, we can rewrite this as

$$f(g) = \int_K \tau(k)\theta(kg) dk$$

for some Haar measure dk on K ([BH06, Corollary 7.6]). Moreover, [BH06, Equation 7.6.2] tells us that there is a left Haar measure db on B such that

$$\int_G \phi(g) dg = \int_K \int_B \phi(bk) db dk$$

for all $\phi \in C_c^\infty(G)$. Using this, our zeta integrals reduce to integrals over B and K . Integration over K is easier to handle using the smoothness of our representations. We can write $db = dn dt$ to view integration over B as integration over T and N . In order to relate $\zeta(\Phi, f, s + \frac{1}{2})$ to zeta functions coming from $\chi : T \rightarrow \mathbb{C}^\times$, we want to express the integrals over B solely in terms of integrals over T . To do so we use the following lemma.

Lemma 5.34. *Let D be the algebra of diagonal matrices in A so that $D^\times = T$. Let $\Phi \in C_c^\infty(A)$. There is a unique function $\Phi_T \in C_c^\infty(D)$ whose restriction to T is given by*

$$\Phi_T(t) = |t_1| \int_N \Phi(tn) dn, \quad t = \begin{pmatrix} t_1 & 0 \\ 0 & t_2 \end{pmatrix}.$$

The map $\Phi \mapsto \Phi_T$ is a linear surjection $C_c^\infty(A) \rightarrow C_c^\infty(D)$.

Proof. The space $C_c^\infty(A)$ is spanned by functions of the form

$$\Phi = (\phi_{ij}) : (a_{ij}) \mapsto \prod_{i,j} \phi_{ij}(a_{ij})$$

for $\phi_{ij} \in C_c^\infty(F)$. For such Φ we compute (identifying $N \cong F$)

$$\begin{aligned} \Phi_T(t) &= |t_1| \int_F \phi_{11}(t_1) \phi_{12}(t_1 n) \phi_{21}(0) \phi_{22}(t_2) dn \\ &= \phi_{11}(t_1) \phi_{22}(t_2) \phi_{21}(0) |t_1| \int_F \phi_{12}(t_1 n) dn \\ &= \phi_{11}(t_1) \phi_{22}(t_2) \phi_{21}(0) \int_F \phi_{12}(n) dn \end{aligned}$$

which uniquely extends to a function in $C_c^\infty(D)$. Surjectivity is now clear. \square

Remark 5.35. The content of the lemma is that the function Φ_T is smooth, for which the introduction of the factor of $|t_1|$ is necessary.

Proof of Proposition 5.33. We first establish the containment $\mathcal{Z}(\pi) \subset \mathcal{Z}(\chi_1)\mathcal{Z}(\chi_2)$. We must show that for any $\Phi \in C_c^\infty(A)$ and $f \in \mathcal{C}(\pi)$ we have $\zeta(\Phi, f, s + \frac{1}{2}) \in \mathcal{Z}(\chi_2)\mathcal{Z}(\chi_2)$. Since $\mathcal{C}(\pi)$ is spanned by the coefficients $\gamma_{\tau \otimes \theta}$, for $\theta \in V, \tau \in \check{V}$, we assume f is of this form.

Formally expanding, for any $\Phi \in C_c^\infty(A)$

$$\begin{aligned} \zeta(\Phi, f, s + \frac{1}{2}) &= \int_G \Phi(g) f(g) |\det g|^{s+\frac{1}{2}} dg \\ &= \int_G \int_K \Phi(g) \tau(k) \theta(kg) |\det g|^{s+\frac{1}{2}} dk dg \\ &= \int_K \int_G \Phi(k^{-1}g) \tau(k) \theta(g) |\det g|^{s+\frac{1}{2}} dg dk \\ &= \int_K \int_K \int_B \Phi(k^{-1}bk') \tau(k) \theta(bk') |\det b|^{s+\frac{1}{2}} db dk' dk. \end{aligned}$$

Smoothness of Φ and θ imply there is some open normal subgroup K_1 of K for which Φ is left and right translation invariant, and θ and τ are right translation invariant. Let $\{k_i\}$ be a finite set of coset representatives of K/K_1 , and let $\Phi^{ij}(x) = \Phi(k_i^{-1}xk_j)$. Then $\zeta(\Phi, f, s + \frac{1}{2})$ can be expressed as a finite linear combination over \mathbb{C} of terms of the form

$$\int_B \Phi^{ij}(b) \tau(k_i) \theta(bk_j) |\det b|^{s+\frac{1}{2}} db.$$

Using the formula $\theta(bk_j) = \delta_B^{-1/2}(t) \chi(t) \theta(k_j)$, we can express the above as

$$\theta(k_j) \tau(k_i) \int_T \int_N \Phi^{ij}(tn) \chi(t) \delta_B^{-1/2}(t) |\det b|^{s+\frac{1}{2}} dt dn.$$

We have $|\det b| = |\det t| = |t_1||t_2|$ and $\delta_B^{-1/2}(t) = |t_2/t_1|^{-1/2}$. Combining with the previous lemma, we deduce that $\zeta(\Phi, f, s + \frac{1}{2})$ can be expressed as a linear combination of terms of the form

$$\theta(k_j)\tau(k_i) \int_T \Phi_T^{ij}(t)\chi(t)|\det t|^s dt.$$

If Φ is of the form (ϕ_{ij}) for $\phi_{ij} \in C_c^\infty(F)$, then the above term is a scalar multiple of $\zeta(\phi_{11}, \chi_1, s)\zeta(\phi_{22}, \chi_2, s)$ so that $\zeta(\Phi, f, s + \frac{1}{2}) \in \mathcal{Z}(\chi_1)\mathcal{Z}(\chi_2)$.

In the other direction, we wish to find $\Phi \in C_c^\infty(A)$ and $f \in \mathcal{C}(\pi)$ such that $\zeta(\Phi, f, s + \frac{1}{2})$ is a constant multiple of $L(\chi_1, s)L(\chi_2, s)$. We will find f of the form $\gamma_{\tau \otimes \theta}$ and reverse the above calculation. Suppose we were in the situation where Φ is left and right invariant under K , and θ and τ are right invariant under K . Then the above computation shows that

$$\zeta(\Phi, f, s + \frac{1}{2}) = \mu(K)^2 \theta(1)\tau(1) \int_T \Phi_T(t)\chi(t)|\det t|^s dt.$$

Therefore, if we could choose Φ left and right invariant under K with $\Phi_T = \phi_1 \otimes \phi_2$, where $\phi_i \in C_c^\infty(F)$ satisfy $\zeta(\phi_i, \chi_i, s) = L(\chi_i, s)$, and also choose $\theta \in \iota_B^G \chi$, $\tau \in \iota_B^G \check{\chi}$, with $\theta(1), \tau(1) \neq 0$ and θ, τ right invariant under K , then we would be done. Unfortunately, if this was the case then

$$\theta(bk) = \chi(b)\delta_B^{-1/2}(b)\theta(1)$$

for all $b \in B, k \in K$. But this is not well defined - we would require $1 = \chi(b)\delta_B^{-1/2}(b) = \chi(b)$ for all $b \in B \cap K$. This only occurs when χ_1 and χ_2 are both unramified.

Instead, let K_1 be any open normal subgroup of K such that χ is trivial on $B \cap K_1$, and let k_i be a finite set of coset representatives of K/K_1 . There are then unique $\theta \in \iota_B^G \chi$ and $\tau \in \iota_B^G \check{\chi}$, each supported on BK_1 , invariant under right translation by K_1 , and with $\theta(1) = 1 = \tau(1)$. Let $f = \gamma_{\tau \otimes \theta}$.

For $\Phi \in C_c^\infty(A)$ left and right invariant under K_1 , our previous computation gives us

$$\zeta(\Phi, f, s + \frac{1}{2}) = \mu(K_1)^2 \sum_{i,j} \int_T \theta(k_j)\tau(k_i)\Phi_T^{ij}(t)\chi(t)|\det t|^s dt.$$

To control the terms over all i, j , we would like to choose Φ such that

$$\theta(k_j)\tau(k_i)\Phi_T^{ij}(t) = \Phi_T(t)$$

for all $t \in T$ and all i, j such that $k_i, k_j \in BK_1$. Then, by construction of θ and τ , each term $\theta(k_j)\tau(k_i)\Phi_T^{ij}(t)$ is either 0 or $\Phi_T(t)$, and at least one is $\Phi_T(t)$, so that

$$\zeta(\Phi, f, s + \frac{1}{2}) = c \int_T \Phi_T(t)\chi(t)|\det t|^s dt$$

for some $c > 0$. If $k_j = b_j k \in BK_1$, then $\theta(k_j) = \chi(b_j)\delta_B^{-1/2}(b_j)\theta(1) = \chi(b_j)$ because $\delta_B = 1$ on $B \cap K$. Similarly, if $k_i = b_i k \in BK_1$, then $\tau(k_i) = \chi(b_i)^{-1}$. The condition

$$\theta(k_j)\tau(k_i)\Phi_T^{ij}(t) = \Phi_T(t),$$

together with the K_1 invariance of Φ , reduces to the condition

$$\chi(b_j)\chi(b_i)^{-1} \int_N \Phi(b_i^{-1}tnb_j)dn = \int_N \Phi(tn)dn$$

for all $b_i, b_j \in B \cap K_1$, as functions of $t \in T$.

To summarise, we want to construct $\Phi \in C_c^\infty(A)$ with the following properties:

- The function Φ is invariant under left and right translation by K_1 .
- For all $b_i, b_j \in B \cap K_1$ and $b \in B$ we have

$$\chi(b_j)\chi(b_i)^{-1}\Phi(b_i^{-1}bb_j) = \Phi(b).$$

- For our chosen $\phi_1, \phi_2 \in C_c^\infty(F)$ satisfying $\zeta(\phi_i, \chi_i, s) = L(\chi_i, s)$, we have $\Phi_T = c \cdot \phi_1 \otimes \phi_2 \in C_c^\infty(D)$ for some $c \neq 0$.

Since we may have chosen any open $K_1 \triangleleft K$, provided χ is trivial on $B \cap K_1$, we are free to shrink K_1 and adjust τ and θ accordingly. We can remove the dependence on K_1 by strengthening the second condition above, and now ask for $\Phi \in C_c^\infty(A)$ with the following properties:

- For all $x, y \in B \cap K$ and $b \in B$ we have

$$\chi(xy)\Phi(xby) = \Phi(b).$$

- For some $\phi_1, \phi_2 \in C_c^\infty(F)$ satisfying $\zeta(\phi_i, \chi_i, s) = L(\chi_i, s)$, we have $\Phi_T = c \cdot \phi_1 \otimes \phi_2 \in C_c^\infty(D)$ for some $c \neq 0$.

If we take Φ of the form $\Phi = (\phi_{ij})$, and set $\phi_{12} = \phi_{21} = \mathbb{1}_K$, then the computation of Lemma 5.34 shows that for $t = \begin{pmatrix} t_1 & 0 \\ 0 & t_2 \end{pmatrix}$,

$$\Phi_T(t) = \mu(\mathcal{O}_F)\phi_{11}(t_1)\phi_{22}(t_2).$$

Taking $\phi_{ii} = \phi_i$, it suffices to find for each $i = 1, 2$ some $\phi_i \in C_c^\infty(F)$ such that

- For all $x, y \in \mathcal{O}_F^\times$ and $a \in F^\times$ we have

$$\chi_i(xy)\phi_i(xay) = \phi_i(a).$$

- We have $\zeta(\phi_i, \chi_i, s) = c \cdot L(\chi_i, s)$ for some $c \neq 0$.

Here we divide into cases. If χ_i is unramified, then we may take $\phi_i = \mathbb{1}_{\mathcal{O}_F}$ by the proof of Proposition 5.14. If χ_i is ramified, and the restriction to U_F^n is trivial, then we take

$$\phi_i = \sum_{u \in \mathcal{O}_F^\times / U_F^n} \chi_i(u)^{-1} \mathbb{1}_{uU_F^n}.$$

One sees that this satisfies the first condition. For the second we have

$$\zeta(\phi_i, \chi_i, s) = \sum_u \int_{U_F^n} \chi_i(u)^{-1} \chi_i(ux) |x|^s d^*x = \mu(\mathcal{O}_F^\times)$$

which is a constant (and $L(\chi_i, s) = 1$ in the ramified case). We have proven $\mathcal{Z}(\chi_1)\mathcal{Z}(\chi_2) \subset \mathcal{Z}(\pi)$.

□

Remark 5.36. The computations of Proposition 5.33 show that each $\zeta(\Phi, f, s)$ converges absolutely and uniformly in vertical strips in some right half plane, and admit analytic continuation to a rational function in q^{-s} .

Definition 5.37. Define the L -function attached to $\pi = \iota_B^G \chi$, where $\chi = \chi_1 \otimes \chi_2$ is a character of T , to be

$$L(\pi, s) = P_\pi(q^{-s})^{-1} = L(\chi_1, s)L(\chi_2, s).$$

We now turn to the functional equations satisfied by the zeta functions $\zeta(\Phi, f, s)$. This involves understanding these zeta functions when we replace Φ with its Fourier transform, $\hat{\Phi}$. From the computations of Proposition 5.33, this boils down to relating the map $\Phi \mapsto \Phi_T$ to the various Fourier transforms over A and D .

Lemma 5.38. For $\Phi \in C_c^\infty(A)$, we have $(\hat{\Phi})_T = \widehat{\Phi_T}$.

Proof. [BH06, Lemma 26.3]. □

Lemma 5.39. For $k_i, k_j \in K$ let Φ^{ij} denote the function $x \mapsto \Phi(k_i^{-1} x k_j)$ for $\Phi \in C_c^\infty(A)$. Then $\hat{\Phi}^{ji} = \widehat{\Phi^{ij}}$.

Proof. We calculate

$$\hat{\Phi}^{ji}(x) = \int_A \Phi(y) \psi_A(k_j^{-1} x k_i y) dy$$

and

$$\widehat{\Phi^{ij}}(x) = \int_A \Phi(k_i^{-1} y k_j) \psi_A(xy) dy = \int_A \Phi(y) \psi_A(x k_i y k_j^{-1}) dy.$$

Since $\psi_A = \psi \circ \text{tr}$ and tr is invariant under conjugation, we have $\psi_A(k_j^{-1} x k_i y) = \psi_A(x k_i y k_j^{-1})$. □

Notation 5.40. If $f \in \mathcal{C}(\pi)$ is a matrix coefficient, denote by $\check{f} \in \mathcal{C}(\check{\pi})$ the matrix coefficient $\check{f}(g) = f(g^{-1})$.

Proposition 5.41. Let $\pi = \iota_B^G \chi$ where $\chi = \chi_1 \otimes \chi_2$ is a character of T . There is a unique $\gamma(\pi, s, \psi) \in \mathbb{C}(q^{-s})$, depending on the additive character $\psi \neq 1$ of F defining the Fourier transform, such that

$$\zeta(\hat{\Phi}, \check{f}, (1-s) + \frac{1}{2}) = \gamma(\pi, s, \psi) \zeta(\Phi, f, s + \frac{1}{2})$$

for all $\Phi \in C_c^\infty(A)$ and $f \in \mathcal{C}(\pi)$. Moreover,

$$\gamma(\pi, s, \psi) = \gamma(\chi_1, s, \psi) \gamma(\chi_2, s, \psi).$$

Proof. Since the zeta function is linear in the matrix coefficients, as is the operation $f \mapsto \check{f}$, it suffices to prove such γ exists for all $\Phi \in C_c^\infty(A)$ and f of the form $\gamma_{\tau \otimes \theta}$ as in the proof of Proposition 5.33. We calculated that

$$f(g) = \int_{B \backslash G} \tau(x) \theta(xg) d\dot{x} = \int_K \tau(k) \theta(kg) dk,$$

for some Haar measure dk on K , so that by right invariance of $d\dot{x}$ we have

$$\check{f}(g) = \int_{B \backslash G} \tau(xg) \theta(x) d\dot{x} = \int_K \tau(kg) \theta(k) dk.$$

The same computation as the proof of Proposition 5.33 gives (for the same K_1 and coset representatives k_i of K/K_1)

$$\begin{aligned}\zeta(\hat{\Phi}, \check{f}, (1-s) + \frac{1}{2}) &= \mu(K_1)^2 \sum_{i,j} \theta(k_j) \tau(k_i) \int_T (\hat{\Phi}^{ji})_T(t) \chi(t)^{-1} |\det t|^{1-s} dt \\ &= \mu(K_1)^2 \sum_{i,j} \theta(k_j) \tau(k_i) \int_T \widehat{(\Phi_T^{ij})}(t) \chi(t)^{-1} |\det t|^{1-s} dt\end{aligned}$$

by Lemma 5.39. Therefore, it suffices to show that

$$\int_{F^\times} \int_{F^\times} \widehat{(\Phi_T^{ij})}(t) \chi_1(t_1)^{-1} \chi_2(t_2)^{-1} |t_1 t_2|^{1-s} dt_2 dt_1 = \gamma(\chi_1, s, \psi) \gamma(\chi_2, s, \psi) \int_{F^\times} \int_{F^\times} \Phi_T^{ij}(t) \chi_1(t_1) \chi_2(t_2) |t_1 t_2|^s dt_2 dt_1$$

where $t = \begin{pmatrix} t_1 & 0 \\ 0 & t_2 \end{pmatrix} \in T$. By Theorem 5.18, this equality holds whenever we replace $\Phi_T^{ij} \in C_c^\infty(D)$ by a function of the form $\phi_{11}(t_1) \otimes \phi_{22}(t_2) \in C_c^\infty(D)$. But such functions span $C_c^\infty(D)$, so we are done by linearity of the integrals. \square

Definition 5.42. Define the Godement-Jacquet local constant $\epsilon(\pi, s, \psi)$ of $\pi = \iota_B^G \chi$ by

$$\epsilon(\pi, s, \psi) = \gamma(\pi, s, \psi) \frac{L(\pi, s)}{L(\tilde{\pi}, 1-s)}.$$

Corollary 5.43. For $\pi = \iota_B^G \chi$ we have

$$\epsilon(\pi, s, \psi) = \epsilon(\chi_1, s, \psi) \epsilon(\chi_2, s, \psi).$$

Proof. This follows from Proposition 5.41 and Proposition 5.33. \square

For context, we state more general versions of these results that hold for any irreducible smooth representation π of G .

Theorem 5.44. Let π be an irreducible smooth representation of G . There is a unique polynomial $P_\pi(X) \in \mathbb{C}[X]$, satisfying $P_\pi(0) = 1$, and

$$\mathcal{Z}(\pi) = P_\pi(q^{-s})^{-1} \mathbb{C}[q^{-s}, q^s].$$

Proof. [BH06, Theorem 24.2.1]. \square

Notation 5.45. Set $L(\pi, s) = P_\pi(q^{-s})^{-1}$.

Theorem 5.46. Let π be an irreducible smooth representation of G . There is a unique rational function $\gamma(\pi, s, \psi) \in \mathbb{C}(q^{-s})$ such that

$$\zeta(\hat{\Phi}, \check{f}, (1-s) + \frac{1}{2}) = \gamma(\pi, s, \psi) \zeta(\Phi, f, s + \frac{1}{2})$$

for all $\Phi \in C_c^\infty(A)$ and $f \in \mathcal{C}(\pi)$.

Proof. [BH06, Theorem 24.2.2]. \square

Definition 5.47. Define the Godement-Jacquet local constant $\epsilon(\pi, s, \psi)$ of an irreducible smooth representation π of G by

$$\epsilon(\pi, s, \psi) = \gamma(\pi, s, \psi) \frac{L(\pi, s)}{L(\check{\pi}, 1 - s)}.$$

Corollary 5.48. *The local constant satisfies the functional equation*

$$\epsilon(\pi, s, \psi) \epsilon(\check{\pi}, 1 - s, \psi) = \omega_\pi(-1).$$

The local constant is of the form

$$\epsilon(\pi, s, \psi) = a q^{bs}$$

for some $a \in \mathbb{C}^\times$, $b \in \mathbb{Z}$.

Proof. The first statement comes from the Fourier inversion formula and Theorem 5.46. The $\omega_\pi(-1)$ term comes from the minus sign in $\hat{\Phi}(x) = \Phi(-x)$ and the observation that for a matrix coefficient $f \in \mathcal{C}(\pi)$ we have $f(-g) = \omega_\pi(-1)f(g)$. The functional equation and Theorem 5.44 implies that ϵ is a unit in $\mathbb{C}[q^{-s}, q^s]$, and the units are precisely the elements of the form aq^{bs} for $b \in \mathbb{Z}$. \square

The Propositions 5.33 and 5.41 prove the Theorems 5.44 and 5.46 in the case that $\pi = \iota_B^G \chi$ and π is irreducible. As in Theorem 4.29, the representations $\pi = \iota_B^G \chi$ are typically irreducible - they are only reducible when $\chi = \phi \delta_B^{\pm 1/2}$ for some character ϕ of F^\times . In this case the composition factors are characters $\phi \circ \det$, and twists of Steinberg ϕSt_G . We state without proof the L -functions and local constants in the case that π is one of these composition factors. For more detail see Sections 26.5 - 26.8 of [BH06]. The results for all principal series representations are summarised in the following table:

Principal series representation π	$L(\pi, s)$	$\epsilon(\pi, s, \psi)$
$\iota_B^G \chi$, $\chi = \chi_1 \otimes \chi_2$, $\chi \neq \phi \delta_B^{\pm 1/2}$	$L(\chi_1, s) L(\chi_2, s)$	$\epsilon(\chi_1, s, \psi) \epsilon(\chi_2, s, \psi)$
$\phi \circ \det$, $\phi : F^\times \rightarrow \mathbb{C}^\times$ ramified	1	$\epsilon(\phi, s - \frac{1}{2}, \psi) \epsilon(\phi, s + \frac{1}{2}, \psi)$
ϕSt_G , $\phi : F^\times \rightarrow \mathbb{C}^\times$ ramified	1	$\epsilon(\phi, s - \frac{1}{2}, \psi) \epsilon(\phi, s + \frac{1}{2}, \psi)$
$\phi \circ \det$, $\phi : F^\times \rightarrow \mathbb{C}^\times$ unramified	$L(\phi, s - \frac{1}{2}) L(\phi, s + \frac{1}{2})$	$\epsilon(\phi, s - \frac{1}{2}, \psi) \epsilon(\phi, s + \frac{1}{2}, \psi)$
ϕSt_G , $\phi : F^\times \rightarrow \mathbb{C}^\times$ unramified	$L(\phi, s + \frac{1}{2})$	$-\epsilon(\phi, s, \psi)$

Figure 1: L -functions and local constants of principal series representations of G

In particular, if π is a composition factor of $\iota_B^G \chi$ then $L(\pi, s) = L(\chi_1, s) L(\chi_2, s)$, unless $\pi = \phi \text{St}_G$ for some unramified character $\phi : F^\times \rightarrow \mathbb{C}^\times$.

5.4 Converse Theorem

Attached to any principal series representation π of G we have an associated L -function $L(\pi, s)$ and local constant $\epsilon(\pi, s, \psi)$. In some sense this is enough information to distinguish them as irreducible smooth representations of G . More precisely, one can also define L -functions and local constants for the cuspidal representations of G , and then we have

Theorem 5.49 (Converse Theorem). *Let $\psi : F \rightarrow \mathbb{C}^\times$ be an additive character with $\psi \neq 1$. Let π_1, π_2 be irreducible smooth representations of $G = \mathrm{GL}_2(F)$. Suppose that*

$$L(\chi\pi_1, s) = L(\chi\pi_2, s) \text{ and } \epsilon(\chi\pi_1, s, \psi) = \epsilon(\chi\pi_2, s, \psi),$$

for all characters $\chi : F^\times \rightarrow \mathbb{C}^\times$. Then $\pi_1 \cong \pi_2$.

Recall that the twist $\chi\pi$ denotes the representation $g \mapsto \chi(\det(g))\pi(g)$.

We take as fact the following result for cuspidal representations.

Proposition 5.50. *Let π be an irreducible cuspidal representation of G . Then $L(\pi, s) = 1$.*

Proof. [BH06, Corollary 24.5]. □

Then we can distinguish between cuspidal and principal series representations as follows.

Proposition 5.51. *An irreducible smooth representation π of G is cuspidal if and only if $L(\phi\pi, s) = 1$ for all characters ϕ of F^\times .*

Proof. Since twisting preserves principal series representations, it preserves cuspidal representations. Proposition 5.50 implies that if π is cuspidal then $L(\phi\pi, s) = 1$ for all ϕ . In the other direction, suppose that π is a composition factor of $\iota_B^G \chi$ for $\chi = \chi_1 \otimes \chi_2$ a character of T . Taking $\phi = \chi_2^{-1}$, $\phi\pi$ is a composition factor of $\iota_B^G \phi\chi$ with $\phi\chi = \chi_1\chi_2^{-1} \otimes 1$. Now, except for the case $\phi\pi$ is a twist of Steinberg by an unramified character, we have $L(\phi\pi, s) = L(\chi_1\chi_2^{-1}, s)L(1, s)$, and then $L(1, s) = (1 - q^{-s})^{-1}$ is nontrivial. In the case it is a twist of Steinberg by an unramified character, the L -function is still nontrivial as seen in Table 1. □

Proof of Theorem 5.49 for principal series representations. Twisting π , we may assume that $L(\pi, s) \neq 1$ as in the proof of Proposition 5.51. Then $L(\pi, s)$ has degree 2 (as a rational function of q^{-s}).

Suppose $L(\pi, s)$ has degree 2. From Table 1, π is either $\iota_B^G \chi$ for some $\chi = \chi_1 \otimes \chi_2$, with $\chi_1\chi_2^{-1} \neq | - |^{\pm 1}$ and χ_i unramified, or $\pi = \phi \circ \det$ for some unramified character $\phi : F^\times \rightarrow \mathbb{C}^\times$. In either case, we have $L(\pi, s) = L(\chi_1, s)L(\chi_2, s)$ for unramified characters χ_i of F^\times , where $\pi = \phi \circ \det$ corresponds to $\chi_i = \phi| - |^{\pm 1}$. But since an unramified character χ is determined by $\chi(\varpi)$, it is determined by $L(\chi, s)$. Since $\iota_B^G(\chi_1 \otimes \chi_2) \cong \iota_B^G(\chi_2 \otimes \chi_1)$, it follows that $L(\pi, s)$ is enough to distinguish all principal series representations π for which $L(\pi, s)$ has degree 2.

Suppose $L(\pi, s)$ has degree 1, and is $L(\theta, s)$ for some unramified character θ of F^\times . From Table 1, π is either $\iota_B^G(\theta' \otimes \theta)$ for some ramified character θ' , or $\pi = \theta' \mathrm{St}_G$ for $\theta' = \theta| - |^{-1/2}$. In the latter case, θ' is unramified and so for any ramified character ϕ we have $L(\phi\pi, s) = 1$. This distinguishes it from the former case where if we take $\phi = (\theta')^{-1}$, a ramified character, we have $\phi\pi = \iota_B^G(1 \otimes \phi\theta)$ so that $L(\phi\pi, s) \neq 1$. To recover θ' in this case, we can choose some ramified character ϕ such that $L(\phi\pi, s) \neq 1$, say $L(\phi\pi, s) = L(\theta'', s)$ for a unique unramified character θ'' of F^\times . Since $\phi\pi = \iota_B^G(\phi\theta' \otimes \phi\theta, s)$, and $\phi\theta$ is ramified, we have $L(\phi\pi, s) = L(\phi\theta', s)$. Therefore $\theta' = \phi^{-1}\theta''$. □

Remark 5.52. The proof of Theorem 5.49 for principal series representations shows that the isomorphism class of π is determined solely by the L -functions $L(\phi\pi, s)$ as we range over all characters $\phi : F^\times \rightarrow \mathbb{C}^\times$. For cuspidal representations, all L -functions are 1 and they are instead distinguished solely by the local constants.

References

- [BH06] C. J. Bushnell and G. Henniart, *The Local Langlands Conjecture for $GL(2)$* , Grundlehren der mathematischen Wissenschaften, Springer Berlin, 2006.
- [Bum97] D. Bump, *Automorphic Forms and Representations*, Cambridge Studies in Advanced Mathematics, Cambridge University Press, 1997.
- [DE09] A. Deitmar and S. Echterhoff, *Principles of Harmonic Analysis*, Universitext, Springer, New York, 2009.
- [GH24] J. R. Getz and H. Hahn, *An Introduction to Automorphic Representations*, Graduate Texts in Mathematics, Springer Cham, 2024.
- [Tat67] J. Tate, *Fourier analysis in number fields and Hecke's zeta-functions*, Algebraic Number Theory, edited by J.W.S. Cassels and A. Frölich, London Mathematical Society, 1967.