

[Brief Review*]

Interior Product, Lie derivative and Wilson Line in the KBc Subsector of Open String Field Theory

Author: Hiroyuki Hata and Daichi Takeda

KBc algebra has been described classical solutions like tachyon condensation and multi- D_{25} branes. These solutions have singularities caused by $K \sim 0$, but reproduce the exact energies expected by string theory.

We addressed how to emerge Chan-Paton factors around the multi- D_{25} brane solution found by Murata and Schnabl [1]. Comparing with the action with Chan-Paton factors, the fluctuation $\Delta\Psi$ around the N -brane solution is expected to be expanded by the modified form of vertex operators $V_a^\dagger \mathcal{O}_F(k) V_b$ with some operators V_a ($a = 1, \dots, N$) carrying no ghost. Then we find that the question becomes whether there exist N operators V_a satisfying

$$V_a(\overleftarrow{Q}_B + \Psi_0) = 0 \quad \text{and} \quad V_a V_b^\dagger = \delta_{ab}, \quad (0.1)$$

where Ψ_0 is the N -brane solution and $A\overleftarrow{Q}_B = -(-1)^{|A|}Q_B A$. The similar equation for the former can be seen in Wilson lines of gauge theories, and the formal resemblance between SFT and Chern-Simons theory has been known, so we are driven to find the correspondence of Wilson line in SFT.

In the process to find this, we established the KBc manifold, which includes infinite sets of distinct (K, B, c) , of course satisfying KBc algebra, as its points. On the manifold, we succeeded in finding the interior product, Lie derivative and Lie bracket. The Lie derivative gives infinitesimal deformations in the space of solutions. By using the interior product, we finally obtained a candidate for the SFT Wilson line. However, the task to solve (0.1) is still left, and the information about KBc manifold may be not enough.

References

- [1] M. Murata and M. Schnabl, *On Multibrane Solutions in Open String Field Theory*, *Prog. Theor. Phys. Suppl.* **188** (2011) 50 [1103.1382].

* The reviewer: Daichi Takeda (takedai.gauge@gmail.com)