Dataset Information:

| Dataset IIIIOI | |
|--------------------------|---|
| Title | Burning - crop residues |
| Abstract | Greenhouse Gas (GHG) emissions from burning crop residues consist of methane and nitrous oxide gases produced by the combustion of a percentage of the crop residues burnt on-site. Computed at Tier 1 following the 2006 IPCC Guidelines for National GHG Inventories (IPCC, 2006); available by country, with global coverage and relative to the period 1990 onwards, with annual updates. |
| Supplemental | This domain contains data on GHG emissions, associated emission factors and underlying activity data. The FAOSTAT Emissions data are estimates by FAO and do not coincide with GHG data reported by member countries to UNFCCC. The database is intended primarily as a service to help member countries assess and report their emissions, as well as a useful international benchmark. The FAOSTAT Emissions data are disseminated publicly to facilitate continuous feedback from member countries. |
| Creation Date | 2012 |
| Last Update | 2013 |
| Data Type | Climate Change - Greenhouse Gases |
| Category | Environment |
| Time Period | 1990 onwards |
| Periodicity | Annual |
| Geographical Coverage | World |
| Spatial Unit | Country |
| Language | Multilingual (EN, FR, ES) |

Methodology and Quality Information:

Methods and processing

Greenhouse Gas (GHG) emissions from burning crop residues consist of methane (CH $_4$) and nitrous oxide (N $_2$ O) gases produced by the combustion of a percentage of crop residues burnt on-site. The mass of fuel available for burning should be estimated taking into account the fractions removed before burning due to animal consumption, decay in the field, and use in other sectors (e.g., biofuel, domestic livestock feed, building materials, etc.). The FAOSTAT data are estimated at Tier 1 following IPCC, 2006, Vol. 4, Ch. 2 and 5.

The CH₄ and N₂O emissions are estimated at country level, using the formula:

Emission = A * EF

where:

Emission = GHG emissions in units of g CH₄ and g N₂O;

A = Activity data, representing the total amount of biomass burned, kg of dry matter (1); EF = Tier 1, default IPCC emission factors, expressed in gr CH₄/kg of dry matter and gr N₂O/kg of dry matter (2).

- (1) Harvested area statistics are taken from FAOSTAT and used to estimate the amount of biomass burned using mean default crop values of mass of fuel available for combustion (M_B) and combustion factor (C_f) in IPCC, 2006: Vol.4, Ch. 2, Tab. 2.4. The crops included are: wheat, maize, rice and sugarcane. The mass is then reduced by the fraction of crop residue burnt on-site—assumed to be 10%, following IPCC, 2000: Ch.4, Section 4A.2.1.1.
- (2) Global default EF values taken from IPCC, 2006: Vol. 4, Ch. 2, Tab. 2.5.

Dimensionless conversion factors used are:

10⁻⁹, to convert the emissions from g CH₄ to Gg CH₄ and g N₂O to Gg N₂O;

GWP- CH_4 = 21 (100-year time horizon global warming potential), to convert $Gg CH_4$ to $Gg CO_2$ eq and

GWP-N₂O = 310 (100-year time horizon global warming potential), to convert Gg N₂O to

Gg CO₂eq (IPCC, 1996: Technical Summary, Tab. 4 pg. 22).

The burning crop residues domain contains the following data categories available for download: country-level GHG emissions, provided as total, in Gg CH_4 , Gg N_2O and Gg CO_2eq ; implied emission factors; and activity data. Data are available for all countries and territories, as well as for standard FAOSTAT regional aggregations, plus Annex I and non-Annex I groups. The data period is 1990 onwards, with annual updates.

Uncertainties in estimates of GHG emissions are due to uncertainties in emission factors and activity data. They may be related to, inter alia, natural variability, partitioning fractions, lack of spatial or temporal coverage, spatial aggregation. In the case of burning crop residues more detailed information is available in the guidelines (IPCC, 2006: Vol.4, Ch. 5, Section 5.2.4.4).

References

IPCC. 2000. Good practice guidance and uncertainty management in national greenhouse gas inventories. In: J. Penman et al. (Eds.), IPCC National Greenhouse Gas Inventories Programme, Technical Support Unit, Hayama, Japan.

IPCC. 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (Eds), IGES, Hayama, Japan.

Data Collection

Method

Completeness

100%

Computed

Links

www.fao.org/climatechange/micca/ghg/www.ipcc-nggip.iges.or.jp/public/

Distribution Information:

| Owner | FAO |
|----------|-----|
| Provider | FAO |
| Source | FAO |

Copyright Policy

Content on www.fao.org, its affiliated websites and specific pages (collectively "the FAO website") is protected by copyright. To ensure wide dissemination of its information, FAO is committed to making its content freely available and encourages the use, reproduction and dissemination of the text, multimedia and data products presented. Except where otherwise indicated, content may be copied, printed and downloaded for private study, research and teaching purposes, or for use in noncommercial products or services, provided that appropriate acknowledgement of FAO as the source and copyright holder is given and that FAO's endorsement of users' views, products or services is not stated or implied in any way. FAO encourages unrestricted use of news releases provided on the FAO website, and no formal permission is required to reproduce these materials.

All requests for translation and adaptation rights, and for resale and other commercial use rights should be addressed to copyright@fao.org or submitted via the online Licence Request Form when downloading.

Citation

FAO, 2013. FAOSTAT Emissions Database

http://faostat.fao.org/

Acknowledgements

The FAOSTAT Emissions database was produced by the FAO Monitoring and Assessment of Greenhouse Gas Emissions and Mitigation Potential in Agriculture project (MAGHG), with generous funding from the Governments of Norway and Germany, trust funds GCP/GLO/286/GER and GCP/GLO/325/NOR.