### **Dataset Information:**

Title	Rice cultivation
Abstract	Greenhouse gas (GHG) emissions from rice cultivation consist of methane gas from the anaerobic decomposition of organic matter in paddy fields. Computed at Tier 1 following the Revised 1996 IPCC Guidelines for National GHG Inventories (IPCC, 1997) and the IPCC 2000 Good Practice Guidance and Uncertainty Management in National GHG Inventories (IPCC, 2000); available by country, with global coverage and relative to the period 1990-2010, with annual updates.
Supplemental	This domain contains data on GHG emissions, associated emission factors and underlying activity data.  The FAOSTAT Emissions data are estimates by FAO and do not coincide with GHG data reported by member countries to UNFCCC. The database is intended primarily as a service to help member countries assess and report their emissions, as well as a useful international benchmark. The FAOSTAT Emissions data are disseminated publicly to facilitate continuous feedback from member countries.
<b>Creation Date</b>	2012
Last Update	2012
Data Type	Climate Change - Greenhouse Gases
Category	Environment
Time Period	1990 - 2010
Periodicity	Annual
Geographical Coverage	World
Spatial Unit	Country
Language	Multilingual (EN, FR, ES)

## **Methodology and Quality Information:**

# Methods and processing

GHG emissions from rice cultivation consist of methane gas ( $CH_4$ ) emitted by anaerobic decomposition of organic matter in paddy fields. The FAOSTAT data are computed at Tier 1 following IPCC, 1997 Vol. 3, Ch. 4 and IPCC, 2000, Ch. 4.

The emissions are estimated at country level, using the formula:

Emission = A \* EF

where:

Emission = GHG emissions in g CH<sub>4</sub> m<sup>-2</sup> yr<sup>-1</sup>;

A = Activity data, representing rice paddy annual harvested area in m<sup>-2</sup> (1); EF = Tier 1, default IPCC emission factors, in g CH<sub>4</sub> m<sup>-2</sup> yr<sup>-1</sup> (2).

- (1) Country area statistics are taken from FAOSTAT (domain: Production) and disaggregated into irrigated, rain-fed and upland water regimes, using default IPCC percentages in IPCC, 1997, Vol. 3, Ch. 4, Tab. 4.11.
- (2) Seasonally integrated EF values are specified for key rice producing countries in the guidelines (IPCC, 1997: Vol. 3, Ch. 4, Tab. 4.13). For countries with no default value for EF, the following strategy is applied: for Asia, countries are assigned an area-weighted average EF (15.7 g CH<sub>4</sub> m<sup>-2</sup> yr<sup>-1</sup>); in all other regions, countries are either assigned the IPCC EF value reported for a neighbouring country, where one existed, or they are assigned the IPCC global default EF value (20 g CH<sub>4</sub> m<sup>-2</sup>yr<sup>-1</sup>). Furthermore, seasonally integrated EF values are further modified by the application of a dimensionless scaling factor for water regime (i) and a dimensionless correction factor for organic amendments (ii).
- (i) The scaling factors for rice paddy water regime (IPCC, 1997: Vol. 3, Ch. 4, Tab. 4.12) are in the range 0-1. Specifically for all countries a scaling factor of 0.7 is used for rain-fed rice

and 0 for upland rice or dry conditions (IPCC, 2000: Tab. 3, page 403).

(ii) The correction factor for organic amendments is the default value of 2 for all countries, corresponding to the assumption that 40% of farmers use organic amendments (IPCC, 2000: Tab. 3, page 403).

Dimensionless conversion factors used are:

10<sup>-4</sup>, to convert the activity data from m<sup>-2</sup> to ha;

10<sup>-9</sup>, to convert the emissions from g CH<sub>4</sub> to Gg CH<sub>4</sub>; and

GWP-CH<sub>4</sub> = 21 (100-year time horizon global warming potential), to convert Gg CH<sub>4</sub> to Gg CO<sub>2</sub>eq. (IPCC, 1996: Technical Summary, Tab.4, pg. 22).

The rice cultivation domain contains the following data categories available for download: country-level GHG emissions both in Gg  $CH_4$  and Gg  $CO_2$ eq; implied emission factors; and activity data. Data is available for 126 individual countries and territories, as well as for standard FAOSTAT regional aggregations, plus Annex I and non-Annex I groups. The data period is 1990-2010, with annual updates.

Uncertainties in estimates of GHG emissions are due to uncertainties in emission factors and activity data. They may be related to, inter alia, natural variability, partitioning fractions, lack of spatial or temporal coverage, spatial aggregation. In the case of rice cultivation, more detailed information are available in the guidelines (IPCC, 2000: Ch. 4, Section 4.9.1.6).

#### References

IPCC. 1996. Climate Change 1995 - The Science of Climate Change: Contribution of Working Group I to the Second Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge.

IPCC. 1997. Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories, OECD, Paris.

IPCC. 2000. Good practice guidance and uncertainty management in national greenhouse gas inventories. In: J. Penman et al. (Eds.), IPCC National Greenhouse Gas Inventories Programme, Technical Support Unit, Hayama, Japan.

IPCC. 2002. Background Papers, IPCC Expert Meetings on Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. IPCC-NGGIP, p.399-417.

Data Collection

Computed

Method Completeness

100%

Links

www.fao.org/climatechange/micca/ghg/www.ipcc-nggip.iges.or.jp/public/

### **Distribution Information:**

OwnerFAOProviderFAOSourceFAO

**Copyright Policy** 

All rights reserved. FAO encourages the reproduction and dissemination of material published on this Web site. Non-commercial uses will be authorized free of charge, upon request. Reproduction for resale or other commercial purposes, including educational purposes, may incur fees. Applications for permission to reproduce or disseminate FAO copyright material, and all queries concerning rights and licences, should be addressed by e-mail to copyright@fao.org or to the:

Chief, Publishing Policy and Support Branch,

Office of Knowledge Exchange, Research and Extension

FAO, Viale delle Terme di Caracalla

	00153 Rome, Italy
	Note: Hyperlinks to non-FAO Internet sites do not imply any official endorsement of or responsibility for the opinions, ideas, data or products presented at these locations, or guarantee the validity of the information provided. The sole purpose of links to non-FAO sites is to indicate further information available on related topics.
Citation	FAOSTAT, 2012. FAOSTAT Emissions Database. www.faostat.fao.org
Acknowledgement	The FAOSTAT Emissions database was produced by the Monitoring and Assessment of Greenhouse Gas Emissions and Mitigation Potential in Agriculture project of the MICCA Programme, with generous funding from the Governments of Norway and Germany, trust funds GCP/GLO/286/GER and GCP/GLO/325/NOR.