Both diet and sociality affect primate brain-size evolution

Mark Grabowski[a,b], Bjørn T. Kopperud[c,d], Masahito Tsuboi[b,e], Thomas F. Hansen[b]

[a]Research Centre in Evolutionary Anthropology and Palaeoecology, Liverpool John Moores University, 3 Byrom Street, Liverpool L3 3AF, UK

[b]Department of Biology, University of Oslo, Blindernveien 31, 0371 Oslo, Norway

[c]GeoBio-Center LMU, Ludwig-Maximilians-Universität München, Richard-Wagner-Str. 10 80333 Munich, Germany

[d]Department of Earth and Environmental Sciences, Paleontology and Geobiology, Ludwig-Maximilians-Universität München, Richard-Wagner Straße 10, 80333 Munich, Germany

[e]Department of Biology, Lund University, Ekologihuset, Sölvegatan 37, 223 62 Lund, Sweden

Corresponding Author: Mark Grabowski, m.w.grabowski@ljmu.ac.uk

Acce

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

© The Author(s) 2022. Published by Oxford University Press, on behalf of the Society of Systematic Biologists. All rights reserved. For permissions, please email: journals.permissions@oup.com

Abstract

Increased brain size in humans and other primates is hypothesized to confer cognitive benefits but brings costs associated with growing and maintaining energetically expensive neural tissue. Previous studies have argued that changes in either diet or levels of sociality led to shifts in brain size, but results were equivocal. Here we test these hypotheses using phylogenetic comparative methods designed to jointly account for and estimate the effects of adaptation and phylogeny. Using the largest current sample of primate brain and body sizes with observation error, complemented by newly compiled diet and sociality data, we show that both diet and sociality have influenced the evolution of brain size. Shifting from simple to more complex levels of sociality resulted in relatively larger brains, while shifting to a more folivorous diet led to relatively smaller brains. While our results support the role of sociality, they modify a range of ecological hypotheses centered on the importance of frugivory, and instead indicate that digestive costs associated with increased folivory may have resulted in relatively smaller brains.

Keywords: primate brain size, social-brain hypothesis, energetic constraints, adaptation, phylogenetic comparative methods, allometry, evolutionary trend, *Slouch, bayou*

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Brain size varies greatly among primates (e.g. Jerison 1979) and a wealth of comparative studies have attempted to determine the factors that drove brain-size evolution (Table 1). These factors generally fit into two categories - ecological and social – with different studies supporting the importance of one category over the other. Pioneering work by Clutton-Brock and Harvey (1980) pointed to the role of ecology, with factors such as diet and home-range size explaining a proportion of interspecific variation in relative brain size. Ecological hypotheses posit that large brains with presumed greater cognitive ability (e.g. Deaner et al. 2007) are adaptations to deal with ecological problems, such as remembering the location of fruiting plants (see also Harvey et al. 1980). Later studies focused on the role of sociality in promoting larger brains and reported relationships between relative brain size and different metrics of social behavior (e.g. Dunbar 1998). Sociality and ecology may also interact. The social-brain hypothesis (*sensu stricto*) emphasizes large brains as an adaptation to ecological problem solving through social interactions (Dunbar 1992, 1998; Barton 1996; Dunbar and Schultz 2017). In addition, brain-size adaptation must balance energetic constraints. Large brains are energetically expensive, in humans accounting for about 20% of resting metabolic rate in adults, ~ 52% in infants, and peaking at ~66% during childhood (Kuzawa et al. 2014). Large-brained species may be those that have been better able to meet or have evolved solutions to these constraints (Aiello and Wheeler 1995; Martin 1996; Wrangham et al. 1999; Isler and Van Schaik 2009); for example through greater maternal investment (Martin 1996; Powell et al. 2019). Despite decades of research, however, the drivers of primate brain-size evolution remain in dispute (Table 1; see also Whiting and Barton 2003; Dunbar and Schultz 2017).

Conflicting results from comparative studies are partly attributable to a variety of methodological issues, some of which have only recently begun to be addressed. First, small sample sizes (both intra- and inter-specific) and sub-optimal quality of the underlying data can lead to errors (e.g. Smith and Jungers 1997; Borries et al. 2013; Sandel et al. 2016; Gilbert and Jungers 2017). Second, misinterpretation of statistical significance or model-selection criteria as measures of biological effect can be misleading. This is particularly pertinent when the absence of statistical

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

significance is misinterpreted as evidence for the absence of a biological effect. Third, testing factors one at a time rather than in a multivariate fashion ignores associations that can be driving results (Dunbar and Shultz 2017). Fourth, few comparative studies have accounted for observation or measurement error in species means (but see Grabowski et al. 2016), which can cause inaccurate parameter estimates (Hansen and Bartoszek 2012; Garamszegi 2014; Morrissey 2016). Finally, statistical techniques to correct for shared ancestry vary dramatically in underlying models of trait evolution, and differences in those models can lead to different conclusions and interpretations (Garamszegi and Mundry 2014; Hansen 2014).

Phylogenetic generalized least squares with Brownian-motion assumptions for the residuals (i.e. assuming phylogenetic covariances proportional to shared branch length) is currently the most widely used approach to test hypotheses about brain-size evolution (Table 1) but is not a suitable model to study adaptation to external factors (Hansen 1997, 2014; Hansen and Orzack 2005; Labra et al. 2009; Uyeda et al. 2018; Moen et al. 2022). The main problem is that this approach implicitly assumes that adaptation is instantaneous. This is not only incompatible with the simultaneous assumption of phylogenetic correlations in the residuals, but also neglects the past history of associations between species and the niches they are adapting to. A species that has only had a short association with a given selective regime is expected to be less influenced by this regime than a species with a long history of association. Furthermore, past associations may still have an influence on the current state of the trait if the species have only recently shifted to their current niches. These effects all depend on the rate of adaptation. It is essential to estimate this rate and then use it to set up a consistent model of the evolutionary process and a correct comparative analysis (Hansen, 1997; Hansen et al. 2008; and see methods below). As adaptation is at the core of most major hypotheses about brain-size evolution, the common reliance on an unsuitable evolutionary model could be a major issue in comparative studies of brain size.

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Here we test hypotheses on the role of ecology and sociality in primate brain-size evolution using phylogenetic comparative methods specifically designed to jointly estimate and account for the historical effects of adaptation and shared ancestry (Hansen 1997; Hansen et al. 2008). Our analysis uses the largest current sample of wild-caught primate average brain and body sizes with observation error – 183 species – complemented by newly compiled diet data from field studies and sociality data from published compilations. Our goal is to move beyond testing whether particular predictors had a significant effect, toward a quantitative understanding of how diet and sociality have affected brain-size evolution through the history of primates.

Materials and Methods

Brain- and Body-Size Data

Average endocranial volume and body mass from 183 primate species based on 3,506 individual observations for brain size and 6,608 individual observations for body mass were taken from Isler et al. (2008). For our main analyses, each species mean was represented by at least one male and one female. Measurement variance due to estimation error in species means for both log endocranial volume (in cubic centimeters or *cc*) and log body mass (in grams or *g*), which is quantified by the standard error of the estimate, was calculated following Grabowski et al. (2016; see also Tsuboi et al. 2018).

Diet Metrics

We used percentage feeding time on food items compiled from field studies, supplemented by published compilations of field studies (Campbell et al. 2011; Powell et al. 2017), and divided into four categories following Nunn and Van Schaik (2002): (1) fauna (e.g. insects and other small animals), (2) plant reproductive parts, (3) plant vegetative parts, and (4) plant exudates including gum. Percentage feeding time in each category was converted into two broader categorical variables

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

– frugivorous/non-frugivorous and folivorous/non-folivorous – based on whether plant reproductive parts (e.g. fruit, flowers, seeds, buds, nectar), or plant vegetative parts (e.g. leaves, roots, and other plant parts) occupied a larger percentage of feeding time than the three other broad categories listed above (see also Sailer et al. 1985; Powell et al. 2017), and this scheme was kept constant across species. We chose this simple categorical scheme rather than analyzing continuous percentages or more detailed classification schemes for two reasons. First, because percentage data are likely to violate the unboundedness, homoscedascity, and normality assumptions of standard comparative methods based on Brownian-motion or Ornstein-Uhlenbeck processes, and second, because classification into more dietary categories will generate more parameters relative to the available number of species (see Text S1). Following Powell et al. (2017) we also ran separate analyses in which folivores were restricted to only those species with clear physiological adaptations for folivory (Hladik 1978; Chivers and Hladik 1980; Clutton-Brock and Harvey 1980).

Sociality Metrics

We used average group size both as a continuous variable and as a discrete variable with five categories (groups of 1-5, 6-10, 11-25, 26-50, and >50 individuals) based on data from Powell et al. (2017) and supplemented by data from DeCasien et al. (2017). We used a four-way social system (solitary, pair-living, uni-male and multi-female, multi-male and multi-female) and a five-way mating system (spatial polygyny, monogamy, polyandry, harem polygyny, polygynandry) following the classification of DeCasien et al. (2017), supplementing their published dataset with 32 additional species from the same published compilations (Plavcan 1999; Thorén et al. 2006; Shultz et al. 2011). Finally we reclassified social system into the metric of social complexity following DeCasien and Higham (2019) using a three-way category scheme based on general group type - solitary, pair-living, or group-living. We did not run models with group size categories, social system, and mating system together because these factors are closely related to each other. We also did not run models with

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

different categorizations of sociality or average group size and group size categories at the same time for the same reason.

Activity Period

Although not a primary focus of our study, we included activity period as a covariate because previous analyses have suggested that it has a role in brain evolution (Barton 1996, 1999; Barton et al. 1999; Powell et al. 2017; DeCasien et al. 2019). Based on data from Powell et al. (2017) and supplemented by data from published compilations (MacLean et al. 2009; Shultz et al. 2011; Mittermeier and Wilson 2013; Santini et al. 2015), we treated activity period as a dichotomous variable (diurnal/nocturnal). Following Powell et al. (2017) the few cathemeral species were classified as diurnal. While activity period is generally grouped with ecological variables such as diet (e.g. Barton 1996), it has been shown to interact with sociality metrics (Shultz et al. 2011). It is thus not clear to which group it should be assigned.

Note that we specifically focus on factors that underlie the role of ecology and sociality; hence behavioral factors previously identified as influencing brain size, such as cognitive performance (e.g. MacLean et al. 2014) are not included here, though they are undoubtedly important.

Phylogeny

We used the dated consensus tree from version 3 of 10Ktrees website using the Genbank taxonomy (Arnold et al. 2010), as has been done in numerous recent studies on brain-size evolution in primates (DeCasien et al. 2017; DeCasien and Higham 2019; Todorov et al. 2019). Our individual analyses were based on subsets of species due to missing data for some predictor variables. The

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

main set of analyses were based on 128 species with complete diet, sociality, and activity-period data; our univariate analyses were subsets of the 183 species dataset. We note that while there are by some accounts more than 500 living species of primates (Estrada et al. 2018), the phylogeny used here is based on 301 species. Our complete data on endocranial volume, body mass, diet, sociality, and activity period are provided in the supplementary material.

Comparative Approach

Following Hansen (1997) we model evolution of log endocranial volume as an Ornstein-Uhlenbeck process around a (primary) optimal state that is a function of different predictor variables (diet, social system, etc.) mapped as regimes on the phylogeny for the categorical predictors and of a randomly changing predictor variable for log group size (Hansen et al. 2008). Following Grabowski et al. (2016) we augment this with a "direct" effect of log body mass such that evolutionary changes in body size are associated with an immediate correlated response in brain size. The model is described by the stochastic differential equations:

$$dy = -\alpha(y - \theta(z, g))dt + bdx + \sigma_y dB_y,$$

$$dg = \sigma_g dB_g ,$$

$$dx = \sigma_x dB_x,$$

where y is log endocranial volume, g is log group size, and x is log body mass. The dB_y dB_g and dB_x are independent white-noise processes multiplied by their standard deviations σ_y, σ_g, and σ_x. The α-parameter quantifies the rate of adaptation toward a niche-specific optimum, θ, modeled as a function of ecological predictors, z, such as diet and social system, mapped on the phylogeny, as well

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

as *g*, log group size, modeled as evolving according to a Brownian-motion process. The α-parameter can be expressed as a phylogenetic half-life, $t_{1/2} = \ln(2)/\alpha$, the time it takes for the response variable to evolve in expectation half the distance from its ancestral state toward the current optimum. The parameter *b* gives the direct effect of changes in body size on brain size, as may be expected from a within-species, static, allometry. In the results, we report σ_y by reparametrizing as the stationary variance of the Ornstein-Uhlenbeck process as $v_y = \sigma_y^2/2\alpha$, the expected residual variance when species have evolved under constant adaptive regimes for a long period of time.

In many of our analyses the phylogenetic half-life becomes very large at the same time as the optima are estimated to be far outside the range of the species data. This is not uncommon in Ornstein-Uhlenbeck models of adaptation but requires a reparameterization and some reinterpretation (Hansen 1997). An infinite half-life is a characteristic of a Brownian motion, but when this happens simultaneously with infinitely large optima, the deterministic (adaptation) part of the model does not vanish, and the model converges on a Brownian motion with niche-specific deterministic trends. These trends can be reliably estimated as $\tau = \lim_{\alpha \to 0} \alpha\theta$, even when the α and θ parameters are individually inaccurate (Hansen 1997). The biological interpretation is that the species are far from their niche-specific optima and continuously evolving toward them at niche-specific rates. We estimate these "adaptive" trends, $\tau(z, g)$ with units of trait change per tree height, as functions of ecological variables such as diet and social systems in the same way as for the primary optima described above. Note that this trend model is different from standard Brownian-motion-based PGLS in that the deterministic prediction for each species is a weighted sum of the different trends it has experienced through its history. It thus incorporates (or models) historical information about past selective regimes.

Since all species are extant, the trends are only meaningful as contrasts with each other, and not as estimates of absolute trends (Hansen 1997). As our phylogeny is scaled to unit length = 1 (original height 73 million years or *myr*), a contrast of 50% per tree height when switching from

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

spatial polygyny to polygynandry would mean that relative endocranial volume is expected to be 50% larger in the polygynandric regime than it would have been in the spatial polygynic regime for species that have remained in those regimes over their entire history from the root of the tree.

We start by using *bayou* version 2.2.0 (Uyeda and Harmon 2014) to search for shifts in the adaptive optima of the Ornstein-Uhlenbeck process and allometric parameters across the phylogeny without an *a priori* hypothesis. Here we use the latest version of *bayou* to detect shifts in allometric exponents for the relationship between log endocranial volume and log body mass (see also Miller et al. 2019), a function that was not available prior to version 2.0. We ran six chains, two for each model, each for three million generations. Convergence was measured using Gelman's R statistic and was less than 1.005 for α, σ_y^2, and log*L*. Following Uyeda et al. (2017), we initialized the MCMC without any shift proposals for the first 10,000 generations to improve the fit of the model. Priors for the parameters as defined above were set to:

$\alpha \sim \log N(\mu = \log(0.5), \sigma = 2.0)$

$\sigma_y^2 \sim$ half-Cauchy ($\mu = 0$, $\sigma = 0.1$)

$b \sim N(\mu = 0.6, \sigma = 0.1)$,

Here, the prior on α is a distribution of $t_{1/2}$ in which the lower 10% is equivalent to 1 million years or less, and the upper 10% starts at two times the height of the tree. The prior on b is based on the global slope estimate for primates found in Grabowski et al. (2016). Model comparisons were performed using Bayes factors following Uyeda et al. (2017).

We next tested for relationships between relative brain size and metrics of diet and sociality with *slouch* 2.1.4 (Hansen et al. 2008; Kopperud et al. 2020), using the location of the *bayou*-

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

discovered regime shifts to inform our analyses. *Slouch* uses generalized least squares to estimate primary optima or trends in the Ornstein-Uhlenbeck framework conditionally on the other parameters of the model, which are estimated by maximum likelihood. Contrasts between trends with standard errors are calculated in *slouch*, and we give a specific example of how this is done in Text S4.

Apart from some colinear combinations indicated above, we compared all possible combinations of factors hypothesized to influence brain size with likelihood using AICc. All models accounted for measurement error in both log endocranial volume and log body mass following previous approaches (Hansen and Bartoszek 2012; Grabowski et al. 2016), a feature that is built into *slouch* (Kopperud et al. 2020). We assessed the fit of the model for the primary optimum (i.e. the adaptive landscape) with a residual R^2; i.e., the percent of residual variance explained by the predictors after accounting for body mass. We did this because log body mass itself explains a large proportion of the variance in log endocranial volume – around 80% on average. Thus, explaining half of the remaining 20%, though only 10% of the total variance, is arguably a substantial result, which is made clear by this metric.

To assess uncertainty in the evolutionary history of the discrete predictors, diet, social categories, and activity period were first mapped on the full phylogeny using stochastic character mapping using the make.simmap function from phytools (Revell 2011) to fit a continuous time-reversible Markov-chain model, which was then used to simulate character states on the phylogeny. For each model including discrete predictors, we used AICc to choose the structure of the transition rate matrix from the possibilities of equal rates, symmetric rates, or all rates different (Table S1). After running all simulated mappings in *slouch*, their AICc distributions were used to qualitatively inform model selection. The final regime assignments were the maximum a posteriori regime estimates for each node in the Markov-chain model (Figs. S1-S8) with the transition rate matrix with

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

the lowest AICc score. None of the results were qualitatively changed by the analyses of stochastic maps and we do not discuss these further (see Text S2).

Results

Shifts in Optima

The *bayou* analysis favored two shifts in the relative brain-size optimum with posterior probabilities (*pp*) higher than 0.5. One near the origin of the haplorhines after the split from tarsiers and the other within platyrrhines (*pp* = 0.90, and *pp* = 0.78, respectively; Fig. 1; Fig. S9; Table S2). Therefore, we used the two *bayou*-located shifts to inform our *slouch* analyses. There was no evidence for changes in the allometric slope, 0.56 (95% highest posterior density (HPD) of 0.5-0.62), and the shifts in the intercept model were supported with a Bayes Factor of 11.04 over a model with no change and 9.67 over a model with shifts in both intercept and slope. The posterior median phylogenetic half-life for this model was $t_{1/2}$ = 67% of tree height with a 95% HPD of 32%-∞ (Table S2; Text S3). A third possible shift, along the lineage leading to the Aye-aye *Daubentonia madagascariensis*, had a posterior probability slightly below our 0.5 cutoff (*pp* = 0.389). As pointed out by Uyeda et al. (2017), it is not possible to determine if shifts on branches leading to singleton taxa are the result of shifts in the slope or in the intercepts of allometric relationships.

Adaptation to Diet and Social Parameters

The best model for the evolution of brain size across primates as a whole included distinct regimes for diet (strict folivorous/non-folivorous) and mating system, with group size added as a continuous variable. Together these factors explained 31% of the residual variance in brain size after accounting for body size (Table 2; Tables S3, S4). The phylogenetic half-life for this model was 263%

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

percent of the tree height (2-unit support interval 43% of tree height-∞), which means that relative brain-size evolution behaves roughly as Brownian motion with distinct evolutionary trends in the different selective regimes (Figure 2a).

Based on reconstructed shifts in diet and mating systems (Fig. S10), one of the largest upward shifts in trends occurred with a change from spatial polygyny to polygynandry in species maintaining a non-folivorous diet (Fig. 2a). The estimated contrast for this shift was 1.11±0.27 log *cc* per tree height, corresponding to an Exp[1.11] = 303% difference or a threefold change over the time span of the phylogeny (Table S4; Text S4). This coincides with one of the shifts located by *bayou* (Fig. 1), which occurred after tarsiers split off from other haplorhines. The other *bayou*-located shift, near the origin of the Callitrichidae, is the largest downward shift with a contrast of -2.85±0.52 log *cc* per tree height, corresponding to a 94% downward shift in the trend associated with a change from polygynandry to polyandry while maintaining a non-folivorous diet, but we caution that this latter shift is based on a small number of polyandrous species.

The largest predicted shifts were for species switching to non-folivorous harem polygyny from any of the other regimes, which occurred multiple times across the phylogeny. The second largest shift is associated with switching to non-folivorous polygynandry (Fig. 2a; Table 2; Fig. S10).

Group size had a small negative effect on relative brain size with an evolutionary slope of -0.08±0.03 log *cc* per tree height, meaning that a doubling of group size would lead to a 2.7% decrease in brain size over the time span of the phylogeny (Text S4).

Because shifts in diet happen in the context of a variety of mating systems, it is difficult to fully disentangle the two, but it is clear that a folivorous diet is associated with a downward shift in the brain-size trend (Fig. 2a; Fig. S10). Shifting from non-folivorous polygynandry to the folivorous variant is predicted to decrease the adaptive trend with a contrast of -0.67±0.54 log *cc* per tree height – an almost 50% decrease over the time span of the phylogeny. Similarly, shifting from non-

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

folivorous to folivorous spatial polygyny decreases the trend with -0.44±0.53 log *cc* per tree height – an 36% reduction over the time span of the phylogeny (Fig. 2a; Table S4).

A model with only the two *bayou*-located regime shifts alone or combined with group size did not improve the fit (Table S5). This shows that while some shifts may correspond to those located by *bayou*, the combination of diet categories and sociality has better explanatory power.

Using male or female averages separately (Text S7) or rerunning analyses with single predictors (Text S8) gave results consistent with the main findings both for primates as a whole and for the subclades. The smaller residual R^2 values for single predictors illustrates the importance of using multiple predictors in explaining relative brain size (Text S8).

Variation within Taxonomic Groups

Based on the *bayou*-located shifts, we divided the phylogeny into three major subsets – strepsirrhines plus tarsiers, haplorhines minus tarsiers, and platyrrhines. We used these subsets to determine if relative brain size evolved differently in the different groups. As only one haplorhine species included here was nocturnal (*Aotus trivirgatus*), activity period was not included as a predictor in the haplorhine or platyrrhine subsets.

Strepsirrhines. For strepsirrhines plus tarsiers (n = 28), the best model included group size, mating system, and activity period, together explaining 65% of the residual variance and 95% of the total variance in brain size (Table 3, Tables S6, S7). The phylogenetic half-life for this model was infinity with a 2-unit support interval of 43% - ∞, which means that evolution behaves as a Brownian motion with distinct trends in the different selective regimes (Fig 2b). Based on reconstructed shifts (Fig. S11), we found large changes associated with two regime switches. Going from nocturnal

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

monogamy to diurnal polygynandry was associated with a contrast of 3.15±0.71 log *cc* per tree height, which would correspond to a 23-fold difference in brain size over the time span of the phylogeny. Switching from nocturnal to diurnal monogamy was associated with a contrast of 1.86±0.46 log cc per tree height, which would correspond to a six-fold difference over the time span of the phylogeny. These extreme differences are of course not realized, as most species have spent relatively short amounts of time in the diurnal monogamous and polygynandric regimes (Fig. S11) and are thus predicted to be far from their optimal brain sizes (Fig. 2b). Within mating-system and activity-period categories, group size had a negative effect, with an evolutionary slope of -0.53±0.11 log *cc*, meaning that a doubling of group size would lead to a 17% reduction over the time span of the phylogeny (Text S5). Because this negative effect of group size did not appear in the other subclades, it is likely that this result in strepsirrhines drives the effect found for all primates.

Haplorhines. For haplorhines excluding tarsiers (n = 100), the best model included diet categories (strict folivorous/non-folivorous) and mating system explaining 32% of the residual variance (Table 4; Tables S8, S9). As the shift to polygynandry from spatial polygyny had already taken place at the root of the haplorhine clade, it is likely that diet and mating system may be playing a further role in brain-size patterning within the haplorhines. The phylogenetic half-life for this model was also infinity (2-unit support interval 43% - ∞). Overall trends (Fig. 2c; Fig. S12) were similar to the results for all primates though effects associated with non-folivorous spatial polygyny had higher standard error due to the small number of species in this regime. In addition, species displaying folivorous polygynandry and folivorous harem polygyny were distant from their predicted brain sizes (Fig. 2c) as expected from the short amount of time they have spent in these regimes (Fig. S12). The model predicts that brain size is still decreasing in these species.

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Platyrrhines. For platyrrhines (*n* = 32), diet categories (strict folivorous/non-folivorous) and mating system explained 93% of the residual variance, and together with body mass, 98% of the total variance in log brain size (Table 5; Tables S10, S11). In contrast to the other clades, the phylogenetic half-life was much shorter, 6% (2-unit support interval 6%-92%) of the total height of the platyrrhine phylogeny (22 *myr*) corresponding to 1.32 myr. This suggests that brain size is evolving around estimable evolutionary optima influenced by the combination of diet and mating system (Fig. 2d; Fig. S13). Species displaying folivorous harem polygyny, non-folivorous monogamy, non-folivorous polyandry, and non-folivorous polygynandry are all near their adaptive optima, though shifting from the latter to any of the former decreases the optimal brain size in platyrrhines by 52% (Table 5; Text S6). Alternatively, shifting to non-folivorous harem polygyny from non-folivorous polygynandry increases the brain-size optima by 223%, though species in the former regime are distant from their predicted brain-size optima (Fig. 2d) as expected from the short amount of they have spent in these regimes (Fig. S13; Text S6).

Discussion

Overall, we found a relative decrease in brain size when shifting from frugivory to folivory, and a relative increase when shifting to more complex social systems. As the predictors of brain size for primates in general appear to be generated by a combination of differences between and within suborders, we discuss each clade in turn rather than attempting to interpret our detailed results as being general to all primates. This being said, our finding that both diet and sociality effect brain-size evolution stands in contrast to a number of recent analyses that favored ecological rather than social factors (e.g. DeCasien et al. 2017; Street et al. 2017), or social rather than ecological factors (e.g. Todorov et al. 2019). The phylogenetic heterogeneity of these factors makes it likely that at least some of the disparate results may be an outcome of differences in taxonomic sampling. Our results also illustrate how choice of models for phylogenetic comparative analysis can affect the outcome.

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Comparisons with Previous Studies and Biological Interpretations

*Strepsirrhines***.** Our finding that group size, mating system, and activity period affected brain-size evolution, both complement and contrast those of previous studies. In a study of 19 different lemur species, MacLean et al. (2009) concluded that diet and activity period (cathemeral, diurnal, or nocturnal), rather than social group size or pair bonding, were associated with relatively larger brains. Like us they found a tendency for a negative effect of group size on brain size but discounted the result due to a lack of statistical power. The negative effect of group size in our study was also relatively weak. While a negative effect of group size may not support the social-brain hypothesis, it must be noted that this is a residual effect from a model with stronger effects of social system. For example, brain evolution was accelerated by a shift to diurnal polygynandry from diurnal monogamy, with the former arguably a more complex social system than the latter. We also note that the negative group-size effect in primates as a whole appears to be solely due to the negative relationship within the strepsirrhines.

MacLean et al. (2009) also reported an effect of diet on brain size based on reduced major axis regression, but no effect when using independent contrasts. This latter finding is mirrored in our results because the relationship between diet and brain size in strepsirrhines disappears if we assume strong phylogenetic signal (i.e., Brownian motion). Brain size is adapting to diet on a short time scale relative to the time span of the phylogeny, however, taking on average 4.4 *myr* to evolve half the distance to the new optima. This model indicates that folivory is associated with a 28% reduction in optimal brain size (Text S5).

*Haplorhines***.** Overall, we found that harem polygyny combined with a non-folivorous diet is associated with the fastest increases in relative brain size. This result differs from other findings of a

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

relationship between brain size and mating system in haplorhines, which pointed to monogamous (Schillaci 2006), or polygynandrous (Shultz and Dunbar 2007) species as possessing the largest brains. One explanation for these divergent results is that when mating system is considered independently from diet, the pattern is reversed, with shifting to harem polygyny from polygynandry resulting in a 48% decrease in the rate of relative brain-size evolution over the time span of the phylogeny (Text S9). Both harem polygyny and polygynandry are complex mating systems when compared to spatial polygyny or monogamy, lending support to Dunbar's (1998) hypothesis that the ability to manage social interactions within these systems required larger brains.

On the other hand, haplorhines adapted to eating foliage appear to have relatively smaller brains. As frugivory was likely the ancestral diet for haplorhines (Supplementary Fig. S10 and see Soligo and Martin 2006), haplorhines that evolved away from this diet may have done so recently. Together with long phylogenetic half-lives, this finding suggests that brain size in folivores may continue to diverge downward relative to non-folivores. Contrary to suggestions by Clutton-Brock and Harvey (1980) and DeCasien et al. (2017) that brain sizes tend to increase in frugivorous species, our results are more consistent with a decrease in folivorous species. As discussed further below, this does not rule out a role for frugivory in the origination or maintenance of large brains in some lineages.

Our findings illustrate how studying adaptation using methods that account for the evolutionary history of the selective environment, such as non-folivory vs. folivory, can affect the results and their biological interpretation. Our findings are consistent with suggestions that the smaller brains of folivores are the result of folivores having evolved a large stomach and digestive system (Hladik 1967), which may require shifts in energy allocation away from the brain (Aiello and Wheeler 1995; Isler and Van Schaik 2009; Fonseca-Azevedo and Herculano-Houzel 2012). An alternative interpretation is that large brains in primates compared to other mammals are maintained as an adaptation to ancestral frugivory and the associated ecological problem solving. If

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

so, this could be an example of “stasis is data” (Gould and Eldredge 1977), where energetically expensive large brains are maintained by stabilizing selection in frugivorous lineages while lineages that no longer required these functions shifted resources to digestion.

*Platyrrhines***.** The rapid adaptation of brain size given the best-fitting model (strict diet categories + mating system) reveals the distinctive nature of platyrrhines compared to other primates. The fact that diet categories, mating system, and body mass together explain 98% of brain-size variance in platyrrhines may reflect both their narrow ecological range and their relatively recent adaptive radiation - the last common ancestor of all living platyrrhines is dated to the early Miocene, about 20 Ma (Fernandez-Duque et al. 2012), and there may not have been enough time for species to adapt to divergent selection pressures since they arose. This is illustrated by the brain sizes of species exhibiting non-folivorous harem polygynandry being much smaller than their predicted primary optimum – given time, we predict that species in this regime will approach their optima. Our findings are complementary to Aristide et al. (2016), who suggested that brain shape, rather than size as studied here, is related to diet, locomotion, and social group size. In fact, we found that group size alone was the third best supported model, after a model including diet categories, mating system, and group size. Platyrrhines are noted for having a wide range of social systems and variation in group size and composition, but a narrow ecological range (Fernandez-Duque et al. 2012).

Our ancestral-state reconstruction (Fig. S13) indicates that non-folivorous polygynandry was ancestral for platyrrhines and maintained in other platyrrhine lineages while the family Callitrichidae shifted to monogamy and polygynandry. The Callitrichidae appear to follow the same brain-body relationship found across strepsirrhines and tarsiers (Fig. S13). Martin (1992) referred to the Callitrichidae as a dwarfed lineage, and it is possible that the brain-body relationship is constrained to evolve in a narrow range, with Callitrichidae mirroring strepsirrhines and tarsiers through convergent evolution. This result is further supported by the *Alouatta* clade, which appears to have

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

experienced a similar reduction in brain size in response to a shift to folivory (Fig. S13). Brain-size variation may then reflect variation in the evolutionary allometric intercept, as found in other group of vertebrates (Tsuboi et al. 2018) as well as in allometric relationships of other morphological traits (Voje et al. 2014).

Caveats and Conclusions

Brains are multifunctional organs interacting with all aspects of an animal's life. Consequently, we expect multiple sources of selection and constraint to have potential impact on brain size and structure. However, only a small set of potentially relevant factors have been investigated and interpreted, often in isolation from each other. While past studies have sometimes suggested a lack of relationship with this or that factor, many suffer from the issues mentioned in the introduction, including small sample sizes, misinterpretation of statistical significance as measures of biological effect, and using statistical models that do not match the biological context and question. Most likely, brain evolution has been influenced by a multitude of ecological, social, behavioral, physiological, and sensory factors that may have varied in their impact through the history of the species. The multitude of influences may be reflected in our finding of a Brownian-motion-like pattern in the residual deviations from our main models. Such a pattern is expected from many quasi-independent factors shifting constraints and local optima stochastically around the main trends. Hypothetically, evolutionary lags and phylogenetic correlations would diminish if more factors were included in the main model (see Labra et al. 2009).

A particular shortcoming for a comparative study of brain-size evolution is that we have little quantitative information on the behavioral, cognitive, and sensory features that mediate selection on brain size. As just one example, vocalizations are essential in the life of most primates and their production and interpretation require complex neuronal processing. Vocalizations are affected by

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

body size (Hauser 1993) and reflect and influence a variety of social and ecological factors (Fitch and Hauser 1995). It is thus to be expected that shifts in these factors may alter selection on vocalizations with associated, but unfortunately unknown, effects on brain evolution.

The brain is a complex structure with many parts, and overall brain size is a crude measure of its capabilities. The size of individual parts will covary with overall brain size (Clark et al. 2001; Finlay et al. 2001; Falk 2007; but see Barton and Harvey 2000), and the hope is that a study of overall size may capture some similar insights as would have been captured by studies of individual parts. In a recent study, DeCasien and Higham (2019) tested how 33 different brain regions correlated with ecological and sociality traits, noting that past studies focused on only a limited number of regions (e.g. neocortex size) and that findings were inconsistent (e.g. Dunbar 1998; Schillaci 2008). Their results suggested that both dietary and social complexity were factors influencing the size of individual brain regions, which differed among primate groups. Such studies reveal more detail about brain evolution, but also have some shortcomings relative to studies of the whole brain in that they tend to be based on smaller sample sizes (between 17 and 58 species depending on the region analyzed in the DeCasien and Higham study), and the chosen phylogenetic comparative methods are problematic in our view. As we have illustrated, different results may appear from the assumption of different evolutionary models. It would be instructive to also base comparative analyses of brain parts on methods consistent with adaptive evolution.

There is debate as to whether cognitive abilities are best captured by absolute or relative brain size (Striedter 2005; Deaner et al. 2007; MacLean et al. 2014). In our study we used absolute brain size as a response variable, but we included a direct (allometric) effect of body size on its evolution. While we estimated the allometric effect of body size jointly with the adaptive effects, and indeed found that it explained the majority of the variance in brain size, our model of adaptation to ecological and dietary factors then apply to deviations from brain-body allometry, and in this sense to relative brain size. It is of course possible that absolute brain size and its cognitive

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

correlates may also adapt by inducing selection on body size, or evolve as a side effect of changes in body size driven by other means (e.g. Lande 1979).

Overall, our results contradict the idea that there is one general "rule" for the causes of primate brain-size variation – both ecology and sociality appear to have influenced the evolution of brain size, but to different degrees in different clades. Each taxon has its own evolutionary history and ecology, constraints, and selection pressures. Our findings, as well as the overall results of previous studies (e.g. Table 1), reinforce the idea that as for locomotory, dietary, and physiological adaptations, primate brains are flexible and able to adapt to the challenges of their environment.

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Funding

This work was supported by the Fulbright U.S. Scholar program to MG.

Acknowledgements:

We thank Chet Sherwood, four anonymous reviewers, associate editors Dr. Jacob Esselstyn and Dr. Lisa Barrow for comments that greatly improved the manuscript.

Disclosure Statement:

The authors have no conflicts of interest to declare.

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

References

Aiello L.C., Wheeler P. 1995. The expensive-tissue hypothesis: The brain and the digestive system in human and primate evolution. Curr. Anthropol. 36:199–221.

Aristide L., dos Reis S.F., Machado A.C., Lima I., Lopes R.T., Perez S.I. 2016. Brain shape convergence in the adaptive radiation of New World monkeys. Proc. Natl. Acad. Sci. 113:2158–2163.

Arnold C., Matthews L.J., Nunn C.L. 2010. The 10kTrees website: A new online resource for primate phylogeny. Evol. Anthropol. 19:114–118.

Barton R.A. 1996. Neocortex size and behavioural ecology in primates. Proc. R. Soc. B. 263:173–177.

Barton R. A. 1999. The evolutionary ecology of the primate brain. In: Lee P.C., editor. Comparative primate socioecology. Cambridge University Press. p. 167-194.

Barton R.A., Harvey P.H. 2000. Mosaic evolution of brain structure in mammals. Nature 405:1055–1057.

Barton R.A. 1999. The evolutionary ecology of the primate brain. In: Lee P.C., editor. Comparative Primate Socioecology. Cambridge University Press. p. 167–194.

Barton R.A., Purvis A., Harvey P.H. 1995. Evolutionary radiation of visual and olfactory brain systems in primates, bats and insectivores. Phil. Trans. Biol. Sci. 348: 381–392.

Benson-Amram S., Dantzer B., Stricker G., Swanson E.M., Holekamp K.E. 2016. Brain size predicts problem-solving ability in mammalian carnivores. Proc. Natl. Acad. Sci. 113:2532–2537.

Borries C., Gordon A.D., Koenig A. 2013. Beware of primate life history data: A plea for data standards and a repository. PLoS ONE. 8:e67200-12.

Campbell C., Fuentes A., MacKinnon K., Bearder S., Stumpf R. 2011. Primates in perspective (2nd edition). Oxford, UK: Oxford University Press.

Chivers D.J., Hladik C.M. 1980. Morphology of the gastrointestinal tract in primates: Comparisons with other mammals in relation to diet. Journal of Morphology 166: 337–386.

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Ciochon R.L., Corruccini R.S. 1975. Morphometric analysis of platyrrhine femora with taxonomic implications and notes on two fossil forms. J. Hum. Evol. 4:193–217.

Clark D.A., Mitra P.P., Wang S.S. 2001. Scalable architecture in mammalian brains. Nature 411: 189–193.

Clutton-Brock T.H., Harvey P.H. 1980. Primates, brains and ecology. J. Zool. 190:309–323.

Deaner R.O., Isler K., Burkart J., van Schaik C. 2007. Overall brain size, and not encephalization quotient, best predicts cognitive ability across non-human primates. Brain. Behav. Evol. 70:115–124.

DeCasien A.R., Higham J.P. 2019. Primate mosaic brain evolution reflects selection on sensory and cognitive specialization. Nature Ecology & Evolution. 3:1483–1493.

DeCasien A.R., Williams S.A., Higham J.P. 2017. Primate brain size is predicted by diet but not sociality. Nature Ecology & Evolution. 1:0112.

Dunbar R.I.M. 1992. Neocortex size as a constraint on group size in primates. J. Hum. Evol. 22:469–493.

Dunbar R.I.M. 1998. The social brain hypothesis. Evol. Anthrop. 6:178–190.

Dunbar R.I.M., Shultz S. 2007. Understanding primate brain evolution. Phil. Trans. R. Soc. Lond. B 362: 649–658.

Dunbar R.I.M., Shultz S. 2017. Why are there so many explanations for primate brain evolution? Phil. Trans. R. Soc. Lond. B 372: 20160244–14.

Estrada A. et al. 2018. Primates in peril: the significance of Brazil, Madagascar, Indonesia and the Democratic Republic of the Congo for global primate conservation. PeerJ. 6:e4869.

Falk D. 2007. Evolution of the Primate Brain. Handbook of Paleoanthropology. Springer Berlin Heidelberg.

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Fernandez-Duque E., Di Fiore A., Huck M. 2012. The behavior, ecology, and social evolution of New World monkey. In: Mitani J.C., Call J., Kappeler P.M., Palombit R.A., Silk J.B., editors. The evolution of Primate societies. Chicago: University of Chicago Press.

Finlay B.L., Darlington R.B., Nicastro N. 2001. Developmental structure in brain evolution. Behavioral and Brain Sciences. 24:298–304.

Fitch W.T., Hauser M.D. 1995. Vocal production in nonhuman primates: Acoustics, physiology, and functional constraints on “honest” advertisement. Am. J. Primat. 37:191–219.

Fonseca-Azevedo K., Herculano-Houzel S. 2012. Metabolic constraint imposes tradeoff between body size and number of brain neurons in human evolution. Proc. Natl. Acad. Sci. 109:18571–18576.

Ford S.M. 1980. Callitrichids as phyletic dwarfs, and the place of the callitrichidae in platyrrhini. Primates. 21:31–43.

Garamszegi L.Z. 2014. Uncertainties due to within-species variation in comparative studies: Measurement errors and statistical weights. In: Garamszegi L.Z., editor. Modern phylogenetic momparative Methods and their application in evolutionary biology. Berlin, Heidelberg: Springer. p. 157–199.

Garamszegi L.Z., Mundry R. 2014. Multimodel-inference in comparative analyses. In Garamszegi L.Z., editor. Modern phylogenetic comparative methods and their application in evolutionary biology. Berlin, Heidelberg: Springer. p. 305–331.

Gilbert C.C., Jungers W.L. 2017. Comment on relative brain size in early primates and the use of encephalization quotients in primate evolution. J. Hum. Evol. 109:79–87.

Gould S.J., Eldredge N. 1977. Punctuated equilibria: the tempo and mode of evolution reconsidered. Paleobiology:115–151.

Grabowski M., Voje K.L., Hansen T.F. 2016. Evolutionary modeling and correcting for observation error support a 3/5 brain-body allometry for primates. J. Hum. Evol. 94:106–116.

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Hansen T.F. 1997. Stabilizing selection and the comparative analysis of adaptation. Evolution. 51:1341–1351.

Hansen T.F. 2014. Use and misuse of comparative methods in the study of adaptation. In: Garamszegi L.Z., editor. Modern phylogenetic comparative methods and their application in evolutionary biology. Berlin, Heidelberg: Springer. p. 351–379.

Hansen T.F., Bartoszek K. 2012. Interpreting the evolutionary regression: the interplay between observational and biological errors in phylogenetic comparative studies. Syst. Biol. 61:413–425.

Hansen T.F., Orzack S.H. 2005. Assessing current adaptation and phylogenetic inertia as explanations of trait evolution: the need for controlled comparisons. Evolution. 59: 2063–2072.

Hansen T.F., Pienaar J., Orzack S.H. 2008. A comparative method for studying adaptation to a randomly evolving environment. Evolution. 62: 1965–1977.

Harvey P.H., Clutton-Brock T.H., Mace G.M. 1980. Brain size and ecology in small mammals and primates. Proc. Natl. Acad. Sci. 77: 4387–4389.

Harvey P.H., Krebs J.R. 1990. Comparing brains. Science. 249: 140–146.

Hauser M.D. 1993. The evolution of nonhuman primate vocalizations: Effects of phylogeny, body weight, and social context. Am. Nat. 142: 528–542.

Hershkovitz P. 1970. Notes on Tertiary platyrrhine monkeys and description of a new genus from the late Miocene of Colombia. FPR. 12: 1–37.

Hladik C.M. 1967. Surface relative du tractus digestif de quelques primates, morphologie des villosites intestinales et correlations avec le regime alimentaire. Mammalia 31: 120–147.

Hladik C.M. 1978. Adaptive strategies of primates in relation to leaf eating. In: Montgomery G.G., editor. The ecology of arboreal folivores. Smithsonian Institution Press, Washington. Pp. 373–395.

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Houle D., Pélabon C., Wagner G.P., Hansen T.F. 2011. Measurement and meaning in biology. Quart. Rev. Biol. 86: 3–34.

Isler K., Christopher Kirk E., Miller J.M.A., Albrecht G.A., Gelvin B.R., Martin R.D. 2008. Endocranial volumes of primate species: scaling analyses using a comprehensive and reliable data set. J. Hum. Evol. 55: 967–978.

Isler K., Van Schaik C.P. 2009. The expensive brain: a framework for explaining evolutionary changes in brain size. J. Hum. Evol. 57: 392–400.

Jerison H.J. 1979. Brain, body and encephalization in early primates. J. Hum. Evol. 8: 615–635.

Joffe T.H., Dunbar R.I. 1997. Visual and socio-cognitive information processing in primate brain evolution. Proc. R. Soc. B. 264: 1303–1307.

Kappeler P.M. 2012. The behavioral ecology of strepsirrhines and tarsiers. In: Mitani J.C., Call J., Kappeler P.M., Palombit R.A., Silk J.B., editors. The evolution of primate societies. Chicago: University of Chicago Press. p. 17–42.

Kappeler P.M., Heymann E.W. 1996. Nonconvergence in the evolution of primate life history and socio-ecology. Biol. J. Linn. Soc. 59: 297–326.

Kopperud B.T., Pienaar J., Voje K.L., Orzack S.H., Hansen T.F. 2020. Slouch: Stochastic Linear Ornstein-Uhlenbeck Comparative Hypotheses. R package version 2.1.4.

Kudo H., Dunbar R.I.M. 2001. Neocortex size and social network size in primates. Animal Behaviour 62: 711–722.

Kuzawa C.W., Chugani H.T., Grossman L.I., Lipovich L., Muzik O., Hof P.R., Wildman D.E., Sherwood C.C., Leonard W.R., Lange N. 2014. Metabolic costs and evolutionary implications of human brain development. Proc. Natl. Acad. Sci. 111:13010–13015.

Labra A., Pienaar J., Hansen T.F. 2009. Evolution of thermal physiology in *Liolaemus* lizards: adaptation, phylogenetic inertia, and niche tracking. Am. Nat. 174:204–220.

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Lande, R. 1979. Quantitative genetic analysis of multivariate evolution, applied to brain:body size allometry. Evolution 33: 402-416.

Lindenfors P. 2005. Neocortex evolution in primates: the “social brain” is for females. Biol. Lett. 1: 407–410.

MacLean E.L., Barrickman N.L., Johnson E.M., Wall C.E. 2009. Sociality, ecology, and relative brain size in lemurs. J. Hum. Evol. 56: 471–478.

MacLean E.L. et al. 2014. The evolution of self-control. Proc. Natl. Acad. Sci. 111: E2140–E2148.

Martin R.D. 1992. Goeldi and the dwarfs: the evolutionary biology of the small New World monkeys. J. Hum. Evol. 22: 367–393.

Martin R.D. 1996. Scaling of the mammalian brain: the maternal energy hypothesis. News in Physiological Sciences. 11: 149–156.

Miller I.F., Barton R.A., Nunn C.L. 2019. Quantitative uniqueness of human brain evolution revealed through phylogenetic comparative analysis. eLife. 8:317.

Mittermeier R.A., Wilson D.E. 2013. Handbook of the mammals of the world: Vol. 3: Primates.

Moen D.S., Cabrera-Guzman, E., Caviedes-Solis I.W., Gonzalez-Bernal E., Hanna A.R. 2022. Phylogenetic analysis of adaptation in comparative physiology and biomechanics: overview and a case study of thermal physiology in treefrogs. J. Exp. Biol. 225: jeb243292.

Morrissey M.B. 2016. Meta-analysis of magnitudes, differences, and variation in evolutionary parameters. J. Evol. Biol. 29:1882–1904.

Nunn C.L., Van Schaik C.P. 2002. A comparative approach to reconstructing the socioecology of extinct primates. In: Plavcan J.M., Kay R.F., Jungers W.L., Van Schaik C.P., editors. Reconstructing behavior in the primate fossil record. Boston, MA: Springer. p. 159–215.

Perez-Barberia, F.J., Shultz S., Dunbar, R.I.M. 2007. Evidence for coevolution of sociality and relative brain size in three orders of mammals. Evolution 61: 2811-2821.

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Plavcan J.M. 1999. Mating systems, intrasexual competition and sexual dimorphism in primates. In: Lee P.C., editor. Comparative primate socioecology. Cambridge University Press. p. 241–260.

Powell L.E., Barton R.A., Street S.E. 2019. Maternal investment, life histories and the evolution of brain structure in primates. Proc. R. Soc. B. 286:20191608–9.

Powell L.E., Isler K., Barton R.A. 2017. Re-evaluating the link between brain size and behavioural ecology in primates. Proc. R. Soc. B. 284: 20171765–8.

Reader S.M., Laland K.N. 2002. Social intelligence, innovation, and enhanced brain size in primates. Proc. Natl. Acad. Sci. 99: 4436–4441.

Revell L.J. 2011. phytools: an R package for phylogenetic comparative biology (and other things). Meth. Ecol. Evol. 3: 217–223.

Sailer L.D., Gaulin S.J.C., Boster J.S., Kurland J.A. 1985. Measuring the relationship between dietary quality and body size in primates. Primates. 26: 14–27.

Sandel A.A., Miller J.A., Mitani J.C., Nunn C.L., Patterson S.K., Garamszegi L.Z. 2016. Assessing sources of error in comparative analyses of primate behavior: Intraspecific variation in group size and the social brain hypothesis. J. Hum. Evol. 94: 126–133.

Santini L., Rojas D., Donati G. 2015. Evolving through day and night: origin and diversification of activity pattern in modern primates. Behav. Ecol. 26: 789–796.

Schillaci M.A. 2006. Sexual selection and the evolution of brain size in primates. PLoS ONE. 1: e62-5.

Schillaci M.A. 2008. Primate mating systems and the evolution of neocortex size. J. Mammalogy 89: 58–63.

Shultz S., Dunbar R.I.M. 2007. The evolution of the social brain: anthropoid primates contrast with other vertebrates. Proc. R. Soc. B. 274: 2429–2436.

Shultz S., Opie C., Atkinson Q.D. 2011. Stepwise evolution of stable sociality in primates. Nature. 479:219–222.

Smith R.J., Jungers W.L. 1997. Body mass in comparative primatology. J. Hum. Evol. 32: 523–559.

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Soligo C., Martin R.D. 2006. Adaptive origins of primates revisited. J. Hum. Evol. 50: 414–430.

Street S.E., Navarrete A.F., Reader S.M., Laland K.N. 2017. Coevolution of cultural intelligence, extended life history, sociality, and brain size in primates. Proc. Natl. Acad. Sci. 114:7908–7914.

Striedter G.F. 2005. Principles of brain evolution. Sunderland (MA): Sinauer.

Thorén S., Lindenfors P., Kappeler P.M. 2006. Phylogenetic analyses of dimorphism in primates: Evidence for stronger selection on canine size than on body size. Am. J. Phys. Anthropol. 130: 50–59.

Todorov O.S., Weisbecker V., Gilissen E., Zilles K., de Sousa A.A. 2019. Primate hippocampus size and organization are predicted by sociality but not diet. Proc. R. Soc. B. 286: 20191712.

Tsuboi M., van der Bijl W., Kopperud B.T., Erritzøe J., Voje K.L., Kotrschal A., Yopak K.E., Collin S.P., Iwaniuk A.N., Kolm N. 2018. Breakdown of brain–body allometry and the encephalization of birds and mammals. Nature Ecology & Evolution:1–12.

Uyeda J.C., Harmon L.J. 2014. A novel Bayesian method for inferring and interpreting the dynamics of adaptive landscapes from phylogenetic comparative data. Syst. Biol. 63: 902–918.

Uyeda J.C., Pennell M.W., Miller E.T., Maia R., McClain C.R. 2017. The evolution of Eenergetic scaling across the vertebrate tree of life. Am. Nat. 190: 185–199.

Uyeda J.C., Zenil-Ferguson R., Pennell M.W. 2018. Rethinking phylogenetic comparative methods. Syst. Biol.. 106:13410–19.

Voje K.L., Hansen T.F., Egset C.K., Bolstad G.H., Pélabon C. 2014. Allometric constraints and the evolution of allometry. Evolution. 68: 866–885.

Whiting B.A., Barton R.A. 2003. The evolution of the cortico-cerebellar complex in primates: anatomical connections predict patterns of correlated evolution. J. Hum. Evol. 44: 3–10.

Wrangham R.W., Jones J.H., Laden G., Pilbeam D., Conklin-Brittain N. 1999. The raw and the stolen: Cooking and the ecology of human origins. Curr. Anthro. 40: 567–594.

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Figure Captions

Figure 1: Time-calibrated primate phylogeny showing regimes for best fitting bayou model of log ECV (cc) on log Body Mass (g). Regime shifts are shifts in the intercept of the allometric equation with a posterior probability greater than 0.5. Diameter of circles are proportional to the posterior probability of a shift. The tree height is 73 *myr*. Inset shows estimated optima for the two regime shifts with the black dot and the box indicating the median and the interquartile range of the posterior distribution. Number of species in that clade are shown in parenthesis.

Figure 2: Brain size as a function of ecological factors. Mean brain size of each species is plotted as a function of its current ecological or social factors for a) all primates; b) strepsirrhines plus tarsiers; c) haplorhines minus tarsiers; and d) platyrrhines. In a), b) and c) we show the predicted brain sizes from the best-fitting evolutionary trend model if the species had evolved in the same regime from the root of the phylogeny assuming for illustration that they all started from the global mean brain and body mass. For platyrrhines the best model was an Ornstein-Uhlenbeck model with fixed optima and not a trend model. Hence, in d) we show the estimated primary optima for the regimes. In all cases the evolutionary model included body size and the plotted species mean brain sizes are residuals from the allometric model. Note that the species values are often systematically different from the evolutionary predictions. This is because the species may not have evolved in their current selective regime for a long time, and thus display a lag in adaptation. The error bars are standard errors, conditional on the maximum likelihood estimates for α and σ_y^2. Abbreviations are Folivorous Spatial Polygyny (FSP), Folivorous Monogamy (FM), Folivorous Polygynandry (FPG), Folivorous Harem Polygyny (FHP), Non-Folivorous Spatial Polygyny (NFSP), Non-Folivorous Monogamy (NFM), Non-Folivorous Polygyny, (NFP), Non-Folivorous Polygynandry (NFPG), Non-Folivorous Harem Polygyny (NFHP) in a), b) and c) with subsets in b) along with a nocturnal (N) or diurnal (D) activity period. Folivores are squares, non-folivores are circles.

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Table 1: Studies testing the role of Ecological and/or Sociality metrics as an explanation for primate brain evolution

Study	Focus	Broad Hypotheses Tested	Factors Found to Affect Brain Size	N	Statistical Approach
Clutton-Brock and Harvey 1980	Brain	Ecology, Sociality	Diet, Home-Range Size	<= 30 species	OLS and MA
Dunbar 1992	Neocortex	Ecology, Sociality	Group Size	38 genera	RMA
Barton 1996	Neocortex	Ecology, Sociality	Group Size, Diet	43 species	RMA
Joffe and Dunbar 1997	Primary visual cortex	Sociality	Social Group Size	27 species	RMA
Kudo and Dunbar 2001	Neocortex	Sociality	Social Partners (Grooming Clique Size), Total Group Size	31 species	RMA
Reader and Laland 2002	Executive brain ratio, volume, residual*	Sociality	Social Learning, Innovation, Tool Use	29-30 Species	RMA
Lindenfors 2005	Neocortex	Sociality	Female Group Size	20 species	RMA
Schillaci 2006	Brain	Sociality	Mating System, Body Mass Dimorphism	31 species	RMA
Shultz and Dunbar 2007	Brain	Diet, Sociality	Diet, Group Size, Social System, Stratification, Diurnal vs. Nocturnal	45 species	PGLS with BM
Dunbar and Shultz 2007	Brain	Ecology, Behavior, Life-History	Diet, Home-Range Size, Day Range Size, Group Size	24 species	OLS
Pérez-Barberia et al. 2007	Brain	Sociality	Social Group Associations, Group Size	42 species	PGLS with BM
Maclean et al. 2009	Brain	Ecology, Sociality	Diet and Activity Pattern	19 lemur species	RMA and Independent Contrasts
Sandel et al. 2016	Neocortex	Sociality	Group Size	23 species	PGLS with BM

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

DeCasien et al. 2017	Brain	Ecology, Sociality	Diet	144 species	PGLS with BM
Street et al. 2017	Brain	Sociality, Life History	Longevity, Social Learning, Social Group Size	186 species	Bayesian Mixed Models
Powell et al. 2017	Brain	Ecology, Sociality	Home Range Size, Activity Period, Diet	99 species	PGLS with BM
Todorov et al. 2019	Hippocampus	Ecology, Sociality	Group size, Home Range Size	43 species	PGLS with BM
Current study	Brain	Ecology, Sociality	Diet, Social/Mating System, Group Size	128 species	PGLS with OU

*All metrics focused on regions that function in innovation and social learning - neocortex and striatum

Key: OLS - ordinary least squares regression, equivalent to a model of instant adaptation, MA - major axis regression, RMA - reduced major axis regression, PGLS - phylogenetic generalized least squares, BM - Brownian-motion, OU - Ornstein-Uhlenbeck

Table focuses on external selective factors driving brain evolution

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Table 2: *Slouch* results for all primates showing the best model, its parameter estimates and type, and the allometric model (i.e. only log body mass as a predictor)[a,b]. In each case, log endocranial volume is the response variable and log body mass is included as a direct effect predictor.

Group	*N*	Predictors	Parameters	Regime Trends	Direct Effect	Evolutionary					
				Intercepts	Slope	Slope	$t_{1/2}$	V_y	R^2 (%)	Res. R^2 (%)	*ΔAICc*
All Primates	12 8	Diet (Strict Folivorous/Non-folivorous)+Group Size+Mating System	Folivorous Harem Polygyny	-2.87±1.72			2.63 [0.43-∞]	0.18 [0.04-0.29]	87		0
			Folivorous Monogamy	-1.7±1.79							
			Folivorous Polygynandry	0.41±2.19							
			Folivorous Spatial Polygyny	-3.17±2.05							
			Non-folivorous Harem Polygyny	4.19±2.37							
			Non-folivorous Monogamy	-3.29±1.55							
			Non-folivorous Polyandry	-7.77±1.99							
			Non-folivorous Polygynandry	3.25±1.01							
			Non-folivorous Spatial Polygyny	-1.32±0.2							
			log Body Mass		0.56±0.02						
			log Average Group Size			-0.04±0.02					
All	12	Body Mass	Intercept	-1.3±0.24			∞ [0.86-∞]	0.35 [0.12-	80.0	0	21.5

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Primates	8			0.45]	7
		log Body Mass	0.59±0.03		

a Results include sample size (*N*), estimated parameters, the phylogenetic half-life ($t_{1/2}$) in units of tree height with 2-unit support interval shown in brackets, the stationary variance (v_y) in units of squared trait units (log ECV in *cc*) per unit tree height with 2-unit support interval shown in brackets, the variance explained by the model (R^2) shown in %, the extra variance explained by the model after accounting for the variance explained by body mass (Res. R^2), and the difference between the *AICc* score of the best model and the Brownian-motion model with body mass as the sole predictor (*ΔAICc*).

b Estimated parameters include Regime Optima or Trends for the effects of the categorical predictors - the former from an OU model [in trait units (log ECV in *cc*)], the latter from an OU model as half-life goes to infinity [in units of trait units (log ECV in *cc*) per tree height)], the Direct Effect Slope for the relationship between brain and body mass [in units of trait units (log ECV in *cc*) per change in log Body Mass], and the Evolutionary Slope for the relationship between brain mass and group size [in units of trait units (log ECV in *cc* per change in log Group Size].

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Table 3: *Slouch* results for strepsirrhines showing the best model, its parameter estimates and type, and the allometric model (i.e. only log body mass as a predictor)[a,b]. In each case, log endocranial volume is the response variable and log body mass is included as a direct effect predictor.

Group	*N*	Predictors	Parameters	Regime Trends/ Intercepts	Direct Effect Slope	Evolutionary Slope	$t_{1/2}$	V_y	R^2 (%)	Res. R^2 (%)	*ΔAICc*
Strepsirrhines	28	Group Size + Mating System + Activity Period	Monogamy Diurnal	0.14±0.49			∞ [0.43-∞]	0.1 [0.02-0.22]	94.5	64.5	0
			Monogamy Nocturnal	-1.72±0.48							
			Polygynandry Diurnal	1.42±0.77							
			Spatial Polygyny Nocturnal	-1.26±0.23							
			log Body Mass		0.56±0.03						
			log Average Group Size			-0.53±0.11					
Strepsirrhines	28	Body Mass	Intercept	-1.47±0.35			0.18 [0.06-∞]	0.06 [0.04-0.43]	84.5	0	13.6
			log Body Mass		0.59±0.05						

a Results include sample size (*N*), estimated parameters, the phylogenetic half-life ($t_{1/2}$) in units of tree height with 2-unit support interval shown in brackets, the stationary variance (v_y) in units of squared trait units (log ECV in *cc*) per unit tree height with 2-unit support interval shown in brackets, the variance explained by the model (R^2) shown in %, the extra variance explained by the model after accounting for the variance explained by body mass (Res. R^2), and the Brownian-motion model with body mass as the sole predictor (*ΔAICc*).

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

b Estimated parameters include Regime Optima or Trends for the effects of the categorical predictors - the former from an OU model [in trait units (log ECV in *cc*)], the latter from an OU model as half-life goes to infinity [in units of trait units (log ECV in *cc*) per tree height)], the Direct Effect Slope for the relationship between brain and body mass [in units of trait units (log ECV in *cc*) per change in log Body Mass], and the Evolutionary Slope for the relationship between brain mass and group size [in units of trait units (log ECV in *cc* per change in log Group Size].

c Plus tarsiers

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Table 4: *Slouch* results for haplorhines showing the best model, its parameter estimates and type, and the allometric model (i.e. only log body mass as a predictor)[a,b]. In each case, log endocranial volume is the response variable and log body mass is included as a direct effect predictor.

Group	*N*	Predictors	Parameters	Regime Trends	Direct Effect	Evolutionary					
				Intercepts	Slope	Slope	$t_{1/2}$	V_y	R^2 (%)	Res. R^2 (%)	*ΔAICc*
Haplorhines	100	Diet (Strict Folivorous/Non-folivorous) + Mating System	Folivorous Harem Polygyny	-6.77±1.75			∞ [0.43-∞]	0.2 [0.04-0.29]	84.2	32.2	0
			Folivorous Polygynandry	-7.17±3							
			Non-folivorous Harem Polygyny	0.86±2.37							
			Non-folivorous Monogamy	-7.1±1.8							
			Non-folivorous Polyandry	-13.28±2.18							
			Non-folivorous Polygynandry	-0.02±0.29							
			Non-folivorous Spatial Polygyny	-1.21±3.43							
			log Body Mass		0.52±0.03						
Haplorhines	100	Body Mass	Intercept	-0.74±0.31			∞ [0.67-∞]	0.33 [0.08-0.43]	76.6	0	22.9
			log Body Mass		0.58±0.03						

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

a Results include sample size (*N*), estimated parameters, the phylogenetic half-life ($t_{1/2}$) in units of tree height with 2-unit support interval shown in brackets, the stationary variance (v_y) in units of squared trait units (log ECV in *cc*) per unit tree height with 2-unit support interval shown in brackets, the variance explained by the model (R^2) shown in %, the extra variance explained by the model after accounting for the variance explained by body mass (Res. R^2), and the Brownian-motion model with body mass as the sole predictor (*ΔAICc*).

b Estimated parameters include Regime Optima or Trends for the effects of the categorical predictors - the former from an OU model [in trait units (log ECV in *cc*)], the latter from an OU model as half-life goes to infinity [in units of trait units (log ECV in *cc*) per tree height)] and the Direct Effect Slope for the relationship between brain and body mass [in units of trait units (log ECV in *cc*) per change in log Body Mass].

c Minus tarsiers

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Table 5: *Slouch* results for platyrrhines showing the best model, its parameter estimates and type, and the allometric model (i.e. only log body mass as a predictor)[a,b]. In each case, log endocranial volume is the response variable and log body mass is included as a direct effect predictor.

Group	*N*	Predictors	Parameters	Regime Optima	Direct Effect	Evolutionary					
				Intercepts	Slope	Slope	$t_{1/2}$	V_y	R^2 (%)	Res. R^2 (%)	*ΔAICc*
Platyrrhines	32	Diet (Strict Folivorous/Non-folivorous) + Mating System	Folivorous Harem Polygyny	-1.53±0.3			0.06 [0.06-0.92]	0.01 [0.01-0.05]	98.3	92.5	0
			Folivorous Polygynandry	-0.79±0.45							
			Non-folivorous Harem Polygyny	-0.15±0.35							
			Non-folivorous Monogamy	-1.57±0.21							
			Non-folivorous Polyandry	-1.66±0.21							
			Non-folivorous Polygynandry	-0.95±0.27							
			log Body Mass		0.62±0.03						
Platyrrhines	32	Body Mass	Intercept	-1.84±0.51			∞ [0.18-∞]	0.53 [0.06-0.96]	77.6	0	18.9
			log Body Mass		0.71±0.07						

a Results include sample size (*N*), estimated parameters, the phylogenetic half-life ($t_{1/2}$) in units of tree height with 2-unit support interval shown in brackets, the stationary variance (v_y) in units of squared trait units (log ECV in *cc*) per unit tree height with 2-unit support interval shown in brackets, the

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022


variance explained by the model (R^2) shown in %, the extra variance explained by the model after accounting for the variance explained by body mass (Res. R^2), and the difference between the *AICc* score of the best model and the Brownian-motion model with body mass as the sole predictor (*ΔAICc*).

b Estimated parameters include Regime Optima or Trends for the effects of the categorical predictors - the former from an OU model [in trait units (log ECV in *cc*)], the latter from an OU model as half-life goes to infinity [in units of trait units (log ECV in *cc*) per tree height)], and the Direct Effect Slope for the relationship between brain and body mass [in units of trait units (log ECV in *cc*) per change in log Body Mass].

Accepted Manuscript

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022


Fig. 1



Accepted

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022

Fig. 2



Accepted

Downloaded from https://academic.oup.com/sysbio/advance-article/doi/10.1093/sysbio/syac075/6862053 by University of Arizona user on 18 December 2022